I is the presence or absence of significant intrinsic sympathomimetic activity, and S is the presence or absence of significant cardioselectivity. Similarly, α is a constant, δ describes the overall treatment effect, β describes the effect of intrinsic sympathomimetic activity, and γ describes the effect of cardioselectivity.

- Smith J, Channer KS. Increasing prescription of drugs for secondary prevention after myocardial infarction. *BMJ* 1995;311:917-8.
- Eccles M, Bradshaw C. Use of secondary prophylaxis against myocardial infarction in the north of England. *BMJ* 1991;302:91-2.
- Viskin S, Barron HV. β -Blockers prevent cardiac death following myocardial infarction: so why are so many infarct survivors discharged without β -blockers? *Am J Cardiol* 1996;78:821-2.
- Yusuf S, Peto R, Lewis J, Collins R, Sleight P. β -Blockade during and after myocardial infarction: an overview of the randomized trials. *Prog Cardiovasc Dis* 1985;27:335-71.
- Smith TC, Spiegelhalter DJ, Thomas A. Bayesian approaches to random-effects meta analysis: a comparative study. *Stats Med* 1995;14:2685-99.
- Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias: dimensions of methodological quality associated with estimates of treatment effects in controlled trial. *JAMA* 1995;273:408-12.
- Rothman KJ. *Modern epidemiology*. Boston: MA Little and Brown, 1986.
- Robins J, Breslow N, Greenland S. Estimators of the Mantel-Haenszel variance consistent in both sparse data and large strata models. *Biometrics* 1986;42:311-23.
- Fleiss J, Gross AJ. Meta-analysis in epidemiology, with special reference to studies of the association between exposure to environmental tobacco smoke and lung cancer: a critique. *J Clin Epidemiol* 1991;44:127-39.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clin Trials* 1986;7:177-88.
- Carlin JB. Meta analysis for 2 $\times$ 2 tables: a Bayesian approach. *Stats Med* 1992;11:141-58.
- Hardy RJ, Thompson SG. A likelihood approach to meta analysis with random effects. *Stats Med* 1996;15:619-29.
- Spiegelhalter D, Thomas A, Best N, Gilks W. BUGS: Bayesian inference using Gibbs sampling, Version 0.50. Cambridge: Medical Research Council Biostatistics Unit, 1995.
- Ioannidis JPA, Cappelleri JC, Lau J, Skolnik PR, Melville B, Chalmers TC, et al. Early or deferred zidovudine therapy in HIV-infected patients without an AIDS defining illness. *Ann Intern Med* 1995;122:856-66.
- Freemantle N, Mason JM, Eccles M. Deriving treatment recommendations from evidence within randomised trials: the role and limitation of meta analysis. *Intern J Technol Assess Health Care* (in press).
- Second International Study of Infarct Survival Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both or neither among 17 187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* 1988;ii:349-60.
- The Thrombolysis in Myocardial Infarction Study Group. Comparison of invasive and conservative strategies after treatment with intravenous tissue plasminogen activator in acute myocardial infarction: results of the thrombolysis in myocardial infarction (TIMI) phase II trial. *N Engl J Med* 1989;320:618-27.
- Wilhelmsson C, Vedin JA, Wilhelmsson L, Tibblin G. Reduction of sudden deaths after myocardial infarction by treatment with alprenolol: preliminary results. *Lancet* 1974;ii:1157-60.
- Feely J, de Vane PJ, Maclean D. β -Blockers and sympathomimetics. *BMJ* 1983;286:1043-7.
- McDevitt DG. The assessment of β -adrenoceptor-blocking drugs in man. *Br J Clin Pharmacol* 1977;4:413-25.
- Owen A. Intravenous β -blockade in acute myocardial infarction: should be used in combination with thrombolysis. *BMJ* 1998;317:226-7.
- Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy. I. Prevention of death, myocardial infarction and stroke by prolonged antiplatelet therapy in various categories of patients. *BMJ* 1994;308:81-106.
- β -Blocker Pooling Project Research Group. The β -blocker pooling project (BBPP): subgroup findings from randomized trials in post infarction patients. *Eur Heart J* 1988;9:8-16.
- Gottlieb SS, McCarter RJ, Vogel RA. Effect of β -blockade on mortality among high risk and low risk patients after myocardial infarction. *N Engl J Med* 1998;339:489-97.
- First International Study of Infarct Survival Collaborative Group. Randomised trial of intravenous atenolol among 16 027 cases of suspected acute myocardial infarction: ISIS-1. *Lancet* 1986;ii:57-67.
- Doughty RN, Rodgers A, Sharpe N, MacMahon S. Effects of β -blocker therapy on mortality in patients with heart failure: a systematic overview of randomized controlled trials. *Eur Heart J* 1997;18:560-5.
- Heidenreich PA, Lee TT, Massie BM. Effect of β -blockade on mortality in patients with heart failure: a meta analysis of randomized clinical trials. *J Am Coll Cardiol* 1997;30:27-34.
- Cleland JGF, Freemantle N, McGowan J, Clark A. The evidence for β blockers in heart failure. *BMJ* 1999;318:824-5.
- Latest trials on statins show large benefits? *Lancet* 1997;350:1525.
- The Multicenter Diltiazem Postinfarction Trial (MDPIT) Research Group. The effect of diltiazem on mortality and reinfarction after myocardial infarction. *N Engl J Med* 1988;319:385-92.
- Smith P, Arnesen H, Holme I. The effect of warfarin on mortality and reinfarction after myocardial infarction. *N Engl J Med* 1990;323:147-52.
- Anticoagulants in the Secondary Prevention of Events in Coronary Thrombosis (ASPECT) Research Group. Effect of long term oral anticoagulant treatment on mortality and cardiovascular morbidity after myocardial infarction. *Lancet* 1994;343:499-503.

(Accepted 6 April 1999)

Risk factors for human hantavirus infection: Franco-Belgian collaborative case-control study during 1995-6 epidemic

N S Crowcroft, A Infuso, D Illef, B Le Guenno, J-C Desenclos, F Van Loock, J Clement

Puumala hantavirus is the most common human hantavirus infection in Europe.^{1,2} It is transmitted to humans by inhalation or contamination of skin breaches by urine or faeces of infected bank voles. Infection ranges from subclinical to a severe influenza-like illness progressing to acute renal failure.³ We carried out a case-control study in an endemic area in France and Belgium to estimate knowledge of hantavirus and identify possible risk factors for infection.

Subjects, methods, and results

National reference laboratories in each country identified cases for the study. A case was defined as someone with laboratory confirmed IgM positive *Puumala hantavirus* infection between 1 April 1996 and 31 July 1996 in the French departments Nord, Ardennes, and Aisne and Belgian provinces of Hainaut, Namur, and Luxembourg. Controls were matched by sex, community (village), and age group. They were randomly selected from the telephone book. Interviews were carried out by telephone using a standardised question-

naire covering knowledge of hantavirus, distance of the home to a forest, refuse disposal, rodent infestation and control, gardening activities, use of wood for heating or cooking, activities in forests, and entry into rodent infested buildings.

In all, 69/88 (78%) eligible cases were included in the study and 125 controls were recruited. Most cases were in men (51) and those aged 15-65 years (64). Two cases and one control were forestry workers—no others were in occupations thought to be at risk. Forty seven per cent (91/194) of those interviewed had heard of hantavirus infection before becoming ill or being interviewed. Friends were the commonest source of information (44/91, 48%); 63/75 (84%) had heard of the disease in the past 3 years.

The table shows the results of logistic regression. Cases and controls often went walking in forests (odds ratio 0.5, 95% confidence interval 0.1 to 2.7; $P = 0.64$). Cases were more likely to have entered a building where there might be rodents (1.9, 1.0 to 3.6; $P = 0.05$) and were more likely to have cleaned (4.2, 1.1 to 15.7;

Epidemiology Department, Scientific Institute of Public Health—Louis Pasteur, Brussels, Belgium

N S Crowcroft, fellow, European programme for intervention epidemiology training
J-C Desenclos, head of infectious diseases unit

Réseau National de Santé Publique, Saint-Maurice, France

A Infuso, fellow, European programme for intervention epidemiology training
F Van Loock, epidemiologist

continued over

BMJ 1999;318:1737-8

Cellule
Inter-régionale
d'Epidémiologie
d'Intervention,
Direction Régionale
des Affaires
Sanitaires et
Sociales du Nord
Pas-de-Calais, Lille,
France

D Illef,
*regional
epidemiologist*

Centre National de
Référence pour les
Fièvres
Hémorragiques et
les Arbovirus,
Institut Pasteur,
Paris, France

B Le Guenno,
director

Belgian Zoonosis
Workgroup,
National Reference
Center for
Hantavirus
Infections, Queen
Astrid Military
Hospital, B-1120,
Brussels, Belgium

J Clement,
head

Correspondence to:
Dr N S Crowcroft,
Immunisation
Division, Public
Health Laboratory
Service,
Communicable
Disease
Surveillance Centre,
London NW9 5EQ
ncrowcro@ogks,
co.uk

Conditional logistic regression model for infection with hantavirus. Adjusted odds ratios for activities undertaken by cases and controls

Exposure	Unadjusted odds ratio (95% CI)	Adjusted odds ratio (95% CI)	P value
Significant exposure in a forest (spent >16 hours in a forest; fetched, picked up, carried, or worked with wood; or exposed to dust or earth in a forest)	3.1 (1.6 to 6.0)	6.1 (1.9 to 19.5)	0.003
Interaction term between living <50 m from forest and seeing rodents	—	19.4 (1.2 to 308.2)	0.04
Lives <50 m from forest	3.5 (1.5 to 7.9)	1.9 (0.4 to 9.2)	0.43
Saw rodents at home	2.2 (1.1 to 4.3)	1.8 (0.5 to 6.1)	0.34
Entered a building where there may be rodents	2.7 (1.4 to 5.0)	3.0 (1.0 to 9.1)	0.046
Carries out rodent control at home	0.7 (0.3 to 1.4)	0.3 (0.1 to 1.1)	0.06
Home is cleaned more than once a week	2.9 (1.1 to 7.3)	3.8 (0.9 to 16.3)	0.07
Had been digging earth	3.6 (1.5 to 8.9)	3.1 (0.8 to 11.8)	0.09

P=0.04 and raised dust there (15.7, 2.4 to 651; P=0.01.) Two variables were constructed to refine the logistic regression analysis. For forests the variable was defined by those who spent more than 16 hours a month in forests, who went to forests for wood, or who picked up wood or were exposed to dust or earth during a leisure visit. For exposure in buildings where they may have been rodents the variable was defined by those who spent more than 2 hours there and who cleaned, raised dust, or made a vigorous physical effort. In the final model of the conditional logistic regression analysis, cases were more likely to live less than 50 metres from a forest and have seen rodents in or around their home, to have been digging, to have spent long periods in forests and been in contact with wood or disturbed earth or dust (table). Rodent control was more common among controls. Cases were more likely than controls to both live near a forest and see rodents at home (66.1, 5.7 to 768.9).

Comment

We did not test controls to ensure that they had never been infected subclinically, but in a previous case-control study in Belgium all 69 controls were seronegative⁴ and the general population of the French Ardennes has a seroprevalence of only 0.45%.⁵ The interaction between living near a forest and seeing rodents at home has not been previously reported—bank voles are thought to prefer empty buildings.

Rodent control at home was protective. This simple, cheap measure can be recommended to those living near forest in an endemic area.

We thank all the participating clinicians for their contribution to the study.

Contributors: NSC, AI, DI, and BleG contributed substantially to conception and design, analysis and interpretation, drafts of the article, and final approval of the version to be published. NSC led the writing of the paper and acts as guarantor for the study. J-CD contributed substantially in supervising the project at design stage, discussion of results and in revising the manuscript. FVL had the idea to do a study, supervised NSC during the study, and carried out some telephone interviews. JC provided his support for the study and the reference laboratory results for Belgium. Dr D Kadi, Dr Van Hoof, Mrs I Thomas, and Ms S Ghoos assisted with telephone interviews.

Funding: AI and NSC were fellows in the European programme for intervention epidemiology training funded by the European Community under agreement number SOC 96 202584 05GF01.

Competing interests: None declared.

- 1 Le Guenno B, Camprasse MA, Guilbaut JC, Lanoux P, Hoen B. Hantavirus epidemic in Europe, 1993. *Lancet* 1994;343:114-5.
- 2 Clement J, Heyman P, McKenna P, Colson P, Avsic-Zupanc T. The hantaviruses of Europe: from the bedside to the bench. *Emerging Infect Dis* 1997;3:205-11.
- 3 Le Guenno B, Coudrier D, Camprasse MA. Les hantavirus: aspects virologiques et diagnostiques. Données sur l'épidémiologie de la fièvre hémorragique avec syndrome rénal en France. *Méd Mal Infect* 1994;24 (spécial):512-6.
- 4 Van Loock F, Thomas I, Clement J, Ghoos S, Colson P. A case-control study after a hantavirus outbreak in the south of Belgium: who is at risk? *Clin Infect Dis* (in press).
- 5 Le Guenno B. Les hantavirus. *Méd Mal Infect* 1997;27:703-10.

(Accepted 18 December 1998)

Improvement in clinical work through feedback: intervention study

Rolf Jorde, Arne Nordøy

Department of
Internal Medicine,
University Hospital
of Tromsø, 9038
Tromsø, Norway
Rolf Jorde,
professor
Arne Nordøy,
professor
continued over

We have frequently found an unacceptable number of hospital records and discharge reports lacking even the most basic information. To improve on this, we reviewed our hospital records and discharge reports on a regular basis, and we sent out questionnaires on quality of care to patients shortly after discharge. The hospital's medical staff received the results as a summarised report every other week. We deliberately disclosed only half of the variables studied. At the end of one year, the results were compared with those obtained before the intervention.

Subjects, methods, and results

Our study took place from September 1994 to October 1995 at the Department of Internal Medicine, University Hospital of Tromsø (120 beds and 45 doctors). We reviewed the hospital records for two sets of information: variables that were disclosed to the staff (past or present occupation, smoking habits, general physical condition, and blood pressure) and variables that were not disclosed (marital status, alcohol consumption, glandular enlargements, and