- Jameson A, Paris C, Tasso C, eds. *Proceedings of the sixth international conference on user modelling*. New York: Springer Wien, 1997:107-18.
- 10 Kreuter MW, Strecher VJ. Do tailored behavior change messages enhance the effectiveness of health risk appraisal? Results from a randomized trial. *Health Educ Res* 1996;11:97-105.
  - 11 Department of Health. *NHS information for health*. London: DoH, 1998.
  - 12 Hathaway D. Effect of preoperative instruction on postoperative outcomes: a meta-analysis. *Nurs Res* 1986;35:269-75.
  - 13 Schwartz LP, Brenner ZR. Critical care unit transfer: reducing patient stress through nursing interventions. *Heart Lung* 1979;8:540-7.
  - 14 Gilhooly MLM, McGhee SM. Medical records: practicalities and principles of patient possession. *J Med Ethics* 1991;17:138-43.
  - 15 Hinds C, Streater A, Mood D. Functions and preferred methods of receiving information related to radiotherapy. *Cancer Nurs* 1995;18:374-83.
  - 16 Jones RB, Pearson J, McGregor S, Cawsey A, Barrett A, Gilmour H, et al. Cross sectional survey of patients' satisfaction with information about cancer. *BMJ* 1999;319:1247-8.
  - 17 Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-70.
  - 18 Watson M, Greer S, Young J, Inayat Q, Burgess C, Robertson B. Development of a questionnaire measure of adjustment to cancer: the MAC scale. *Psychol Med* 1988;18:203-9.
  - 19 Jones RB, Knill-Jones RP. Electronic patient record project: direct patient input to the record. Report for the Strategy Division of the Information Management Group of the NHS ME Glasgow: University of Glasgow, 1994.
  - 20 Glimelius B, Birgegard G, Hoffman K, Kvale G, Sjoden P. Information to and communication with cancer patients: improvements and psychological correlates in a comprehensive care program for patients and their relatives. *Patient Educ Counsell* 1995;25:171-82.
  - 21 Luker KA, Beaver K, Leinster SJ, Owens RG. Information needs and sources of information for women with breast cancer: a follow-up study. *J Adv Nurs* 1996;23:487-95.
  - 22 Jones RB, Navin LM, Murray KJ. Use of a community-based touch-screen public access health information system. *Health Bull* 1993;51:34-42.
  - 23 McGarry E, Jones R, Cowan B, White J. A multimedia system for personalised treatment of anxiety in primary care. In: Richards B, ed. *Current perspectives in healthcare computing*. Weybridge: BJHC Books, 1998:277-85.
  - 24 Morton AR, Patterson L, Jones R, Atkinson JM, Coia D. Personalised patient information for patients with schizophrenia living in the community. In: Richards B, ed. *Current perspectives in healthcare computing*. Weybridge: BJHC Books, 1998:94-104.
  - 25 Bailey AJ, Parmar MKB, Stephens RJ for the CHART Steering Committee. Patient-reported short-term and long-term physical and psychologic symptoms: results of the continuous hyperfractionated accelerated radiotherapy (CHART) randomised trial in non-small-cell lung cancer. *J Clin Oncol* 1998;16:3082-93.
  - 26 Rapoport Y, Kreitler S, Chaitchik S, Algor R, Weissler K. Psychosocial problems in head-and-neck cancer patients and their change with time since diagnosis. *Ann Oncol* 1993;4:69-73.
  - 27 Hall A, A'Hern R, Fallowfield L. Are we using appropriate self-report questionnaires for detecting anxiety and depression in women with early breast cancer? *Eur J Cancer* 1999;35:79-85.
  - 28 Dunn SM, Butow PN, Tattersall MHN, Jones QI, Sheldon JS, Taylor JJ, et al. General information tapes inhibit recall of the cancer consultation. *J Clin Oncol* 1993;11:2279-85.
  - 29 Tattersall MHN, Butow PN, Griffin AM, Dunn SM. The take home message: patients prefer consultation audiotapes to summary letters. *J Clin Oncol* 1994;12:1305-11.
  - 30 Miller SM. Monitoring versus blunting styles of coping with cancer influence the information patients want and need about their disease: implications for cancer screening and management. *Cancer* 1995;76:167-77.
  - 31 Street RL Jr, Voigt B, Geyer C Jr, Manning T, Swanson GP. Increasing patient involvement in choosing treatment for early breast cancer. *Cancer* 1995;76:2275-85.
  - 32 McHugh P, Lewis S, Ford S, Newlands E, Rustin G, Coombes C, et al. The efficacy of audiotapes in promoting psychological well-being in cancer patients: a randomised, controlled trial. *Br J Cancer* 1995;71:388-92.

(Accepted 11 October 1999)

Cross sectional survey of patients' satisfaction with information about cancer

Ray Jones, Janne Pearson, Sandra McGregor, W Harper Gilmour, Jacqueline M Atkinson, Ann Barrett, Alison J Cawsey, Jim McEwen

Most patients with cancer want as much information as possible appropriate to their personal needs and circumstances.^{1,2} We surveyed the views of cancer patients entering a randomised trial of computer based information.³ We examined their need for information and their satisfaction with information received and how these varied with their demographic, social, and psychological characteristics.

Patients, methods, and results

Eligible patients were those planned to receive radical radiotherapy, who knew their diagnosis, were without visual or mental handicap, and were without severe pain or symptoms causing distress. Of 715 patients asked to take part, 190 refused, 25 stating they did not want more information. Of the 525 participants, 309 had breast cancer, 129 had prostate cancer, 22 had cervical cancer, and 65 had laryngeal cancer.

Data were collected at the recruitment interview, from a questionnaire the patients completed at home shortly after, and from their case notes. Data included the information patients would like,² a hospital anxiety and depression scale,⁴ the newspaper patients read, and deprivation category (derived from postcode).⁵ Using ² tests and multiple logistic regression analysis, we compared the patients' sources and perceived quantity of information received and their satisfaction with this information, as binary variables, with their age, sex, cancer site, newspaper read (tabloid v broadsheet), deprivation category, and anxiety and

depression scores (table). Information need² ("as much as possible" v other) was considered both as a response variable and as a predictor of sources and satisfaction.

Four out of five patients wanted as much information as possible. In univariate analysis, newspaper read, deprivation category, having a connection with the health service, age, and time since diagnosis were predictors of information need. In multiple logistic regression analysis, however, only newspaper read and age remained predictors.

One in five patients were not satisfied with the information given. Univariate analysis showed that dissatisfied patients were much more likely to be depressed and were marginally more likely to be anxious or to want as much information as possible. In multiple logistic regression age, sex, and depression were predictors of dissatisfaction. Fifteen per cent of patients said there had been many differences in what they had been told by health professionals. Multiple logistic regression showed that these patients were more likely to be anxious.

Patients with breast cancer and readers of broadsheets had received more information and from more people than patients with other cancers but were not significantly more likely to be satisfied. The location of the clinician who gave the diagnosis had no effect on how much information patients had received. Younger depressed patients who wanted as much information as possible were less likely to be satisfied even though they had received more information than others. A third of patients said there were other things that they

Department of Public Health, University of Glasgow, Glasgow G12 8RZ

Ray Jones
senior lecturer in health informatics

Janne Pearson
research assistant
Sandra McGregor
research assistant

W Harper Gilmour
senior lecturer in medical statistics
Jacqueline M Atkinson
senior lecturer in behavioural sciences

Jim McEwen
professor of public health

Beatson Oncology Centre, Western Infirmary, Glasgow G11 6NT

Ann Barrett
professor of radiation oncology

continued over

BMJ 1999;319:1247-8

Percentage (number) of cancer patients agreeing with various statements about the information they had received before starting radiotherapy

	Given information from few people (v many)	Not given much information	Many differences in information given (v few or none)	Would like to have been given other information	Non-committal or dissatisfied with information given (v satisfied)
Age:					
<60	73 (193)	49 (112)	18 (45)	36 (93)	21 (56)†
> 60	82 (207)*	59 (119)	13 (31)	32 (77)	17 (43)
Sex:					
Female	71 (240)	46 (131)	17 (55)	33 (109)	17 (58)
Male	89 (160)*	69 (100)*	12 (21)	35 (61)	23 (41)†
Cancer site:					
Breast	70 (213)	44 (116)	16 (48)	33 (99)	17 (52)
Other	89 (187)*†	69 (115)*†	14 (28)	35 (71)	23 (47)
Time since diagnosis (months):					
< 3	79 (205)	57 (118)	13 (32)	29 (73)	19 (48)
4-12	73 (162)	48 (94)	14 (29)	38 (82)	19 (41)
>12	97 (28)*	64 (16)	10 (3)	43 (12)	30 (9)
Where diagnosed:					
Teaching hospital	81 (137)	51 (75)	13 (21)	35 (57)	18 (30)
Other hospital	77 (215)	56 (133)	16 (43)	35 (95)	21 (56)
Elsewhere	70 (45)	47 (22)	20 (12)	25 (15)	18 (12)
Deprivation category (Carstairs category):					
Affluent (1+2)	79 (84)	42 (38)	21 (22)	37 (39)	19 (20)
Middle (3+4)	76 (159)	52 (94)	14 (28)	33 (67)	17 (35)
Deprived (5+7)	79 (153)	62 (95)*†	14 (26)	34 (64)	23 (44)
Newspaper read:					
Broadsheet	73 (143)	44 (71)	16 (31)	34 (66)	16 (32)
Tabloid	81 (211)†	62 (137)*†	14 (35)	33 (85)	21 (54)
Connection with NHS:					
Yes	73 (149)	46 (77)	16 (31)	35 (68)	18 (36)
No	80 (221)	59 (137)*	15 (39)	33 (90)	21 (56)
Hospital anxiety and depression scale:					
Anxiety:					
No	81 (261)	53 (141)	12 (38)	28 (83)	16 (49)
Borderline	73 (75)	52 (44)	14 (14)	35 (34)	23 (23)
Yes	71 (62)	58 (44)	29 (24)*†	54 (46)*†	27 (23)
Depression:					
No	77 (346)	51 (190)	13 (58)	31 (134)	17 (74)
Borderline	80 (37)	64 (27)	37 (15)*	44 (20)	37 (16)
Yes	88 (15)	86 (12)*	19 (3)	71 (12)*	50 (8)*†
Attitude to information:					
As much as possible	76 (324)	51 (180)	16 (66)	36 (148)*†	21 (88)
Only good news or no details	84 (67)	67 (43)*	12 (9)	20 (15)	12 (9)
All patients	78 (400/516)	54 (231/430)	15 (76/494)	34 (170/503)	19 (99/509)‡

*Significant predictors (P<0.01) in analysis. †Significant predictors (P<0.05) in multiple logistic regression analysis.

Department of Computer Science, Heriot Watt University, Edinburgh EH14 4AS

Alison J Cawsey lecturer in computer science

Correspondence to: R Jones
r.b.jones@udcf.gla.ac.uk

would like to have been told. Patients most commonly required further information on the effects of treatment and prognosis and recovery.

Comment

One in five of 525 patients starting radiotherapy were not satisfied with the information they had received. Breast cancer patients were better provided with information than patients with other cancers but were not necessarily more satisfied with it as a result. More should be done to help patients with other cancers obtain suitable information.

Three of the outcome measures were associated with anxiety or depression. The nature of the relation of anxiety, depression, and information among cancer patients would be worthy of further study.

As in other work,² we found that most cancer patients wanted as much information as possible. However, our percentage slightly overestimates this, as refusal to take part in the study might indicate lack of desire for further information. Although those patients living in affluent areas wanted more information, type of newspaper read was a better indicator of information need. Few studies have used this intuitively obvious indicator, and we suggest its use, among older patients, in both clinical practice and research.

We thank the consultants (particularly Nick Reed and Tim Habeshaw, who were directors of the Beatson Oncology Centre during the study), other medical staff, cancer nurse specialists, radiographers, medical records staff, and other staff at the Beatson Oncology Centre for their collaboration with this project; the patients who took part in the study; Lynn Naven, who worked as a locum for SMcG during three months' sick leave; Ed Carter; Ross Morton and Keith Murray, who helped with various aspects of computing; Charles Gillis and Cathy Meredith, who advised on research design; Sally Tweddle, who made available unpublished protocols and papers; and colleagues in the University of Glasgow and the Beatson Oncology Centre who commented on the manuscript.

Contributors: RJ had the original idea for the study, designed the study, was the main grant holder, supervised the research assistants, analysed the data, and wrote the paper. JP carried out the pilot study, contributed to the design, was a research assistant with day to day responsibility for data collection and running the study, undertook preliminary analysis, and contributed to final analysis and editing of paper. SMcG contributed to the design of the study, was a research assistant with day to day responsibility for data collection and running the study, undertook some preliminary analysis, and edited the paper. WHG advised on design of the study and analysis of the data and edited the paper. JMA advised on the design of the study and choice of psychological measures, edited the paper, and was a grant holder. AB discussed the original idea, set up opportunities for the pilot study and main study, contributed to the design of the study, edited the paper, and was a grant holder. AJC discussed the original idea, contributed to the design of the study, discussed the analysis of data, edited the paper, and was a grant holder. JMcE discussed the original idea, set up opportunities for the pilot study and main study, edited the paper, and was a grant holder. RJ and AB are guarantors for the study.

Funding: Scottish Office Home and Health Department grant number K/OPR/2/2/D248. The views expressed in this paper are those of the authors and do not represent the views of the Scottish Office.

Competing interests: None declared.

- 1 National Cancer Alliance. Patient-centred cancer services? What patients say Oxford: National Cancer Alliance, 1996.
- 2 Meredith C, Symonds P, Webster L, Lamont D, Pyper E, Gillis CR, et al. Information needs of cancer patients in west Scotland: cross sectional survey of patients' views. *BMJ* 1996;313:724-6.
- 3 Jones R, Pearson J, McGregor S, Cawsey AJ, Barrett A, Craig N, et al. Randomised trial of personalised computer based information for cancer patients. *BMJ* 1999;319:1241-7.
- 4 Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-70.
- 5 Carstairs V, Morris R. Deprivation and health in Scotland. Aberdeen: Aberdeen University Press, 1991.

(Accepted 11 October 1999)

Endpiece Bill Gates's strength

The fox knows many things; the hedgehog knows one big thing; the 800 pound gorilla doesn't give a shit what anybody knows. [Bill] Gates's great strength was in combining all these attributes.

John Lanchester, *London Review of Books*
September 1999, p 5.