- 16 Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE, et al. Risk factors for hip fracture in white women. The study of osteoporotic fractures research group. *N Engl J Med* 1995;332:767-73.
- 17 Black DM, Palermo L, Nevitt MC, Genant HK, Epstein R, San Valentin R, et al. Comparison of methods for defining prevalent vertebral fractures: the study of osteoporotic fractures. *J Bone Miner Res* 1995;10: 890-902.
- 18 Bauer DC, Browner WS, Cauley JA, Orwoll ES, Scott JC, Black DM, et al. Factors associated with appendicular bone mass in older women. The study of osteoporotic fractures research group. *Ann Intern Med* 1993;118:657-65.
- 19 Orwoll ES, Bauer DC, Vogt TM, Fox KM. Axial bone mass in older women. Study of osteoporotic fractures research group. *Ann Intern Med* 1996;124:187-96.
- 20 Leone NC, Stevenson CA, Hilbish TF, Sosman MC. A roentgenologic study of human population exposed to high-fluoride domestic water (a 10-year study). In: McClure FJ, ed. *Fluoride drinking waters*. Bethesda, MD: National Institute of Dental Research, 1962 (DHEW Publ No (PHS)62-825).
- 21 Bernstein DS, Sadowsky N, Hegsted DM, Guri D, Stare FJ. Prevalence of osteoporosis in high- and low-fluoride areas in North Dakota. *JAMA* 1966;198:499-504.
- 22 Ansell BM, Lawrence JS. Fluoridation and the rheumatic diseases. A comparison of rheumatism in Watford and Leigh. *Ann Rheum Dis* 1966;25:67-75.
- 23 Kroger H, Alhava E, Honkanen R, Tuppurainen M, Saarikoski S. The effect of fluoridated drinking water on axial bone mineral density—a population-based study. *Bone Miner* 1994;27:33-41.
- 24 Hillier S, Cooper C, Killingray S, Russell G, Hughes H, Coggon D. Fluoride in drinking water and risk of hip fracture in the UK: a case-control study. *Lancet* 2000;335:265-9.
- 25 Ad-Hoc Subcommittee on Fluoride. *Review of fluoride benefits and risks*. Bethesda, MD: Public Health Service, Department of Health and Human Services, 1991:17.
- 26 Whitford GM. The metabolism and toxicity of fluoride. *Monogr Oral Sci* 1996;16(2):1-153.
- 27 Farley JR, Tarbaux N, Hall S, Baylink DJ. Evidence that fluoride-stimulated ³[H]-thymidine incorporation in embryonic chick calvarial cell cultures is dependent on the presence of a bone cell mitogen, sensitive to changes in the phosphate concentration, and modulated by systemic skeletal effectors. *Metabolism* 1988;37:988-95.
- 28 Dabeka RW, McKenzie AD. Survey of lead, cadmium, fluoride, nickel, and cobalt in food composites and estimation of dietary intakes of these elements by Canadians in 1986-1988. *J AOAC Int* 1995;78:897-909.
- 29 Lahti SM, Uusitalo U, Feskens E, Haw U, Tuomilehto J, Luoma H. Fluoride and sugar intake among adults and youth in Mauritius: preliminary results. *Adv Dent Res* 1995;9:21-5.
- 30 Silbergeld EK, Davis DL. Role of biomarkers in identifying and understanding environmentally induced disease. *Clin Chem* 1994;40:1363-7.

(Accepted 5 July 2000)

Audit of oxygen prescribing before and after the introduction of a prescription chart

M E Dodd, F Kellet, A Davis, J C G Simpson, A K Webb, C S Haworth, R McL Niven

North West Lung
Centre, South
Manchester
University Hospitals
Trust, Manchester
M23 9LT

M E Dodd
specialist
physiotherapy
clinician
F Kellet
senior physiotherapist
A Davis
senior physiotherapist

A K Webb
consultant physician
C S Haworth
research fellow
R McL Niven
consultant physician

Stepping Hill
Hospital, Poplar
Grove, Stockport
SK2 7JE

J C G Simpson
senior registrar

Correspondence to:
M E Dodd
doddmary@
hotmail.com

BMJ 2000;321:864-5

Oxygen, used to treat hypoxaemia, may be lethal and should therefore be considered a drug and be prescribed.¹ It is, however, recognised that oxygen is poorly prescribed by doctors.² To ensure the safe and effective delivery of oxygen the prescription should include the flow rate, the concentration, the delivery device, the duration, and the method for monitoring treatment.² We audited the prescription of oxygen to inpatients by doctors before and after the introduction of a specific prescription chart.

Participants, methods, and results

Junior doctors at the North West Lung Centre are given two lectures on practical aspects of oxygen delivery and prescribing at the beginning of their one year's rotation in respiratory medicine. In 1997 and 1998 the doctors were informed that an audit of their prescribing practice would take place some time during the next year. The outcome measures of the audit were whether the oxygen was prescribed and whether the prescription was accurate—that is, that the audit matched patient use in relation to the delivery device and that the flow rate and concentration were appropriate to that device.

The audit was conducted on three respiratory wards over three months. FK and AD identified all patients receiving oxygen within 24 hours of admission, and they recorded the device, oxygen concentration, and flow rate appropriate to the device for each patient. They consulted a drug Kardex for a prescription of the proposed oxygen treatment. If a prescription was present they recorded the device, concentration, and flow rate. After the first audit, a specific prescription chart for oxygen was developed to encompass all the oxygen delivery systems used on the

wards (figure). The methodology for the second audit was identical to that of the first, with the exception that both the chart and the drug Kardex were examined for the prescription. The χ^2 test was used to analyse the prescription of oxygen before and after the introduction of the chart.

Overall, 115 patients were identified as receiving oxygen in the first audit and 121 in the second. In the first audit oxygen was prescribed for 63 of the 115 (55%) patients. After the chart was introduced the number of oxygen treatments prescribed increased to 110 of 121 (91%) patients ($P < 0.001$). The prescription was accurate for eight (7%) patients in the first audit and 93 (77%) in the second. The accuracy of prescription was 94% (73 of 78 patients) with the chart and 63% (20 of 32) with the drug Kardex.

Comment

Our first audit showed that oxygen was infrequently well prescribed, as previously described.^{2,3} Junior doctors poorly understand the effects and dangers of oxygen, and lectures alone were insufficient to ensure safe and effective practice.⁴ The prescription chart for oxygen listed the delivery devices, guided the doctor to prescribe the appropriate concentration and flow rate, and provided additional notes for the specific indications of each device.

The most common omission from the prescriptions was flow rate. The flow rate of fixed concentration masks should be adjusted for patients with high peak inspiratory flows. Flow rate is the only variable that is prescribed with nasal cannulas, and an accurate prescription of flow rate is essential as hypercapnic respiratory failure may occur.⁴ Oxygen for delivery by nasal cannula is often prescribed by

concentration, with the assumption that an inspired oxygen concentration of 24% equates to 2 l/min; concentrations of 24%-35% with 2 l/min have been described.⁵

The prescription of oxygen is complex, and a drug Kardex does not accommodate the precise details required for the variety of delivery devices. This was clearly shown in the second audit, where the accuracy of the prescription was greater with the chart than with the drug Kardex. We have shown that a specific prescription chart for oxygen improved clinical practice in our specialist medical respiratory centre, and we recommend the use of such a chart.

Contributors: MED designed the study, analysed and interpreted the results, and wrote the paper. FK and AD analysed and interpreted the data. JCGS designed the study and the chart. AKW and CSH critically revised the paper. RMcLN designed the study and the chart and critically revised the paper.

Competing interests: None declared.

- 1 British National Formulary 1999;37:153.
- 2 Bateman NT, Leach RM. ABC of oxygen. Acute oxygen therapy. *BMJ* 1998;317:798-801.
- 3 Small D, Duha A, Wieskopf B, Dajczman E, Laporta D, Kreisman H, et al. Uses and misuses of oxygen in hospitalised patients. *Am J Med* 1992;92:591-5.

The North West Lung Centre Oxygen Prescription Chart																						
CONTINUOUS/NIGHT-TIME/DAY-TIME/PRN (please circle)																						
Tick device	NAME	HOSPITAL NUMBER				WARD				AGE												
NASAL CANNULAE/NIPPV		CHANGE 1				CHANGE 2				CHANGE 3				CHANGE 4				CHANGE 5				Nasal Cannulae are best suited for chronic usage. Concentration of oxygen delivered depends on factors other than flow rate.
Circle flow rate L/min		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	
Time & Date																						
Signature																						
VENTIMASK		CHANGE 1				CHANGE 2				CHANGE 3				CHANGE 4				CHANGE 5				Fixed concentration masks are suited to patients where concentration is critical. A common error is to fail to provide sufficient flow. Increased flow rate DOES NOT alter the concn or increase the chance of CO ₂ retention but reduces work of breathing. Flow rate changes for ventimask only can be prescribed without consultation with a doctor.
Tick concentration required and specify flow rate		CONCENTRATION		FLOW		FLOW		FLOW		FLOW		FLOW		FLOW								
Flow should be adjusted to patient comfort.		24%	(2-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)	(4-15)							
Minimum and maximum allowable flow rates indicated		35%	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)	(8-15)							
		40%	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)	(12-15)							
		60%	15	15	15	15	15	15	15	15	15	15	15	15	15							
Time & Date																						
Signature																						
HIGH FLOW																		High flow rates are only required in special circumstances in breathless, hypoxic patients with a high respiratory drive				
Record concentration (33-100%)																						
Time & Date																						
Signature																						

Prescription chart for oxygen introduced after first audit

- 4 Davies RJO, Hopkin JM. Nasal oxygen in exacerbations of ventilatory failure: an underappreciated risk. *BMJ* 1989;299:43-4.
- 5 Bazuaye EA, Stone TN, Corris PA, Gibson GJ. Variability of inspired oxygen concentration with nasal cannulas. *Thorax* 1992;47:609-11.

(Accepted 11 May 2000)

Refused and granted requests for euthanasia and assisted suicide in the Netherlands: interview study with structured questionnaire

Ilinka Haverkate, Bregje D Onwuteaka-Philipsen, Agnes van der Heide, Piet J Kostense, Gerrit van der Wal, Paul J van der Maas

In 1995, physicians in the Netherlands received 9700 explicit requests for euthanasia or physician assisted suicide, of which 37% were granted and carried out.¹ Among the remaining requests, about half were refused by the physician; in the rest of the cases either the patient died before a decision had been reached or the physician's promise of help could be effected, or the patient withdrew the request.² Knowledge of specific characteristics of refused and granted requests for euthanasia or physician assisted suicide may give insight into physicians' decision making and into the role of criteria for prudent practice. We therefore compared the characteristics of refused and granted requests.

Subjects, methods, and results

In 1995 and 1996, 405 Dutch physicians, randomly sampled nationwide and stratified by specialty and region, were interviewed by over 30 specifically trained and experienced physicians using a structured questionnaire. The response rate was 89%. Euthanasia was defined as the administration of drugs with the explicit intention of ending the patient's life, at the patient's explicit request. Assisted suicide was defined as the prescribing or supplying of drugs with the explicit intention of enabling the patient to end his or

her own life. All physicians were asked to describe their most recent case of a granted request (134 physicians had had such a case) and their most recent case of a refused request (148 physicians had had such a case).

Patients whose requests were refused, compared with patients whose requests were granted, were more often female and aged over 80; were less likely to have cancer; were more likely to have depression as a predominant complaint; were more likely to have a remaining life span of over six months; were less likely to have made a highly explicit request; were less likely to be competent; were less likely to be suffering utterly "hopelessly and unbearably," and were more likely to have access to alternatives for treatment (table).

In both the refused and the granted requests "avoiding loss of dignity" (42% (95% confidence interval 31.6% to 52.4%) and 56% (46.3% to 66.2%) respectively) and "unbearable or hopeless suffering" (39% (29.0% to 48.8%) and 74% (64.9% to 82.6%)) were most often mentioned as the patient's reason for requesting euthanasia or physician assisted suicide. Only two reasons were mentioned more often in refused requests than in granted requests: "weariness of life" (40% (29.8% to 50.5%) and 18% (10.2% to 25.5%) respectively) and "not wanting to become

Institute for Research in Extramural Medicine, Vrije Universiteit Amsterdam, van der Boechorststraat 7, 1081 BT Amsterdam, Netherlands
Ilinka Haverkate
psychologist

Department of Social Medicine, Institute for Research in Extramural Medicine, Vrije Universiteit Amsterdam
Bregje D Onwuteaka-Philipsen
researcher
Gerrit van der Wal
professor

continued over

BMJ 2000;321:865-6