- 2 Pew Internet and American Life. The online health care revolution: how the web helps Americans take better care of themselves (26 Nov 2000). www.pewinternet.org/reports/toc.asp?Report=26 (accessed 1 Nov 2001).
- 3 Quintana Y, Feighner JW, Wathen CN, Sangster LM, Marshall JN. Preventive health information on the internet. Qualitative study of consumers' perspectives. *Can Fam Physician* 2001;47:1759-65.
- 4 Boyer C, Selby M, Scherrer JR, Appel RD. The Health on the Net code of conduct for medical and health websites. *Comput Biol Med* 1998;28: 603-10.
- 5 e-Health Ethics Initiative. e-Health code of ethics. *J Med Internet Res* 2000;2:e9. www.jmir.org/2000/2/e9/ (accessed 1 Nov 2001).
- 6 Nielsen J. *Usability engineering*. San Diego, CA: Morgan Kaufmann, 1993.
- 7 Kitzinger J. Qualitative research. Introducing focus groups. *BMJ* 1995;311:299-302.
- 8 Strauss AL, Corbin JM. *Basics of qualitative research: grounded theory procedures and techniques*. Newbury Park, CA: Sage Publications, 1990.
- 9 Eysenbach G. Consumer health informatics. *BMJ* 2000;320:1713-6. (Accepted 30 January 2002)

Breast cancer on the world wide web: cross sectional survey of quality of information and popularity of websites

Funda Meric, Elmer V Bernstam, Nadeem Q Mirza, Kelly K Hunt, Frederick C Ames, Merrick I Ross, Henry M Kuerer, Raphael E Pollock, Mark A Musen, S Eva Singletary

Abstract

Objectives To determine the characteristics of popular breast cancer related websites and whether more popular sites are of higher quality.

Design The search engine Google was used to generate a list of websites about breast cancer. Google ranks search results by measures of link popularity—the number of links to a site from other sites. The top 200 sites returned in response to the query “breast cancer” were divided into “more popular” and “less popular” subgroups by three different measures of link popularity: Google rank and number of links reported independently by Google and by AltaVista (another search engine).

Main outcome measures Type and quality of content.

Results More popular sites according to Google rank were more likely than less popular ones to contain information on ongoing clinical trials (27% v 12%, $P=0.01$), results of trials (12% v 3%, $P=0.02$), and opportunities for psychosocial adjustment (48% v 23%, $P<0.01$). These characteristics were also associated with higher number of links as reported by Google and AltaVista. More popular sites by number of linking sites were also more likely to provide updates on other breast cancer research, information on legislation and advocacy, and a message board service. Measures of quality such as display of authorship, attribution or references, currency of information, and disclosure did not differ between groups.

Conclusions Popularity of websites is associated with type rather than quality of content. Sites that include content correlated with popularity may best meet the public's desire for information about breast cancer.

Introduction

Recent surveys show that 40-54% of patients access medical information via the internet and that this information affects their choice of treatment.¹⁻⁵ Although the quality of medical information on the world wide web has been an area of increasing concern,⁶⁻¹¹ the factors that contribute to popularity of websites have not been systematically studied.

Understanding the determinants of website popularity has implications for clinicians and medical

centres that recognise the need to provide information about themselves via the internet. Website designers who understand the information needs of the public can attract visitors to their site. Knowing what patients are investigating on the web may help clinicians to educate themselves and their patients.

Two measures of website popularity are “click popularity” and “link popularity.”¹² Click popularity is the frequency with which users have visited (clicked on) a site.¹³ Although some search engines, such as Direct Hit, measure click popularity, this information is not publicly available for a large number of websites. Furthermore, click popularity is subject to artificial marketing manipulations.¹⁴ Link popularity, which is less susceptible to manipulation,¹⁵ relies on links from sites to other sites rather than on statistics about usage. High link popularity is thought to dramatically increase traffic to a site.¹⁶ Link popularity, sometimes referred to as “peer review popularity,” has been proposed as an objective way of identifying high quality websites.¹⁷⁻¹⁹ Google ranks results of searches by using a proprietary link popularity algorithm that takes into account the number of links and the “importance” of the linking sites.^{15 17}

Breast cancer is one of the most common health related search topics among users of the internet.²⁰ Previous studies have evaluated use of the internet by women with breast cancer and the quality of selected sites.^{10 21 22} A recent study found information about breast cancer on the web to be more complete and accurate than for other conditions.²² We are not aware, however, of work that attempts to determine what makes some sites more popular than others. The purpose of our study was to identify the determinants of link popularity of websites about breast cancer and to test the hypothesis that more popular sites are of higher quality.

Methods

Selection of websites

We used the search term “breast cancer” on Google (www.google.com accessed 19 Oct 2000) to generate a list of sites. We examined the first 200 of approximately 600 000 English language sites. Of these, 185 (93%) were accessible, but one was excluded as its content was only peripherally related to breast cancer.

Section of Breast Surgery,
Department of
Surgical Oncology,
University of Texas
M D Anderson
Cancer Center,
1515 Holcombe
Boulevard, Box 444,
Houston, TX
77030, USA

Funda Meric
assistant professor

Nadeem Q Mirza
research investigator

Kelly K Hunt
associate professor

Frederick C Ames
professor

Merrick I Ross
professor

Henry M Kuerer
assistant professor

Raphael E Pollock
professor

S Eva Singletary
professor

Stanford Medical
Informatics,
Department of
Internal Medicine,
Stanford University
Medical School,
Stanford, CA, USA
Mark A Musen
associate professor

University of Texas
Health Science
Center at Houston,
School of Health
Information
Sciences, Houston,
TX, USA

Elmer V Bernstam
assistant professor

Correspondence to:
F Meric fmeric@
mail.
mdanderson.org

BMJ 2002;324:577-81

Determination of popularity

Because there is no standard way to assess link popularity, we used three different measures: Google rank and number of links reported by Google and by AltaVista (on www.altavista.com). Of the top 200 sites returned by Google, we defined the first 100 (Google rank 1-100) as "more popular" and the second 100 (Google rank 101-200) as "less popular." We obtained the number of links in Google and AltaVista by entering each site's universal resource locator (URL) into the search string "link:URL".

Google provided the number of linking sites for 162 sites and AltaVista for 148 sites. We excluded from analysis any sites for which the number of links was not available. The median number of links was 51 according to Google and 21 according to AltaVista. We considered sites with a number of links greater than the median to be "more popular" and sites with fewer links to be "less popular." To assess whether popular sites were displayed by multiple search engines, we repeated the search on four search engines often used by patients: Yahoo (categorical), Excite, AltaVista, and Infoseek.⁴

Evaluation of websites

A breast oncologist (FM) evaluated the sites within four weeks of the original search. Links within each site were pursued until all medical information about breast cancer was evaluated. A median of four pages (range 1-11) were evaluated for each site. Type and quality of content were recorded. Affiliation was determined on the basis of the information provided by the site. Sites were divided into professional (government, universities, major medical centres), non-profit organisation, and commercial (all others).

We assessed quality of content by criteria known as the "JAMA benchmarks": display of authorship of medical content; source (attribution or references); date of update; and disclosure of ownership, sponsorship, advertising policies, or conflicts of interest. We also documented whether each site displayed its webmaster's email address or a Health on the Net

Table 1 Characteristics of breast cancer websites evaluated (n=184). Values are numbers (percentages)

Characteristic	Sites
Affiliation:	
Commercial	84 (46)
Non-profit organisation	64 (35)
University or medical centre	22 (12)
Government	14 (8)
Specialisation:	
Breast cancer exclusively	93 (51)
Breast cancer pages of health site	91 (49)
Content type:	
Medical facts	151 (82)
Opportunities for psychosocial adjustment	65 (35)
Human interest stories	47 (26)
Ongoing available trials	35 (19)
Message board service	23 (13)
Results of clinical trials	14 (8)
Chat site	14 (8)
Site for medical questions	11 (6)
Content presentation:	
Images	87 (47)
Audio	15 (8)
Video or animation	14 (8)

Table 2 Medical facts contained in breast cancer websites (n=125). * Values are numbers (percentages)

Topic	Sites covering topic
Risk factors	72 (58)
General information	65 (52)
Screening	54 (43)
Surgical treatment	39 (31)
Chemotherapy	37 (30)
Hormonal treatment	32 (26)
Radiation treatment	29 (23)
Diet or nutrition	28 (22)
General prevention	27 (22)
Chemoprevention	24 (19)
Reconstruction	23 (18)
Prophylactic mastectomy	16 (13)
Complementary medicine	14 (11)
Psychology	12 (10)

*Only sites containing medical facts on the website rather than through links to other sites are included.

Table 3 Quality of medical content (n=184). Values are numbers (percentages)

Quality measure	Sites
Display of authorship	105 (57)
Display of attribution or references	116 (63)
Display of currency	82 (45)
Disclosure	128 (70)
Display of Health on the Net seal	27 (15)
Display of email address	125 (68)

(www.hon.ch/) seal. Health on the Net is a non-profit foundation with an eight point code of conduct for sites providing health information.²⁵ Sites that comply with the Health on the Net code are allowed to display the seal, but continued compliance is not systematically enforced.

Analysis

We used Pearson χ^2 analysis to compare more popular and less popular websites. We performed separate analyses for each of the three measures of link popularity. We considered groups to be significantly different if $P \leq 0.05$ in at least two of three analyses.

Results

Website characteristics

Table 1 lists the characteristics of the top 184 accessible sites returned by Google. Twenty seven (15%) provided medical facts on the site as well as through links to other sites, and 125 (68%) had medical facts displayed at the website only. Table 2 shows the medical facts displayed in this second group of websites.

Table 3 shows indicators of quality. Of the 184 sites, 105 (57%) displayed some evidence of authorship, but only 32 (17%) displayed the name, qualifications, and institutional affiliation of the author. Sixteen (9%) of sites had all four JAMA benchmarks (authorship, references, currency, and disclosure), 48 (26%) had three, 68 (37%) had two, 43 (23%) had one, and 9 (5%) had none. Forty five sites (25%) displayed a disclaimer that the information provided should not substitute for consultation with a physician.

A Health on the Net seal was displayed on 27 (15%) sites. Commercial sites were more likely than sites of professional groups or of organisations to display the

Number of accurate and inaccurate websites, based on number of JAMA benchmarks met. A website was considered inaccurate if it contained one or more inaccurate statements

seal—21/84 (25%) *v* 3/36 (8%) *v* 3/64 (5%) ($P=0.001$). None of the sites with a seal actually complied with all eight Health on the Net criteria or with all four JAMA benchmarks.

Of the 184 sites, 12 (7%) contained inaccurate medical statements. Commercial sites contained inaccurate statements more often than did sites of professional groups or of organisations—11/84 (13%) *v* 1/36 (3%) *v* 0/64 ($P=0.004$). Three (16%) of 19 commercial sites that displayed the Health on the Net seal contained inaccurate statements. Higher quality sites (at least three JAMA benchmarks) were less likely to contain inaccurate information than lower quality sites (fewer than three JAMA benchmarks)—1/64 (2%) *v* 11/120 (10%) ($P=0.047$) (figure). None of the 16 sites that met all four JAMA benchmarks contained inaccurate information.

Determinants of popularity

Type of content differed significantly between more popular and less popular websites (table 4). Sites that were more popular by at least two of three popularity measures were more likely to contain information about ongoing clinical trials, results of randomised clinical trials, results of other breast cancer research,

information on legislation and advocacy, and information on opportunities for psychosocial adjustment and to allow interaction through a message board service.

We then evaluated differences in the topics of medical facts presented between the more popular and less popular sites (table 2). This analysis was carried out only for the 125 sites that displayed medical information. More popular sites were more likely to discuss breast reconstruction—15/57 (26%) *v* 8/68 (12%) by Google rank ($P=0.037$), 15/51 (29%) *v* 5/57 (9%) by number of links in Google ($P=0.002$), and 16/50 (32%) *v* 6/48 (13%) by number of links in AltaVista ($P=0.018$)—and psychology topics such as depression—11/51 (21%) *v* 1/57 (2%) by number of links in Google ($P=0.001$) and 11/50 (22%) *v* 1/48 (2%) by number of links in AltaVista ($P=0.002$).

More popular and less popular websites did not differ in any of the quality measures studied (table 4). Furthermore, the presence of inaccurate information did not differ between more popular and less popular sites.

Evaluation of popularity measures

We evaluated the concordance between our measures of popularity. The median number of linking sites as measured by Google and AltaVista was significantly higher for sites that were more popular by Google rank than for less popular sites (Google: 82 *v* 21, $P<0.001$; AltaVista: 48 *v* 10, $P<0.001$). The number of links as measured by Google strongly correlated with the number of links as measured by AltaVista (Pearson coefficient 0.806, $P<0.001$).

We hypothesised that link popularity would correlate with a site being displayed by multiple search engines. Of the top 184 accessible sites displayed by Google, AltaVista displayed 24 (13%), Yahoo displayed 41 (22%), Infoseek displayed 58 (32%), and Excite displayed 84 (46%). Indeed, more popular sites were more likely than less popular sites to be displayed by multiple search engines (table 5).

Table 4 Content of breast cancer websites by popularity. Values are numbers (percentages) unless stated otherwise

Characteristic	Google rank			No of links (Google)			No of links (AltaVista)		
	More popular (rank 1-100) (n=90)	Less popular (rank 101-200) (n=94)	P value	More popular (>50 links) (n=79)	Less popular (≤50 links) (n=83)	P value	More popular (>20 links) (n=75)	Less popular (≤20 links) (n=73)	P value
Type of content									
Medical facts	73 (81)	78 (83)	0.741	61 (77)	72 (87)	0.114	56 (75)	65 (89)	0.024
Information on ongoing clinical trials	24 (27)	11 (12)	0.010	23 (29)	12 (15)	0.023	22 (29)	11 (15)	0.040
Results of randomised clinical trials	11 (12)	3 (3)	0.021	10 (13)	1 (1)	0.004	9 (12)	4 (6)	0.161
Results of other research	16 (18)	8 (9)	0.062	16 (20)	6 (7)	0.016	15 (20)	6 (8)	0.037
Opportunities for psychosocial adjustment	43 (48)	22 (23)	0.001	37 (47)	23 (28)	0.012	37 (49)	21 (29)	0.010
Human interest stories	28 (31)	19 (20)	0.090	25 (32)	19 (23)	0.211	26 (35)	15 (21)	0.055
Legislation or advocacy information	22 (24)	27 (29)	0.510	27 (34)	17 (21)	0.050	31 (41)	14 (19)	0.003
Message board service	12 (13)	11 (12)	0.738	17 (22)	5 (6)	0.004	16 (21)	4 (6)	0.005
Chat service	8 (9)	6 (6)	0.522	7 (9)	7 (8)	0.923	7 (9)	5 (7)	0.580
Forum for questions	7 (8)	4 (4)	0.314	4 (5)	6 (7)	0.567	6 (8)	5 (7)	0.719
Quality of content									
Authorship	45 (50)	60 (64)	0.058	36 (46)	57 (69)	0.003	36 (48)	43 (59)	0.184
Attribution or references	55 (61)	57 (61)	0.948	42 (53)	55 (66)	0.089	41 (55)	51 (70)	0.057
Currency	40 (44)	42 (45)	0.974	34 (43)	36 (43)	0.966	35 (47)	31 (43)	0.607
Disclosure	61 (68)	67 (71)	0.606	57 (72)	54 (65)	0.331	53 (71)	52 (71)	0.940
Display of Health on the Net seal	14 (16)	13 (14)	0.741	12 (15)	13 (16)	0.934	11 (15)	13 (18)	0.604
Display of email address	67 (74)	58 (62)	0.064	57 (72)	53 (64)	0.258	55 (73)	50 (69)	0.517

Table 5 Display by other search engines on the basis of website link popularity. Values are numbers (percentages) unless stated otherwise

Search engine	Google rank			No of links (Google)			No of links (AltaVista)		
	More popular (rank 1-100) (n=90)	Less popular (rank 101-200) (n=94)	P value	More popular (>50 links) (n=79)	Less popular (≤50 links) (n=83)	P value	More popular (>20 links) (n=75)	Less popular (≤20 links) (n=73)	P value
AltaVista	15 (17)	9 (10)	0.153	20 (25)	3 (4)	<0.001	21 (28)	3 (4)	<0.001
Yahoo	26 (29)	15 (16)	0.035	33 (42)	7 (8)	<0.001	37 (49)	4 (6)	<0.001
Excite	53 (59)	31 (33)	<0.001	58 (73)	21 (25)	<0.001	64 (85)	19 (26)	<0.001
Infoseek	38 (42)	20 (21)	0.002	45 (57)	12 (15)	<0.001	47 (63)	10 (14)	<0.001

On query with the search string "breast cancer," AltaVista returned approximately 300 000 sites but provided links to 200, Yahoo category search returned 361, and Infoseek returned approximately 600 000 sites but provided links to 335. Excite returned approximately 60 000 results, of which 500 were evaluated.

To assess the correlation between click popularity and link popularity, we evaluated the top 10 sites returned by Direct Hit, which ranks sites based on click popularity and duration of visits. Nine of the 10 sites were in the top 200 by Google rank, eight were among the more popular sites by Google rank, and five were among the top 20 by Google rank.

Discussion

To meet the demand for health information on the web, it is important to identify the factors that influence popularity of websites. Our results show that type rather than quality of content determines popularity. To our knowledge, ours is the first study to assess the popularity as well as the quality and accuracy of health related websites.

We found that many breast cancer websites do not comply with the JAMA benchmarks, but we found higher compliance than previously reported.^{10 11} This may reflect an improvement in quality of websites over the past few years or a difference between search engines used in the studies.

Since accessibility and ranking of websites vary with the search engine used, the overlap between Google and other search engines of only 13-46% is not surprising. We found that "more popular" sites were more likely to be displayed by multiple search engines. If a site is not displayed, it is unlikely to be visited by users of the internet; thus the more popular sites by our measures of link popularity should indeed be the more popular sites among users of the internet. Our finding that eight of the top 10 sites according to Direct Hit were among the more popular sites by Google rank also supports this assertion.

Our results confirm those of an earlier study that found no correlation between measures of quality and link popularity.²⁴ We may have selected higher quality sites by examining the top 200 of about 600 000 sites returned by Google. Significant differences in quality might have emerged if we had increased our sample size or compared the top 100 sites with sites of lower popularity, such as those ranked 1000-1100. Using less popular sites, however, would not have allowed us to correlate multiple measures of link popularity, as most sites would have no incoming links.

One limitation of our study is that we performed multiple comparisons. Another limitation is that a single reviewer (FM) assessed quality and accuracy. To mitigate this, we used objective criteria whenever possible. For example, we used the presence or absence of authorship information rather than author authority. Accuracy is inherently subjective, so our results

What is already known on this topic

Patients are using the world wide web to search for health information

Breast cancer is one of the most popular search topics

Characteristics of popular websites may reflect the information needs of patients

What this study adds

Type rather than quality of content correlates with popularity of websites

Measures of quality correlate with accuracy of medical information

should be confirmed by studies using a panel of experts. Multiple, non-expert reviewers, however, may not be better than a single expert reviewer.

In one survey, only 14% of patients expressed uncertainty about the accuracy of medical information on the web.⁴ We found that higher quality sites contain more accurate information. Objective measures of quality may help lay users to assess online health information.

Self regulation has been advocated as a way of maintaining the quality of online medical content. Our finding that sites displaying a Health on the Net seal did not comply with the Health on the Net code emphasises the limitation of self regulation. It remains the responsibility of the medical community to ensure adequate quality of online medical content, to educate the public regarding quality measures, and to direct patients to sites of known quality.

Link popularity, which can be assessed automatically, has been proposed as an indirect measure of quality.¹⁹ This is analogous to citation analysis, a somewhat controversial approach to measuring quality in the printed literature. Although link popularity may identify sites of interest, it does not correlate with quality of content. The growing number of users of the internet searching for health information indicates an unmet need for information. Understanding what patients are looking for on line may help us meet their need for health information.

We thank Valerie Natale and Stephanie Deming for editorial assistance and Herbert Kaizer and Soumya Raychaudhuri for critical reading of the manuscript.

Contributors: FM was involved in the conception and design of the study, reviewed all the websites, performed the analysis

and interpretation of data, and drafted the paper. EVB was involved in the conception and design of the study, collection of data on links, interpretation of data, and drafting of the paper. NQM carried out the statistical analysis. KKH, FCA, MIR, HMK, and REP were involved in interpreting the data and revised the paper for intellectual content. MAM and SES advised on methods and revised the paper for intellectual content. FM and EVB are the guarantors.

Funding: Supported in part by Grant LM06594 from the National Library of Medicine (EVB).

Competing interests: None declared.

- Metz JM, Devine P, DeNittis A, Stambaugh M, Jones H, Goldwein J, et al. Utilization of the internet by oncology patients to obtain cancer related information. *Proc Am Soc Clin Oncol* 2001;20:395a (abstract 1575).
- Yakren S, Shi W, Thaler H, Agre P, Bach PB, Schrag D, et al. Use of internet and other information resources among adult cancer patients and their companions. *Proc Am Soc Clin Oncol* 2001;20:398a (abstract 1589).
- Helft PR, Hlubocky FJ, Gordon EJ, Ratain MJ, Daugherty CK. Hope and the media in advanced cancer patients. *Proc Am Soc Clin Oncol* 2000;19:633a (abstract 2497).
- O'Connor JB, Johanson JF. Use of the web for medical information by a gastroenterology clinic population. *JAMA* 2000;284:1962-4.
- Health on the Net. Survey on the evolution of internet use for health purposes: raw data for the survey February-March 2001. www.honch/Survey/FebMar2001/ (accessed 14 Jan 2002).
- Silberg WM, Lundberg GD, Musacchio RA. Assessing, controlling, and assuring the quality of medical information on the internet: Caveant lector et viewer—let the reader and viewer beware [editorial] [see comments]. *JAMA* 1997;277:1244-5.
- Jadad AR, Gagliardi A. Rating health information on the internet: navigating to knowledge or to Babel? *JAMA* 1998;279:611-4.
- Bichakjian CK, Schwartz JL, Wang TS, Hall JM, Johnson TM, Sybil Biermann J. Melanoma information on the internet: often incomplete—a public health opportunity? *J Clin Oncol* 2002;20:134-41.
- Price SL, Hersh WR. Filtering web pages for quality indicators: an empirical approach to finding high quality consumer health information on the world wide web. *Proc AMIA Symp* 1999;911-5.
- Shon J, Musen MA. The low availability of metadata elements for evaluating the quality of medical information on the world wide web. *Proc AMIA Symp* 1999;945-9.
- Hoffman-Goetz L, Clarke JN. Quality of breast cancer sites on the world wide web. *Can J Public Health* 2000;91:281-4.
- What is popularity: [thewritemarket.com. www.thewritemarket.com/search/popularity.htm](http://thewritemarket.com/search/popularity.htm) (accessed 14 Mar 2001).
- Kerber R. Direct hit uses popularity to narrow internet searches. *Wall Street Journal* 1998;232 (July 2): B4, Op 1.
- Search engines take a quantum leap: 19 out of 20 now use link popularity to determine relevancy. www.webseed.com/page1007.html (accessed 14 Mar 2001).
- Brin S, Page L. The anatomy of a large-scale hypertextual web search engine. *Computer Networks and ISDN Systems* 1998;30:107-17.
- Linkpopularity.com. www.linkpopularity.com/ (accessed 21 Dec 2001).
- Google. <http://google.com/> (accessed 14 Mar 2001).
- Rumsey E. Peer-review popularity vs. dotcom popularity. www.lib.uiowa.edu (accessed 7 Feb 2002).
- Eysenbach G, Diepgen TL. Towards quality management of medical information on the internet: evaluation, labelling, and filtering of information. *BMJ* 1998;317:1496-500.
- Lacroix E-M. Health topics most hit March 2000. www.nlm.nih.gov/pubs/staffpubs/10/medlineplus/sld013.htm (accessed 27 Jan 2001).
- Bateman M, Rittenberg CN, Gralla RJ. Is the internet a reliable and useful resource for patients and oncology professionals: a randomized evaluation of breast cancer information. *Proc Am Soc Clin Oncol* 1998;17:419a (abstract 1616).
- Berland GK, Elliott MN, Morales LS, Algazy JL, Kravitz RL, Broder MS, et al. Health information on the internet: accessibility, quality, and readability in English and Spanish. *JAMA* 2001;285:2612-21.
- Health On the Net Foundation. HON code of conduct (HONcode) for medical and health web sites: principles. www.hon.ch/HONcode/Conduct.html (accessed 25 May 2001).
- Sandvik H. Health information and interaction on the internet: a survey of female urinary incontinence. *BMJ* 1999;319:29-32.

(Accepted 22 January 2002)

Accuracy of information on apparently credible websites: survey of five common health topics

Heinke Kunst, Diederik Groot, Pallavi M Latthe, Manish Latthe, Khalid S Khan

The internet provides an easily accessible forum to disseminate both accurate and inaccurate health information—so it has the potential to facilitate but also to jeopardise healthcare provision.^{1,2} Many criteria have been alleged to capture the quality of health websites,^{3,4} but the validity of these criteria needs to be examined.⁵ The source, currency, and hierarchy of the evidence posted on a website may be used to judge its credibility—that is, the power of inspiring belief. If these criteria were fulfilled, the contents of the website would be expected to be accurate. We determined whether websites that seem to be credible provide accurate health information.

Methods and results

We determined the relation between credibility features and accuracy of contents of 121 websites that provided information on five common health topics: chronic obstructive pulmonary disease (23 sites), ankle sprain (36), emergency contraception (32), menorrhagia (9), and female sterilisation (21). These sites were identified either by searching each of the most commonly used engines (such as Altavista, Excite, Hotbot, Infoseek, Lycos, Northern Light, Webcrawler) or by simultaneously consulting them using a meta-search engine, Copernic 4.1 (www.copernic.com/). We

selected English language websites whose content provided information about the topics. Website selection and data extractions were performed in duplicate, and agreement between the two assessors was high.

The entire contents of the selected websites were assessed for three credibility features (source, currency, and evidence hierarchy) and accuracy of contents. Source and currency are widely used to assess scientific credibility of a website.^{3,4} The source of medical information is usually regarded as the main criterion for its credibility; sites should display the source of the information clearly. Currency is shown by websites that display the date of the original document or content posting on the internet, and that of any updates. We looked at the hierarchy of evidence posted on each website, examining whether the levels assigned to various pieces of information were related to their validity or methodological quality. This allows users to assess the strength of the recommendations being made. Our assessment showed that 113/121 (93%) websites described source, 59 (49%) currency, and 22 (18%) evidence hierarchy.

Accuracy of website contents was judged against rigorously developed, peer reviewed, and published guidelines for each of the five health topics (see table on bmj.com). The data on accuracy were extracted as a proportion of guideline statements included in the

Department of Respiratory Medicine, St Mary's Hospital, London W2 1NY

Heinke Kunst
specialist registrar

Maastricht University, 6200 MD Maastricht, Netherlands

Diederik Groot
medical student

continued over

BMJ 2002;324:581-2

A table of sources of websites appears on bmj.com