findings suggest that the individual's approach and attitude are particularly important.

Attitude has an important influence on clinical competence and future clinical performance.³ It is beneficial to identify students whose learning style and approach are inappropriate, before they fail. Additional targeted help may be provided for these students.

Contributors: NW had the original idea for the study. Both authors wrote the paper. NW is guarantor.

Funding: No additional funding.
Competing interests: None declared.

- 1 Newble DI, Entwistle NJ, Hejka EJ, Jolly BC, Whelan G. Towards the identification of student learning problems: the development of a diagnostic inventory. *Med Educ* 1988;22:518-26.
- 2 Newble DI, Entwistle NJ. Learning styles and approaches: implications for medical education. *Med Educ* 1986;20:162-75.
- 3 Newble DI. Assessing clinical competence at the undergraduate level. *Med Educ* 1992;26:504-11.

(Accepted 6 February 2002)

Drug points

Prolonged erections produced by dihydrocodeine and sildenafil

David Goldmeier, Harpal Lamba, St Mary's Hospital, London W2 1NY

At least 100 million tablets of sildenafil have been prescribed worldwide (Pfizer's data; personal communication). The use of sildenafil has only rarely been associated with priapism,^{1,2} although one such case was also associated with sickle cell trait.² We report two cases of interaction between dihydrocodeine and sildenafil that resulted in prolonged erections.

Case 1—A 49 year old married man presented to our sexual function clinic with erectile dysfunction after a fall in which he sustained thoracic vertebral fractures. His partial paraplegia improved over the course of a year, after which he presented for treatment of erectile dysfunction. He was successfully treated with 100 mg of sildenafil, which produced hard erections that detumescd immediately after intercourse. He later sustained a minor soft tissue injury to his right shoulder, for which he was initially given non-steroidal anti-inflammatory drugs. He continued to experience pain during the treatment, so his family practitioner changed the treatment to dihydrocodeine 30 mg six hourly, which largely controlled the pain. While taking dihydrocodeine he also took 100 mg of sildenafil. Although he had an orgasm and ejaculated, his erection remained hard for five hours and then subsided. Four days later he again took sildenafil while taking dihydrocodeine, after which he had an erection that lasted four hours. He has since stopped taking dihydrocodeine. He continues to use sildenafil 100 mg, which produces erections that subside immediately after orgasm.

Case 2—A 37 year old homosexual man with HIV was being followed up at our unit. His HIV viral load was 4125 RNA copies/ml and his CD4 cell count was 860 cells/mm³ (23%). He was not receiving antiretroviral treatment. The patient was referred to the sexual function clinic, where he was diagnosed as having psychogenic erectile dysfunction. He had recently been treated for moderate depression with cognitive behaviour therapy. While taking 100 mg of sildenafil he had hard erections that subsided after orgasm. He later took dihydrocodeine (30-60 mg six hourly) for a soft tissue injury. During the first seven days of the analgesic treatment he took 100 mg of sildenafil three times, which resulted in a prolonged erection lasting two to three hours after orgasm on each occasion. He continued to take the dihydrocodeine regularly for two more weeks, and the erection he had while taking sildenafil during this time detumescd immediately after orgasm.

The effects of opiates on sexual functioning are inconsistent and variable, but lowered testosterone concentrations, reduced sexual desire, erectile dysfunction, and delayed ejaculation have been frequently reported, particularly among chronic narcotic drug misusers.^{3,4}

The acute opiate intake in our patients on sildenafil produced abnormally high cyclic guanosine monophosphate concentrations in peripheral nerve endings, which resulted in prolonged erections even after orgasm.

The manufacturers have no reports to date of prolonged erections in patients taking acute dose opiates and sildenafil, apart from our two cases. It is important when prescribing sildenafil to ask about current use of opiates, and to warn about such concurrent use in the future.

Funding: None.

Competing interests: DG has received grants from Pfizer and is a consultant to Eli Lilly.

- 1 Sur RL, Kane CJ. Sildenafil citrate-associated priapism. *Urology* 2000; 55:950.
- 2 Kassim AA, Fabry ME, Nagel RL. Acute priapism associated with the use of sildenafil in a patient with sickle cell trait. *Blood* 2000;95:1878-9.
- 3 Gulliford S. Opioid-induced sexual dysfunction. *J Pharm Care Pain Symptom Control* 1998;6:67-74.
- 4 Mirin SM, Meyer RE, Mendelson JH, Ellingboe J. Opiate use and sexual function. *Am J Psychiatry* 1980;137:909-15.

Corrections and clarifications

Male circumcision linked to lower rates of cervical cancer

An editing slip occurred in this news article by Fred Charatan (27 April, p 994), in which we said that researchers tested for the presence of penile human papillomavirus DNA in 1520 men in the study, and found it in 1139 (75%). It should have said that the researchers got a valid result in that number of subjects. The virus was not found in that number.

Homoeopathy use in NHS not justified

In this news article by Mark Hunter (9 March, p 565), we said that a report from the NHS Centre for Reviews and Dissemination, York, maintained that there was no evidence to justify the use of homoeopathy in the NHS. We should have said that the report found that there was insufficient evidence to warrant a significant change in the current provision of homoeopathy in the NHS.

Driving and arrhythmias

The Driver and Vehicle Licensing Agency (DVLA) has asked us to update a point in this editorial by Helen Binns and John Camm (20 April, pp 927-8). The current standard requires only a two year (not five year, as stated in the article) moratorium on driving after an incapacitating event in relation to implanted cardioverter defibrillators. This new advice was published in December 2001 both in the panel minutes of the November panel meeting and in the DVLA handbook, *At a Glance Guide to the Current Medical Standards of Fitness to Drive*. Both these documents are available on the DVLA's website (www.dvla.gov.uk).