Are breast cancer screening programmes increasing rates of mastectomy? Observational study

Eugenio Paci, Stephen W Duffy, Daniela Giorgi, Marco Zappa, Emanuele Crocetti, Vania Vezzosi, Simonetta Bianchi, Luigi Cataliotti, Marco Rosselli del Turco

Unit of
Epidemiology,
Centre for the
Study and
Prevention of
Cancer CSPO, Via
di S Salvi 12, 50135
Florence, Italy

Eugenio Paci
director
Marco Zappa
senior epidemiologist
Emanuele Crocetti
senior epidemiologist

Screening Unit,
Centre for the
Study and
Prevention of
Cancer CSPO
Marco Rosselli del
Turco
director

Dept of
Mathematics,
Statistics and
Epidemiology,
Imperial Cancer
Research Fund,
London
WC2A 3PX
Stephen W Duffy
principal scientist

Unit of
Epidemiology,
Azienda Sanitaria
Locale (ASL) 2,
55100 Lucca, Italy
Daniela Giorgi
senior epidemiologist

Pathology
Department,
University of
Florence, 50134
Florence, Italy
Vania Vezzosi
pathologist
Simonetta Bianchi
senior pathologist

Surgery
Department,
University of
Florence
Luigi Cataliotti
professor of surgery

Correspondence to:
E Paci
Epid1@cspo.it

BMJ 2002;325:418

In the recent Cochrane review of mammographic screening the authors and editors disagreed with respect to reporting the effect of screening on rates of aggressive treatment for breast cancer.¹ The authors maintained that screening increased the number of mastectomies by around 20%, mainly as a result of overdiagnosis. If this were the case, populations in which screening has been introduced should see a subsequent increase in the number of radical treatments for breast cancer and an increase in the incidence of breast cancer in excess of that expected purely from lead time. We report the changes in rates of radical surgery and incidence of breast cancer since the introduction of the Florence mammographic screening programme.

Methods and results

We studied the period from 1990 to 1996. During this time 59 947 women aged 50-69 years were invited to a prevalence screen and to subsequent screens at two year intervals (2.4 years in practice).² At the beginning of 1990 no women had been invited; by the end of 1994 all the women had been invited to at least one screen. Attendance rates averaged 60%. We considered cases of breast cancer with pathological size T₂ or greater to be late stage. Early stage cases included tumours of less than 2 cm size and in situ carcinomas.

We plotted absolute rates of mastectomy and breast conserving surgery—defined as lumpectomy, wide excision, and quadrantectomy—per thousand women in the population (not per cancer case) against time for the years 1990-6. We assessed the significance of changes over time by trend tests in Poisson regression.

The figure shows the absolute rates of mastectomy and breast conserving surgery over time, with the disease incidence by stage. Rates of breast conserving surgery increased by around 0.7 per thousand (P=0.01) over the period 1990-6, and rates of radical surgery declined by 0.5 per thousand (P<0.001). These rates closely paralleled those of early and late stage breast cancer. Early disease and breast conserving surgery peaked in 1993 as the mobile screening units completed the prevalence screen; this was followed by a reduction in 1994 after the prevalence screen had finished, as one would expect.³ The rates were also strongly related to the amount of screening activity. The peak number of mammograms was 17 298 in 1993, followed by only 8965 in 1994 when the prevalence screen was largely completed but the first repeat screen only just beginning. In 1990 the rate of mastectomy was 1.08 (95% confidence interval 0.84 to 1.37) per thousand, whereas in 1996 it was 0.62 (0.44 to 0.86) per thousand. The rates of breast conserving surgery were 1.18 (0.92 to 1.48) per thousand in 1990 and 1.87 (1.54 to 2.21) per thousand in 1996.

No of mammograms per calendar year

1990	1991	1992	1993	1994	1995	1996
2018	8357	10 297	17 298	8965	16 766	10 774

Size of breast cancers and type of operation

Comment

The rate of breast conserving surgery has increased significantly with the advent of screening, and the rate of radical surgery has declined significantly. Similar reductions in mastectomy rates have been observed elsewhere.⁴ This indicates that the introduction of screening brings about a reduction in mastectomy rates, not an increase. Follow up will continue to ascertain whether these findings are maintained.

Contributors: EP, DG, and MZ did the epidemiological analysis. SWD did the statistical analysis. VV, SB, LC, and MRdelT reviewed and interpreted the pathological, surgical, and radiological data. EP and SWD wrote the paper. All the authors contributed to the final version. EP is the guarantor.

Funding: EP was supported by an American Cancer Society grant. The Florence City Programme is supported as a member of the European Breast Cancer Screening Network.

Competing interests: None declared.

- Olsen O, Gotzsche PC. Cochrane review on screening for breast cancer with mammography. *Lancet* 2001;358:1340-2.
- Paci E, Duffy SW, Giorgi D, Prevost TC, Rosselli del Turco M. Population-based breast cancer screening programmes: estimates of sensitivity, overdiagnosis and early prediction of the benefit. In: Duffy SW, Hill C, Esteve J, eds. *Quantitative methods for the evaluation of cancer screening*. London: Arnold, 2001.
- Day NE, Walter SD. Simplified models of screening for chronic disease: estimation procedures from mass screening programmes. *Biometrics* 1984;43:1-13.
- Wingo PA, Guest JL, McGinnis L, Miller DS, Rodriguez C, Cardinez CJ, et al. Patterns of inpatient surgeries for the top four cancers in the United States, National Hospital Discharge Survey, 1988-95. *Cancer Causes Control* 2000;11:497-512.

(Accepted 17 April 2002)