- 1 Hattersley AT, Tooke JE. The fetal insulin hypothesis: an alternative explanation of the association of low birth weight with diabetes and vascular disease. *Lancet* 1999;353:1789-92.
- 2 Lindsay RS, Dabelea D, Roumain J, Hanson RL, Bennett PH, Knowler WC. Type 2 diabetes and low birth weight: the role of paternal inheritance in the association of low birth weight and diabetes. *Diabetes* 2000;49:445-9.
- 3 Rasmussen F, Davey Smith G, Sterne J, Tynelius P, Leon DA. Birth char-

acteristics of offspring and parental diabetes. *Am J Epidemiol* 2001;153(suppl):S47.

- 4 Pennison EH, Eggerman RS. Perinatal outcomes in gestational diabetes: a comparison of criteria for diagnosis. *Am J Obstet Gynecol* 2001;184:1118-21.
- 5 Butler NR, Bonham DG. *Perinatal mortality*. Edinburgh: Livingstone, 1963.

(Accepted 22 July 2002)

Incidence of coronary heart disease in a health authority in London: review of a community register

Steven J Sutcliffe, Kevin F Fox, David A Wood, Angela Sutcliffe, Kathryn Stock, Melissa Wright, Fawaz Akhras, Ed Langford

Cardiovascular Research Group, National Heart and Lung Institute, Faculty of Medicine, Imperial College of Science, Technology and Medicine, Charing Cross Hospital, London W6 8RF

Steven J Sutcliffe
research fellow
Kevin F Fox
research fellow
David A Wood
professor of cardiovascular medicine

Angela Sutcliffe
research nurse
Kathryn Stock
research nurse
Melissa Wright
statistician

Bromley Hospital, Bromley Hospitals NHS Trust, London BR2 9AQ
Fawaz Akhras
consultant cardiologist
Ed Langford
consultant cardiologist

Correspondence to:
D Wood
d.wood@ic.ac.uk

BMJ 2003;326:20

Strategies to reduce deaths from coronary heart disease need to take into account clinical presentation of the disease.¹⁻⁴ We used a coronary heart register to examine non-fatal and fatal presentations of the disease in a health authority in outer London.

Methods and results

Bromley Health Authority employs 151 general practitioners with a total of 295 584 patients on their lists. The Bromley coronary register prospectively identified all incidences of coronary heart disease in people aged 16-74 from September 1996 to August 1997 for men and from September 1996 to May 1998 for women.

The health authority invited general practitioners to refer new cases of suspected exertional angina to a daily "rapid access" clinic for patients with chest pain at Bromley Hospital. All patients using the clinic and cardiology outpatients were screened. Inpatients admitted to the hospital through its emergency department were screened and followed up on the ward until diagnosis. The two coroner's offices for the authority identified sudden deaths in the community and inpatient deaths that had been reported to them. The authority provided data on all Bromley residents admitted to hospitals in England and Wales with ICD-9 (international classification of diseases, ninth revision) codes for coronary artery disease (410-414). We retrospectively audited a random 8/59 general practices for new prescriptions of nitrates (for angina); we found four missed cases during the registration period. Ethical approval was provided by Bromley local research ethics committee.

A panel of three consultant cardiologists reviewed the patients' presenting history, clinical examination, electrocardiogram, results of blood tests, results of tests measuring stress during exercise, myocardial perfusion scans, and necropsy information where available; they did not review coronary angiograms. The cardiologists classified cases as sudden cardiac death (death attributed

to coronary artery disease), acute myocardial infarction (typical history and resting electrocardiogram changes and creatine kinase concentrations twice the normal upper limit), unstable angina (typical chest pain at rest or rapidly worsening exertional pain without raised creatine kinase concentrations), or exertional angina (typical history and information from exercise testing or perfusion scanning, which was available in 92% of cases) (table 1). To test whether decisions were reproducible, 100 randomly selected cases (about 10% of the total) were reclassified; agreement was good (Cohen's $\kappa=0.86$).

We found 378 cases of men and 242 cases of women who first presented with coronary heart disease, all in the age group 25-74; we found no cases aged 16-24. The incidence of the disease was 414/100 000 population per year in men and 147/100 000 population per year in women. Incidences were greater in older people in both sexes, and at all ages they were significantly higher in men.

Comment

Most coronary heart disease presents in treatable form. Considerable potential exists to reduce the risk of recurrent disease and specifically to prevent patients with angina and well preserved ventricular function from progressing to myocardial infarction and death.⁵ We found that 86% of men and 87% of women present alive to medical services with their first presentation of coronary heart disease; 54% of men and 65% of women have preserved myocardium.

We thank the coroners for their help.

Contributors: SJS ran the clinic for people with chest pain; KFF primarily wrote the paper; DAW designed and supervised the study, reviewed cases, and contributed to the manuscript; KS and AS ran the register; MW did the statistical analysis; and FA and EL reviewed the cases. All authors reviewed and revised the manuscript. DAW is guarantor.

Funding: Preventive Cardiology Trust.

Competing interests: None declared.

- 1 Department of Health. *Saving lives: our healthier nation*. London: Stationery Office, 1999.
- 2 Department of Health. *The national service framework for coronary heart disease*. London: Stationery Office, 2000.
- 3 Gandhi MM, Lampe F, Wood DA. Incidence, clinical characteristics and short-term prognosis of angina pectoris. *Heart* 1995;73:193-8.
- 4 Kannel WB, Feinleib M. Natural history of angina pectoris in the Framingham study. *Am Heart J* 1972;29:154-63.
- 5 Wood DA, Durrington P, Poulter N, McInnes G, Rees A, Wray R on behalf of the British Cardiac Society, British Hyperlipidaemia Association, British Hypertension Society, and British Diabetic Association. Joint British recommendations on prevention of coronary heart disease in clinical practice. *Heart* 1998;80(suppl 2):1-265.

(Accepted 27 July 2002)

Incidence of coronary heart disease per 100 000 population per year in people aged 25-74 in Bromley Health Authority

Presentation	Men		Women	
	Incidence (95% CI)	%	Incidence (95% CI)	%
Exertional angina	172 (146 to 201)	41	77 (63 to 91)	52
Acute myocardial infarction	133 (110 to 159)	32	32 (24 to 42)	22
Sudden cardiac death	57 (43 to 75)	14	19 (13 to 27)	13
Unstable angina	52 (39 to 70)	13	19 (13 to 27)	13