- 7 Schellenberg D, Menendez C, Kahigwa E, Font F, Galindo C, Acosta C, et al. African children with malaria in an area of intense *Plasmodium falciparum* transmission: features on admission to the hospital and risk factors for death. *Am J Trop Med Hyg* 1999;61:431-8.
- 8 Shann F, Barker J, Poore P. Clinical signs that predict death in children with severe pneumonia. *Pediatr Infect Dis J* 1989;8:852-5.
- 9 Sehgal V, Sethi GR, Sachdev HP, Satyanarayana L. Predictors of mortality in subjects hospitalized with acute lower respiratory tract infections. *Indian Pediatr* 1997;34:213-19.
- 10 Bhutta ZA, Nizami SQ, Thobani S, Issani Z. Risk factors for mortality among hospitalized children with persistent diarrhoea in Pakistan. *J Trop Pediatr* 1997;43:330-6.
- 11 Sachdev HP, Kumar S, Singh KK, Satyanarayana L, Puri RK. Risk factors for fatal diarrhea in hospitalized children in India. *J Pediatr Gastroenterol Nutr* 1991;12:76-81.
- 12 Collins S, Myatt M. Short-term prognosis in severe adult and adolescent malnutrition during famine: use of a simple prognostic model based on counting clinical signs. *JAMA* 2000;284:621-6.
- 13 Dramaix M, Brasseur D, Donnen P, Bawhere P, Porignon D, Tonglet R, et al. Prognostic indices for mortality in hospitalized children in central Africa. *Am J Epidemiol* 1996;143:1235-43.
- 14 English M, Murphy S, Mwangi I, Crawley J, Peshu N, Marsh K. Interobserver variation in respiratory signs of severe malaria. *Arch Dis Child* 1995;72:334-6.
- 15 Spiegelhalter DJ, Knill-Jones RP. Statistical and knowledge-based approaches to clinical decision-support systems, with an application in gastroenterology. *J R Stat Soc* 1984;147:35-77.
- 16 Straus SE, McAlister FA, Sackett DL, Deeks JJ. The accuracy of patient history, wheezing, and laryngeal measurements in diagnosing obstructive airway disease. CARE-COAD1 Group. Clinical assessment of the reliability of the examination-chronic obstructive airways disease. *JAMA* 2000;283:1853-7.
- 17 WHO. *Management of the child with a serious infection or severe malnutrition. Guidelines at the first referral level in developing countries*. Geneva: WHO, 2000.

(Accepted 7 November 2002)

Cross sectional survey of meningococcal C immunisation in children looked after by local authorities and those living at home

Catherine M Hill, Mary Mather, Jonathan Goddard

University Child Health, Mail point 803, Southampton General Hospital, Southampton SO16 6YD

Catherine M Hill
senior lecturer in
community child
health

Market Street
Health Centre,
London SE18 6QR

Mary Mather
consultant
paediatrician in
community child
health

Health Care
Research Unit,
University of
Southampton
Jonathan Goddard
statistician

Correspondence to
C M Hill
cmh2@soton.ac.uk

BMJ 2003;326:364-5

Children under the care of a local authority by voluntary agreement or by order of a court often have poor physical and mental health.¹ On 31 March 2001, 75 601 children in England, Scotland, and Wales were looked after by local authorities. In 1998, the government invested £885m in social services through the Quality Protects programme² (Children First in Wales) to improve public care. Compliance with immunisation schedules is an important health performance indicator of these programmes, although lack of immunisation may reflect neglect before, rather than after, a child enters public care. The national meningococcal C vaccine campaign³ in 1999 provided a unique opportunity to measure immunisation uptake in public care, unbiased by historical health neglect. We compared immunisation rates in children looked after by local authorities with that in children at home.

Participants, methods, and results

Nine health districts participated in the study: Southampton; Greenwich; Basildon; Shropshire, Telford, and

Wrekin; Stoke-on Trent; South Tyneside; Calderdale and Huddersfield; Edinburgh; and Cardiff and the Vale of Glamorgan. These were selected because they were the districts of members of a national advisory group. We ascertained the meningococcal C immunisation status of all children on 31 March 2001 from district immunisation databases. We collected data using standardised forms by age (0-4, 5-9, 10-15, 16-18 years) and immunisation category (immunised, not immunised, or unknown). Babies under 2 months were classified as appropriately immunised. We identified children looked after by local authorities from social services records and subtracted the number from the total population by age category to determine the number of children at home. Immunisation status was unknown for 6.5% of children in public care. We excluded missing data from the denominator to calculate unbiased vaccination failure rates. Data were analysed using SAS software (version 8).

Overall, 995/3028 (33%) of children in public care did not receive meningococcal C vaccine compared with 72 841/501 516 (15%) of children at home. Uptake decreased with age in both groups. Children in public

Percentages of children in public care and children living at home who were not immunised against meningitis C in nine health districts and risk ratios for not being immunised. Numbers for the percentages are available on bmj.com

District	Age 0-4 years*			Age 5-9 years*			Age 10-15 years†			Age 16-18 years			All ages		
	% in public care	% at home	Risk ratio	% in public care	% at home	Risk ratio	% in public care	% at home	Risk ratio	% in public care	% at home	Risk ratio	% in public care	% at home	Risk ratio
A	14	8	1.82	29	11	2.66	32	11	2.76	48	26	1.85	31	12	2.58
B	31	12	2.48	30	14	2.13	35	20	1.76	34	36	0.95	33	19	1.76
C	30	30	0.98	53	13	4.12	59	9	6.40	71	9	7.66	53	13	4.17
D	6	11	0.54	18	5	3.67	31	5	5.92	41	17	2.45	24	8	2.91
E	6	12	0.53	—	—	—	16	7	2.30	37	11	3.34	17	10	1.76
F	—	—	—	68	30	2.27	48	38	1.27	64	46	1.39	57	37	1.56
G	—	—	—	7	11	0.63	22	7	3.11	49	20	2.45	22	10	2.17
H	—	—	—	—	—	—	37	11	3.28	—	—	—	37	11	3.28
I	—	—	—	—	—	—	35	13	2.68	—	—	—	35	13	2.68
All	22	12	1.79	31	14	2.28	34	14	2.40	46	22	2.11	33	15	2.26

Risk ratios represent the proportion of children in public care not immunised divided by the proportion of children at home not immunised. A risk ratio >1 indicates that children in public care are less likely to have been immunised than children at home.

*Data were not available for ages 0-4 years in districts F-I nor for ages 5-9 years in district E.

†Aggregated data for all school aged children in districts H and I.

Full data for the
table are available
on bmj.com

care were less likely to be vaccinated than those at home unless they were under 5 years of age in districts C, D, and E or aged 5-9 in district G (table). Retrieving data was difficult in all but one district, which had electronic records of looked after children. No district had a joint health and social services database.

Comment

Overall, children looked after by local authorities were more than twice as likely to not receive meningococcal C vaccine than children at home (risk ratio 2.17 adjusting for age and district, 95% confidence interval 2.06 to 2.28, Mantel-Haenszel method). Because universal childhood meningococcal C vaccination was introduced recently, we were able to study the effectiveness of public care without bias from historical health neglect. Although sampling was opportunistic, our findings are likely to be generalisable as immunisation indices for children in public care in the districts surveyed all fell within one standard deviation of the national mean.⁴

We did not examine the reasons for failure to immunise. However, during 2001, 16% of children in public care moved placement more than three times.⁴ This instability creates potent risk factors, including missed school based immunisation and discontinuity of primary care. The reversed risk ratio for young children in some districts may reflect greater stability in their placements, primary care organisation of the pre-

school campaign, or targeting of vulnerable children by health visitors.⁵

We suggest two ways forward. Firstly, health services should be made accountable for immunisation uptake as well as social services. Secondly, effective shared information systems between health and social services need to be introduced. Together these measures would better protect our most vulnerable children from disease.

We thank Kath Burton, Sue Daniel, Anne Grant, Karen Lehner, Helen Palmer, Carolyn Sampeys, and Peter Soe-Wynn from the medical group of the British agencies of adoption and fostering for help with collecting data.

Contributors: MM and CMH conceived the project and coauthored the paper. CMH organised the data collection. JG analysed and presented the data. All authors approved the final manuscript. CMH is the guarantor.

Funding: None.

Competing interests: None declared.

- 1 House of Commons Health Committee. *Children looked after by the local authority*. London: Stationery Office, 1998.
- 2 Department of Health. Quality protects programme. www.doh.gov.uk/qualityprotects/index.htm (accessed 10 May 2002).
- 3 Department of Health. *Start of the new meningococcal C conjugate vaccine immunisation programme*. London: DoH, 1999. (PL/CMO/99/4.)
- 4 Department of Health. Outcome indicators for looked after children, 12 months to 30 September 2001, England. www.doh.gov.uk/public/oi2001.htm (accessed 18 December 2002).
- 5 Department of Health. *Making a difference. Strengthening the nursing, midwifery and health visiting contribution to health and healthcare*. London: DoH, 1999.

(Accepted 16 October 2002)

Herd immunity from meningococcal serogroup C conjugate vaccination in England: database analysis

Mary E Ramsay, Nick J Andrews, Caroline L Trotter, Edward B Kaczmarski, Elizabeth Miller

In November 1999, the United Kingdom introduced routine meningococcal serogroup C conjugate vaccination for infants. The vaccine was also offered to everyone aged under 18 years in a phased catch-up programme.¹ The first to be vaccinated were adolescents, and the entire programme was completed by the end of 2000. On the basis of direct protection provided by the vaccine,^{1 2} this catch-up programme was likely to be cost effective.³

Maiden et al described a 67% reduction (from 0.45% to 0.15%) in the prevalence of nasopharyngeal carriage of serogroup C meningococci in adolescents before and after the vaccination programme.⁴ A fall in meningococcal carriage would be expected to reduce exposure among unvaccinated children and therefore to enhance the effectiveness of meningococcal conjugate vaccine. We present rates of disease in vaccinated and unvaccinated children to provide the first evidence of an indirect effect from meningococcal conjugate vaccine.

Methods and results

Since December 1999 we have investigated the vaccination history of all cases of serogroup C disease confirmed by the meningococcal reference unit of the Public Health Laboratory Service in age groups

targeted for immunisation.¹ We collected data on vaccination coverage from immunisation coordinators and departments of child health in England.⁵ Between 1 July 2001 and 30 June 2002, we identified a total of 37 cases in the cohorts targeted for catch-up vaccination, eight (22%) in vaccinated children and 29 (78%) in unvaccinated children. We compared cases in unvaccinated children from each age group in the 2001-2 cohort with those in the same age groups for the period from 1 July 1998 to 30 June 1999. The denominator was mid-1999 population estimates from the Office for National Statistics for the age group, adjusted for the proportion of each cohort vaccinated.

Overall, in the age groups targeted for catch-up vaccination, a reduction of 67% (95% confidence interval 52% to 77%) in the attack rate occurred, with a range of 48% to 80% across the age groups (table). A smaller fall occurred in adults not eligible for vaccination (aged ≥ 25 years), for whom the incidence declined by 35% (20% to 49%) from 0.53 (193/36 315 726) to 0.34 (123/36 315 726) per 100 000.

Because of possible underestimation of coverage⁵ we recalculated the attack rates, assuming that coverage was 10% higher in adolescents and 5% higher in other children (based on our experience with other vaccines). The resulting overall estimate was of a 52% reduction (95% confidence interval 30% to 77%) in

Public Health Laboratory Service Communicable Disease Surveillance Centre, London, NW9 5EQ

Mary E Ramsay
consultant
epidemiologist

Nick J Andrews
statistician

Caroline L Trotter
research scientist

Elizabeth Miller
head of immunisation
division

Public Health Laboratory Service Meningococcal Reference Unit, Manchester Public Health Laboratory, Manchester M20 2LR
Edward B Kaczmarski
consultant
microbiologist

Correspondence to: M E Ramsay
mramsay@phls.org.uk

BMJ 2003;326:365-6