

Summary points

Opportunistic disease prevention and health promotion are regarded as part of good primary health care

The number of relevant preventive measures has increased greatly over the past two decades

Decisions about preventive measures need proper discussion about both benefits and harms, which takes time

An extensive preventive agenda may divert the dialogue between patient and doctor away from important social and relational issues relevant to the patient's health

Routine opportunistic preventive initiatives may no longer be ethically justifiable in contemporary Western medicine

Future consultations

Clinical inertia in implementation of preventive medical guidelines should not necessarily be taken as a sign of low quality care. It is time to reconsider the extent to which specific, opportunistic initiatives to prevent disease among asymptomatic individuals should remain a core element of everyday consultations in Western medicine. It is certainly good medical practice to identify, emphasise, and support health promoting resources,²⁷ skills, and activities that have a logical link to the patient's reason for coming to see the doctor. Other opportunistic initiatives may also seem appropriate. Doctors could increase patient autonomy by inviting the patient to introduce a topic rather than using a computerised reminding system. An open ended invitation may be one way to proceed. For example, "It could be that you have been considering

other things that might be good for your health? If there is something you would like to discuss, you are welcome."

We thank the general practitioners at Solvangur Health Centre, Iceland, for encouraging us to write this paper and Anne Luise Kirkengen, Magne Nylenna, Pétur Pétursson, Peter Pritchard, Stefan Hjörleifsson and Steinar Westin for constructive comments.

Contributors and sources: The idea to write this paper arose from discussions at the 12th Nordic Congress in General Practice in Norway, September 2002. The content of general practice was the main topic of the congress, and the authors had central roles as congress president (IH), member of the Nordic reference group (JAS), and key-note speaker (LG). LG has worked for several years in academic general practice and is author of a Norwegian continuing medical education textbook. JAS works as a professor and general practitioner. His research was originally on the epidemiology of cardiovascular risk factors, but recently he has become increasingly interested in the topic of medicalisation. IH works as a GP and associate professor. After her PhD research revealed limited adherence to clinical guidelines in preventive medicine among Norwegian GPs, she went on to address the content of general practice and the role of stake-holders who influence the development of the discipline.

Competing interests: None declared.

Corrections and clarifications

Why do children have chronic abdominal pain, and what happens to them when they grow up? Population based cohort study

An error we made five years ago has just surfaced. In this article by Matthew Hotopf and colleagues (*BMJ* 1998;316:1196-2000), we got a number wrong in the Results section. The final sentence of the first paragraph should start: "Of the risk set, 52 [not 32] were followed up to the age of 36 years."

Sexual health

We muddled the start of the "services" section of the summary box in this editorial by Michael Adler (12 July, pp 62-3). The first two bullet points should have been combined and have read: "Urgent review of staffing requirements and an increase in the number of consultant posts."

Rhabdomyolysis

In converting to *BMJ* style the widely used term "9/11" in this editorial by Russell Lane and Malcolm Phillips (19 July, pp 115-6), we inadvertently referred to the attacks on the World Trade Center in New York as taking place on 9 September 2001. The attacks took place, as we all know, on 11 September.

- 1 Metcalfe D. The crucible. *J R Coll Gen Pract* 1986;36:349-54.
- 2 Stott NCH, Davis RH. The exceptional potential in each primary care consultation. *J R Coll Gen Pract* 1979;29:201-5.
- 3 Sackett DL. The arrogance of preventive medicine. *CMAJ* 2002;167:363-4.
- 4 Skolbekken JA. The risk epidemic in medical journals. *Soc Sci Med* 1995;40:291-305.
- 5 Law MR, Wald NJ. Risk factor thresholds: their existence under scrutiny. *BMJ* 2002;324:1570-6.
- 6 Ault A. Latest US hypertension guidelines create new "pre-hypertensive" category. *Lancet* 2003;361:1798.
- 7 Philips LS, Branch WT, Cook CB, Doyle JP, El-Kebbi IM, Gallina DL, et al. Clinical inertia. *Ann Intern Med* 2001;135:825-34.
- 8 Freeman AC, Sweeney K. Why general practitioners do not implement evidence: a qualitative study. *BMJ* 2001;323:1100-2.
- 9 Hetlevik I. The role of clinical guidelines in cardiovascular risk intervention in general practice. Dissertation. Trondheim: Bjaerum, 1999.
- 10 Yarnall KSH, Pollack KI, Ostbye T, Krause KM, Michener JL. Primary care: is there enough time for prevention? *Am J Pub Health* 2003;93:635-41.
- 11 Edwards A, Elwyn G, Mulley A. Explaining risks: turning numerical data into meaningful pictures. *BMJ* 2002;324:827-30.
- 12 Welsh HG. Informed choice in cancer screening. *JAMA* 2001;285:2776-8.
- 13 Marteau TM, Kinmonth AL. Screening for cardiovascular risk: public health imperative or a matter of individual informed choice? *BMJ* 2002;325:78-80.
- 14 Swensen SJ. Screening for cancer with computed tomography. *BMJ* 2003;326:894-5.
- 15 Misselbrook D, Armstrong D. Patient's responses to risk information about benefits of treating hypertension. *Br J Gen Pract* 2001;51:276-9.
- 16 Kronborg O, Fenger C, Olsen J, Jørgensen OD, Söndergaard O. Randomised study of screening for colorectal cancer with faecal-occult-blood test. *Lancet* 1996;348:1467-71.
- 17 Raffle AE, Alden B, Quinn M, Babb PJ, Brett MT. Outcomes of screening to prevent cancer: analysis of cumulative incidence of cervical abnormality and modelling of cases and deaths prevented. *BMJ* 2003;326:901-4.
- 18 Writing Group for the Women's Health Initiative Investigators. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 2002;288:321-33.
- 19 Getz L, Kirkengen AL. Ultrasound screening in pregnancy: advancing technology, soft markers for fetal chromosomal aberrations, and unacknowledged ethical dilemmas. *Soc Sci Med* 2003;56:2045-57.
- 20 Getz L, Nilsson PM, Hetlevik I. A matter of heart: the general practitioner consultation in an evidence-based world. *Scand J Prim Health Care* 2003;21:3-9.
- 21 Kirkengen AL. *Inscribed bodies: health impact of childhood sexual abuse*. Dordrecht: Kluwer Academic Press, 2001.
- 22 Haynes RB, Devereaux PJ, Guyatt GH. Physicians' and patients' choices in evidence based practice. *BMJ* 2002;324:1350.
- 23 Bracken P. *Trauma: culture, meaning and philosophy*. London: Whurr, 2002.
- 24 Moynihan R, Heath I, Henry D. Selling sickness: the pharmaceutical industry and disease mongering. *BMJ* 2002;324:886-91.
- 25 Smith R. Editor's choice: the screening industry. *BMJ* 2003;326 (26 April).
- 26 Tauber AI. Historical and philosophical reflections on patient autonomy. *Health Care Anal* 2001;9:299-319.
- 27 Hollnagel H, Malterud K. From risk factors to health resources in medical practice. *Med Health Care Philos* 2000;3:257-64.

(Accepted 5 June 2003)