Version of the Glasgow coma scale on neurological observation charts in UK hospitals managing patients with traumatic brain injuries. Values are numbers (percentage)

Version of Glasgow coma scale (points)	Emergency departments (n=233)	Wards for observing patients with traumatic brain injuries (n=233)			Neurosurgical units (n=32)
		Emergency department observation wards (n=58)	Other* (n=175)		
15	178 (76.4)	41 (70.7)	90 (51.4)		18 (56.3)
14	55 (23.6)	17 (29.3)	81 (46.3)		13 (40.6)
Not used	0	0	4 (2.3)		1 (3.1)

*Only about a quarter of all emergency departments have their own observation ward. In most British hospitals, patients with traumatic brain injuries are admitted to general surgical or orthopaedic wards. More rarely, patients are observed on medical, otorhinolaryngological, or neurosurgical wards.

scale in 1976 was not accompanied by an explanation and did not result in a clarifying change of name.

No evidence has been published that the continued use of the 14 point scale may have caused harm to any patient. The practice does, however, lead to difficulties. Staff from one hospital told us about their recurrent problems communicating with the local neurosurgical unit because the two were using different scales. In another trust, the neurological observation charts were changed after a recent coroner's inquest: relatives of a patient who had died of severe traumatic brain injury had raised questions about the logic of two different versions of the Glasgow coma scale being used within the same trust.

Users of the Glasgow coma scale need training to ensure consistency and reliability of scoring.³ The continued employment of two different scales can only add to the confusion. Although the 15 point Glasgow coma scale is not perfect, it should be used by everybody who manages patients with traumatic brain

injuries until even better measures of consciousness are devised.

I thank D Wallis for help with the study design and E Glucksman and P Leman for their comments on the drafts. Many thanks also to all the hospital staff.

Contributors: MFW wrote the paper and is guarantor. Nicola Burger collected some data.

Funding: None.

Competing interests: None declared.

- 1 Diring MN, Edwards DF. Does modification of the Innsbruck and the Glasgow Coma Scales improve their ability to predict functional outcome? *Arch Neurol* 1997;54:606-11.
- 2 Teasdale GM, Jennett B. Assessment of coma and impaired consciousness: a practical scale. *Lancet* 1974;2:81-4.
- 3 Teasdale GM, Jennett B. Assessment and prognosis of coma after head injury. *Acta Neurochir* (Wien) 1976;34:45-55.
- 4 American College of Surgeons Committee on Trauma. *Advanced trauma life support student manual*. 6th ed. Chicago: ACS, 1997.
- 5 Lenfant F, Sobraques P, Nicolas F, Combes JC, Honnart D, Freysz M. Use of Glasgow coma scale by anesthesia and intensive care internists in brain injured patients. *Ann Fr Anesth Reanim* 1997;16:239-43. (In French.)

(Accepted 3 July 2003)

Substance misuse in psychiatric inpatients: comparison of a screening questionnaire survey with case notes

Ben Barnaby, Colin Drummond, Annie McCloud, Tom Burns, Nicola Omu

Doctors are poor at taking alcohol histories in many clinical settings.^{1 2} Given the increasing prevalence of drug misuse in the general population and in psychiatric patients, the extent of detection of drug misuse is also important. For patients with comorbid substance misuse and psychiatric problems, the UK Department of Health states, "Individuals with dual problems deserve high quality, patient focused, and integrated care. This should be delivered within mental health services."³ If substance misuse is not detected, however, such care is unlikely to be delivered. We examined data to investigate the prevalence of alcohol and drug misuse in inpatients admitted to psychiatric wards and the extent and accuracy of detection by the psychiatrists doing assessments on admission.

Participants, methods, and results

We aimed to approach all new admissions to six acute psychiatric wards in two London hospitals over three months. Patients who consented completed the alcohol use disorders identification test (AUDIT) and a short version of the substance abuse assessment questionnaire.⁴ An AUDIT score ≥ 8 indicates hazardous alcohol use and ≥ 15 indicates alcohol dependence.

We excluded patients who did not speak English, who were admitted to the psychiatric intensive care unit, or who were younger than 18 or older than 75. We also excluded patients admitted to specialist wards for addiction or eating disorders and those admitted to forensic wards.

We studied case notes to determine whether a history of substance misuse had been recorded on current admission, its comprehensiveness, whether relevant investigations had been performed, and whether any screening tools had been completed. We considered alcohol and illicit drugs separately. A full history was one including the age of first misuse of each substance, and questions about problems related to using substances and symptoms of dependence, periods of abstinence, and use of specialist services. A partial history contained any one or more of these but less than a full history.

Of a total of 364 new admissions, 326 (89%) were eligible; we approached 263 patients, and 200 (76% of those eligible) gave consent. Not seeing patients was usually because they were admitted and discharged over a weekend or because they absconded.

Of the 200 patients who took part, 106 (53%) were men and 94 (47%) were women. Mean age was 41 years;

Department of Addictive Behaviour and Psychological Medicine, St George's Hospital Medical School, University of London, London SW17 0RE

Ben Barnaby
assistant psychologist
Colin Drummond
professor of addiction psychiatry
Annie McCloud
clinical lecturer in addictive behaviour

Department of Psychiatry, St George's Hospital Medical School
Tom Burns
professor of community psychiatry
continued over

BMJ 2003;327:783-4

South West London
and St George's
Mental Health NHS
Trust, London
SW17 7DJ

Nicola Omu
specialist registrar in
psychiatry

Correspondence to:
C Drummond
colin.drummond@
sghms.ac.uk

Substance misuse and related history in the case notes of 200 patients admitted to psychiatric wards in London*

Measure	No (%)
Results of screening questionnaire	
AUDIT score ≥ 8 (hazardous alcohol use)	97 (49)
AUDIT score ≥ 15 (alcohol dependence)	44 (22)
Lifetime illicit drug misuse†:	
Any	110 (58)
Cannabis	104 (52)
Cocaine or crack	56 (28)
Stimulants	52 (26)
Hallucinogens	52 (26)
Inhalants	38 (19)
Opiates	36 (18)
Illicit drug misuse in previous 30 days‡:	
Any	52 (27)
Cannabis	36 (18)
Cocaine or crack	22 (11)
Stimulants	6 (3)
Hallucinogens	2 (1)
Inhalants	1 (0.5)
Opiates	20 (10)
Review of case notes	
Full alcohol history	1 (0.5)
Partial alcohol history	54 (27)
Liver function test	30 (15)
Assessment tool§	1 (0.5)
Referral to specialist alcohol service	7/44 (16)§
No record of alcohol use	146 (73)
Full drug history	3 (1.5)
Partial drug history	49 (25)
Urine toxicology	36 (18)
No record of drug misuse	148 (74)

*Values for misuse of individual drugs do not sum to value for misuse of any drug because some patients misuse more than one.

†Substance abuse assessment questionnaire.

‡Severity of alcohol dependence questionnaire.

§Seven patients with AUDIT scores ≥ 15 .

158 (79%) were white and 22 (11%) were black. A total of 97 (49%) scored ≥ 8 on AUDIT, indicating hazardous alcohol use—53% of men and 44% of women. Forty four (22%) scored ≥ 15 or above on AUDIT, indicating alcohol dependence. More than half of the patients (110; 58%) reported lifetime substance misuse, and 52 (27%) reported taking illicit drugs in the 30 days before admis-

sion. One patient had had a full alcohol history taken; 54 (27%) had partial alcohol histories in their notes. Three quarters of patients (146; 73%) had no record of using alcohol in their notes. Most patients (148; 74%) had no record of drug misuse in their notes. The ratio of women to men who use alcohol hazardously (1:1.2) was considerably higher than in the general population (1:2.5).⁵

Comment

Substance misuse is common in psychiatric inpatients, but most patients have not been asked about these disorders by admitting psychiatrists. The high prevalence of substance misuse and low screening rates could have an important impact on the quality of the treatment provided, including missed opportunities for substance misuse intervention.

Staff in mainstream mental health services, particularly admitting psychiatrists, urgently need training in detecting and managing comorbid substance misuse. "To overlook or neglect substance misuse in the course of mental health treatment will result in poor treatment outcome."³

Contributors: All authors designed the project and drafted the paper. BB, AMcC, and NO collected the data. BB, CD, and AMcC analysed the data. CD is guarantor.

Funding: No additional funding.

Competing interests: None declared.

Ethical approval: Not needed.

- 1 Farrell MJ, David AS. Do psychiatric registrars take a proper drinking history? *BMJ* 1988;296:395-6.
- 2 Taylor S. Changes in alcohol history taking by psychiatric junior doctors: an audit. *Ir J Psychol Med* 1995;12:37-8.
- 3 Department of Health. *Mental health policy implementation guide: dual diagnosis good practice guide*. London: DoH, 2002. www.doh.gov.uk/mentalhealth/dualdiag.pdf (accessed 5 Aug 2003).
- 4 Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M. Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption II. *Addiction* 1993;88:791-804.
- 5 Singleton N, Bumpstead N, O'Brien M, Lee A, Meltzer H. *Psychiatric morbidity among adults living in private households, 2000*. London: Stationery Office, 2001. www.statistics.gov.uk/downloads/theme_health/psychmorlb.pdf (accessed 5 Aug 2003).

(Accepted 21 July 2003)

Submitting articles to the *BMJ*

We are now inviting all authors who want to submit a paper to the *BMJ* to do so via the web (<http://submit.bmj.com>).

We have introduced Benchpress, our new web based manuscript tracking system, with the aim of streamlining our processes and providing better, quicker information for authors, reviewers, and editors.

Benchpress is a website where authors deposit their manuscripts and editors go to read them and record their decisions. Reviewers' details are also held on the system, and when asked to review a paper reviewers will be invited to access the site to see the relevant paper. The system is secure, protected by passwords, so that authors see only their own papers and reviewers see only those they are meant to. The system is run by Highwire Press, who host bmj.com, and is already being used by 30 journals, including most of the BMJ Publishing Group's specialist journals.

For authors in particular the system offers several benefits. The system provides all our guidance and forms and allows authors to

suggest reviewers for their paper—something we'd like to encourage. Authors get an immediate acknowledgement that their submission has been received, and they can watch the progress of their manuscript. The record of their submission, including editors' and reviewers' reports, remains on the system for future reference.

Anyone with an internet connection and a web browser can use the system.

As with all new systems we expect a few teething problems, but the system itself offers extensive help, and the *BMJ*'s editorial office is geared up to help authors and reviewers if they get stuck. We see Benchpress as part of our endeavour to improve our service to authors and reviewers and, as always, we'd welcome feedback.

Benchpress is accessed via <http://submit.bmj.com> or via a link from bmj.com