- recommendations of the International AIDS Society-USA Panel. *JAMA* 2002;288:222-35.
- 13 Centers for Disease Control and Prevention. 1993 revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *JAMA* 1993;269:460.
 - 14 Juni P, Altman D, Egger M. Assessing the quality of controlled clinical trials. *BMJ* 2001;323:42-6.
 - 15 Song F, Altman DG, Glenny AM, Deeks JJ. Validity of indirect comparison for estimating efficacy of competing interventions: empirical evidence from published meta-analyses. *BMJ* 2003;326:472.
 - 16 Cozzi-Lepri A, Phillips AN, d'Arminio Monforte A, Piersantelli N, Orani A, Petrosillo N, et al. Virologic and immunologic response to regimens containing nevirapine or efavirenz in combination with 2 nucleoside analogues in the Italian Cohort Naïve Antiretrovirals (I.C.O.N.A.) study. *J Infect Dis* 2002;185:1062-9.
 - 17 Matthews GV, Sabin CA, Mandalia S, Lampe F, Phillips AN, Nelson MR, et al. Virological suppression at 6 months is related to choice of initial regimen in antiretroviral-naïve patients: a cohort study. *AIDS* 2002;16:53-61.
 - 18 Phillips AN, Pradier C, Lazzarin A, Clotet B, Goebel FD, Hermans P, et al. Viral load outcome of non-nucleoside reverse transcriptase inhibitor regimens for 2203 mainly antiretroviral-experienced patients. *AIDS* 2001;15:2385-95.
 - 19 Van Leth F, Hassink E, Phanuphak P, Miller S, Gazzard B, Cahn P, et al. Results of the 2NN study: a randomized comparative trial of first-line antiretroviral therapy with regimens containing either nevirapine alone, efavirenz alone or both drugs combined, together with stavudine and lamivudine. 10th conference on retroviruses and opportunistic infections, Boston, MA, 10-14 Feb 2003.
 - 20 Richman DD, Havlir D, Corbeil J, Looney D, Ignacio C, Spector SA, et al. Nevirapine resistance mutations of human immunodeficiency virus type 1 selected during therapy. *J Virol* 1994;68:1660-6.
 - 21 Weidle PJ, Malamba S, Mwebaze R, Sozi C, Rukundo G, Downing R, et al. Assessment of a pilot antiretroviral drug therapy programme in Uganda: patients' response, survival, and drug resistance. *Lancet* 2002;360:34-40.
- (Accepted 6 January 2004)
- doi 10.1136/bmj.37995.435787.A6

Commentary: Indirect comparisons: a novel approach to assessing the effect of anti-HIV drugs

Jens D Lundgren, Andrew N Phillips

The need to evaluate rapidly and provide access to anti-HIV drugs led, in 1997, to an expedited drug approval process, based on short term trials using viral load and CD4 cell counts as surrogate end points for clinical AIDS. The evidence for efficacy of many drugs is therefore based solely on trials using such end points, but it is useful to evaluate studies using clinical end points where available.

Yazdanpanah and coworkers used an indirect comparison of clinical outcomes from randomised controlled trials to compare the effects of drugs from either the protease inhibitor or the non-nucleoside reverse transcriptase inhibitor (NNRTI) class with two nucleoside reverse transcriptase inhibitors (nucleosides).¹ This approach introduces a novel concept to improve further our understanding of the relative efficacy of the two classes. This review suggests a better efficacy of the protease inhibitors than the NNRTIs.

It is important to understand the context of the results to draw conclusions relevant to today. Most of the randomised controlled trials focused on viral end points and were not designed to capture clinical events after virological failure.² Also, many of the drugs from the NNRTI and protease inhibitor classes are considered obsolete, although the NNRTI drug most represented, nevirapine, is still widely used. Furthermore, most trials were based on people with previous exposure to nucleosides and thus likely to harbour virus with resistance to the nucleosides at enrolment. Since the genetic barrier for NNRTIs is lower than for protease inhibitors (with a single nucleotide mutation sufficient to create resistance), it would be predicted that resistance would develop more rapidly with NNRTI based regimens than with protease inhibitor based regimens in this situation, and hence that the clinical outcome would be poorer. The review seems to confirm this prediction, finding little beneficial effect of NNRTIs at all. Importantly, the results for viral load and clinical outcomes are broadly consistent with a better effect of protease inhibitors. In patients starting anti-HIV therapy for the first time, however, several randomised controlled trials with surrogate end points have directly shown that the efficacy of NNRTIs is

comparable and perhaps even superior to protease inhibitors.³⁻⁵

None the less, there are situations in which the findings of Yazdanpanah and coworkers are relevant to today. Frequently, exclusive resistance to nucleosides is seen at failure of current regimens, including triple nucleoside regimens. Furthermore, the WHO has recently launched its 3 by 5 motto of providing anti-HIV therapy to 3 million people by the end of 2005. Hopefully, the therapy will be state of the art, but some may receive inferior regimens of 1-2 nucleosides, increasing the number of patients with resistance to these drugs. It will be critical to start randomised controlled trials with clinical outcomes to establish a rational order of utilisation of the available anti-HIV drugs in this situation. Hopefully, the WHO or other organisations will ensure that this critical knowledge is generated. Yazdanpanah and coworkers provide a strong rationale that this is important.¹

Contributors: JDL wrote the commentary, which was critically revised by ANP.

Funding: None.

Competing interests: JDL has received grant support and fees from Abbott, Bristol-Myers-Squibb, Roche, Boehringer Ingelheim, Pfizer, Merck, and GlaxoSmithKline. ANP has received consultancy fees, grants or funds for meetings from Abbott, Bristol-Myers-Squibb, Roche, Boehringer Ingelheim, and GlaxoSmithKline.

- 1 Yazdanpanah Y, Sissoko D, Egger M, Mouton Y, Zwahlen M, Chêne G. Clinical efficacy of antiretroviral combination therapy based on protease inhibitors or non-nucleoside analogue reverse transcriptase inhibitors: indirect comparison of controlled trials. *BMJ* 2004;328:249-53.
- 2 Kirk O, Pedersen C, Law M, Gulick RM, Moyle G, Montaner J, et al. Analysis of virological efficacy in trials of anti-HIV regimens: drawbacks of not including viral load measurements after premature discontinuation of therapy. *Antivir Ther* 2002;7:271-81.
- 3 Van Leeuwen R, Katlama C, Murphy RL, Squires K, Gatell J, Horban A, et al. A randomized trial to study first-line combination therapy with or without a protease inhibitor in HIV-1-infected patients. *AIDS* 2003;17:987-99.
- 4 Staszewski S, Morales-Ramirez J, Tashima KT, Rachlis A, Skiest D, Stanford J, et al. Efavirenz plus zidovudine and lamivudine, efavirenz plus indinavir, and indinavir plus zidovudine and lamivudine in the treatment of HIV-1 infection in adults. Study 006 Team. *N Engl J Med* 1999;341:1865-73.
- 5 Robbins GK, De Gruttola V, Shafer RW, Smeaton LM, Snyder SW, Pettinelli C, et al. Comparison of sequential three-drug regimens as initial therapy for HIV-1 infection. *N Engl J Med* 2003;349:2293-303.

Copenhagen HIV Programme,
Section 044,
Hvidovre University
Hospital, 2650
Hvidovre, Denmark
Jens D Lundgren
chief physician,
Copenhagen HIV
Programme

Royal Free Centre
for HIV Medicine
and Department of
Primary Care and
Population
Sciences, Royal Free
and University
College Medical
Schools, London
NW3 2PF

Andrew N Phillips
professor of
epidemiology

Correspondence to:
J D Lundgren
jdl@cphiv.dk