

Funding: RMS has received research funding from ECETOC (European Consortium on the Ecotoxicology of Chemicals) and AstraZeneca within the past five years.

Competing interests: RMS and DSI have both received reimbursement of travel expenses for attending meetings sponsored by pharmaceutical companies or the chemical industry.

- Damstra T, Barlow S, Bergman A, Kavlock R, Van Der Kraak G, eds. Global assessment of the state-of-the-science of endocrine disruptors. International Programme on Chemical Safety, 2002. <http://ehp.niehs.nih.gov/who/> (accessed 3 Feb 2004).
- Czene K, Lichtenstein P, Hemminki K. Environmental and heritable causes of cancer among 9.6 million individuals in the Swedish family-cancer database. *Int J Cancer* 2002;99:260-6.
- Jeyaratnam J. Acute pesticide poisoning: a major global health problem. *World Health Stat Q* 1990;43:139-44.
- Second Health and Nutrition Examination Survey (NHANES). *Report on human exposure to environmental chemicals*. Atlanta: National Center for Environmental Health, 2003. (Publication No 02-0716.) www.cdc.gov/nceh/dls/report/ (accessed 3 Feb 2003).
- Mylchreest E, Sar M, Cattle R, Foster PM. Disruption of androgen-regulated male reproductive development by di(n-butyl) phthalate during late gestation in rats is different from flutamide. *Toxicol Appl Pharmacol* 1999;156:81-95.
- Blount BC, Silva MJ, Caudill SP, Needham LL, Pirkle JL, Sampson EJ, et al. Levels of seven urinary phthalate metabolites in a human reference population. *Environ Health Perspect* 2000;108:979-82.
- Koch HM, Drexler H, Angerer J. An estimation of the daily intake of di(2-ethylhexyl) phthalate (DEHP) and other phthalates in the general population. *Int J Hyg Environ Health* 2003;206:1-7.
- Surveillance, Epidemiology, and End Results (SEER) Program SEER*Stat Database: Incidence—SEER 9 Regs Public-Use, Nov 2002 Sub (1973-2000), National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released April 2003, based on the November 2002 submission. <http://seer.cancer.gov/> (accessed 3 Feb 2003).
- Huyghe E, Matsuda T, Thonneau P. Increasing incidence of testicular cancer worldwide: a review. *J Urol* 2003;170:5-11.
- Skakkebaek NE, Rajpert-De Meyts E, Main K. Testicular dysgenesis syndrome: an increasingly common developmental disorder with environmental aspects. *Hum Reprod* 2001;16:972-8.
- Calle E, Frumkin H, Henley J, Savitz DA, Thun MJ. Organochlorines and breast cancer risk. *CA Cancer J Clin* 2002;52:301-9.
- N Evans, ed. Environmental health: what is the connection between chemicals and breast cancer. Breast Cancer Fund and Breast Cancer Action, 2003. www.breastcancerfund.org/environment_evidence_main.htm (accessed 3 Feb 2003).
- Rajapakse N, Silva E, Kortenkamp A. Combining xenoestrogens at levels below individual no-observed-effect concentrations dramatically enhances steroid hormone action. *Env Health Perspect* 2002;110:919-921.
- Endogenous Hormones and Breast Cancer Collaborative Group. Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J Natl Cancer Inst* 2002;94:606-16.
- Hardell L, van Bavel B, Lindstrom G, Carlberg M, Dreifaldt AC, Wijkstrom H, et al. Increased concentrations of polychlorinated biphenyls, hexachlorobenzene and chlordanes in mothers to men with testicular cancer. *Env Health Perspect* 2003;111:930-4.
- Fisher JS, Macpherson S, Marchetti N, Sharpe RM. Human 'testicular dysgenesis syndrome': a possible model based on in utero exposure of the rat to dibutyl phthalate. *Hum Reprod* 2003;7:1383-94.
- Sharpe RM, Franks S. Environment, lifestyle and infertility—an inter-generational issue. *Nature Cell Biol* 2002;4(suppl 1):s33-40.
- Saiyed H, Dewan A, Bhatnagar V, Shemov U, Shenoy R, Rajmohan H, et al. Effect of Endosulfan on male reproductive development. *Env Health Perspect* 2003;111:1958-62.
- Swan SH, Kruse RL, Liu F, Barr DB, Drobnis EZ, Redmon JB, et al. Semen quality in relation to biomarkers of pesticide exposure. *Env Health Perspect* 2003;111:1478-84.
- Duty SM, Silva MJ, Barr DB, Brock JW, Ryan L, Chen Z, et al. Phthalate exposure and human semen parameters. *Epidemiology* 2003;14:269-77.
- Cobellis L, Latini G, De Felice C, Razzi S, Paris I, Ruggieri F, et al. High plasma concentrations of di(2-ethylhexyl)-phthalate in women with endometriosis. *Hum Reprod* 2003;18:1512-5.
- Latini G, De Felice C, Giuseppe P, Del Vecchio A, Paris I, Ruggieri F, et al. In utero exposure to di(2-ethylhexyl)-phthalate and human pregnancy duration. *Environ Health Perspect* 2003;111:1783-5.
- Foster W, Chan S, Platt L, Hughes C. Detection of endocrine disrupting chemicals in samples of second trimester human amniotic fluid. *J Clin Endocrinol Metab* 2000;85:2954-7.
- Cohn BA, Cirillo PM, Wolff MS, Schwing PJ, Cohen RD, Sholtz RI, et al. DDT and DDE exposure in mothers and time to pregnancy in daughters. *Lancet* 2003;361:2205-6.
- Storgaard L, Bonde JP, Ernst E, Spano M, Andersen CY, Frydenberg M, et al. Does smoking during pregnancy affect sons' sperm counts? *Epidemiology* 2003;14:278-86.

Lesson of the week

Bradycardia in acute haemorrhage

Ian Thomas, John Dixon

We describe three patients who all underwent elective surgery and had acute haemorrhage after the operation. All three patients had hypotension but did not develop the typically associated sign of tachycardia. The assumption that occult bleeding is always associated with tachycardia is incorrect and may lead to a delay in diagnosis.

The accepted and traditional cardiovascular signs of acute blood loss are tachycardia, hypotension, and poor peripheral perfusion. In a healthy adult, tachycardia is a heart rate of more than 100 beats/min. The maximum heart rate a person can sustain is generally accepted to be 220 minus the patient's age. Tachycardia occurs in the initial stage of a biphasic cardiovascular response, which attempts to maintain cardiac output in the face of hypovolaemia and a decreased stroke volume. Neural and hormonal mechanisms mediate this tachycardia after arterial baroreceptors are stimulated. In some patients with acute haemorrhage, however, this initial tachycardic response is absent; this may lead to confusion or a delay in diagnosis. The phenomenon has been called relative bradycardia,¹⁻⁴ absence of tachycardic response,¹⁻⁵ or paradoxical bradycardia.⁶ We describe three patients for whom

doctors did not see initial tachycardia during acute haemorrhage.

Case reports

Case 1

A 59 year old woman had an elective total abdominal hysterectomy and bilateral salpingo-oophorectomy. Apart from mild asthma, for which she took a ventolin inhaler, she was otherwise fit and well. She was taking no regular drugs. Her preoperative blood pressure was 145/75 mm Hg, pulse rate was 76 beats/min, and haemoglobin concentration was 147 g/l. Surgery was uneventful. In the recovery room, the woman's blood pressure fell to 70/34, 84/42, and 78/48 mm Hg at 2.5, 4, and 5.5 hours after the operation. At each occasion, doctors prescribed a fluid bolus, which increased systolic blood pressure to greater than 100 mm Hg. Her pulse rate never exceeded 70 beats/min (fig 1). Six hours after the operation, her haemoglobin concentration was 55 g/l. During laparotomy, doctors drained 1500 ml of blood from her peritoneal cavity, securing haemostasis. Subsequent recovery was uneventful.

Patients with acute haemorrhage may not exhibit the typically associated initial tachycardia

Department of Anaesthetics, Royal United Hospital, Bath BA1 3NG
Ian Thomas
specialist registrar

Department of Anaesthetics, Weston General Hospital, Weston-Super-Mare BS23 4TQ
John Dixon
consultant

Correspondence to: I Thomas
ThmsIa@aol.com

BMJ 2004;328:451-3




Fig 1 Case 1: pulse rate and blood pressure every 15 minutes after total abdominal hysterectomy and bilateral salpingo-oophorectomy




Fig 2 Case 2: pulse rate and blood pressure every 30 minutes after laparoscopic cholecystectomy

Case 2

A 54 year old woman had an elective laparoscopic cholecystectomy. She was fit and well and taking no regular drugs. Before the operation, her blood pressure was 122/56 mm Hg, pulse rate was 72 beats/min, and haemoglobin concentration was 111 g/l. Surgery was uneventful. Three hours after the operation, she had a blood pressure of 85/40 mm Hg and a pulse rate of 70 beats/min. Despite having fluids replaced, her systolic blood pressure did not exceed 90 mm Hg and her




Fig 3 Case 3: pulse rate and blood pressure every hour after abdominal hysterectomy and bilateral salpingo-oophorectomy

pulse rate remained fewer than 70 beats/min over the next nine hours (fig 2).

At 12 hours after the operation, the patient had increasing pain in her right shoulder and abdomen. Examination showed diffuse abdominal tenderness. Her haemoglobin was 71 g/l. At laparotomy, surgeons removed 600 ml of blood together with large clots. Her bleeding point was a tear at the insertion of the falciform ligament into the liver. Doctors secured haemostasis, and subsequent recovery was uneventful.

Case 3

A 46 year old woman had an elective abdominal hysterectomy and bilateral salpingo-oophorectomy. She was fit and well and taking no regular drugs. Before surgery, her blood pressure was 145/82 mm Hg, pulse rate was 84 beats/min, and haemoglobin concentration was 139 g/l. After uneventful surgery, she was persistently oliguric with a systolic blood pressure of about 100 mm Hg. Doctors gave 4500 ml of crystalloid and colloid over 18 hours, during which the woman's pulse rate remained fewer than 80 beats/min (fig 3). Her haemoglobin concentration was 59 g/l. At laparotomy, surgeons removed 700 ml of blood and removed large clots. Surgeons identified the source of bleeding as venous ooze from the edge of her vagina and secured haemostasis. Transient renal failure and myocardial ischaemia complicated recovery.

Discussion

Patients with acute haemorrhage may not show the expected initial tachycardic response. Clinicians should consider a diagnosis of acute haemorrhage for patients with hypotension and a normal or low heart rate after surgery. The three patients presented to us in a single year; for patients with acute haemorrhage, lack of initial tachycardia may be more common than previously thought.

Experimental evidence and clinical observations show that, without resuscitation, the response of heart rate to a reduction in the volume of circulating blood is usually biphasic. In the first phase, baroreceptor mediated reflex vasoconstriction and cardio-acceleration maintain arterial pressure despite falling cardiac output. This phase is therefore characterised by tachycardia and normotension. The second phase occurs when about a third of the volume of circulating blood has been lost. Sympathetically mediated vasoconstriction and cardiac drive fall abruptly, and cardiac vagal drive increases. This simultaneously reduces mean arterial pressure and heart rate. Early recognition of this vasodepressor-cardioinhibitory reaction to a reduced circulating volume is vital; the associated bradycardia and hypotension precede circulatory collapse. These phases, seen with progressive central hypovolaemia, are reversed during resuscitation of circulatory volume. The distinction between a bradycardia seen in the second phase of a normal response to acute haemorrhage and a relative bradycardia is that patients with relative bradycardia do not exhibit the initial tachycardia in the first phase.

Bradycardic responses to acute haemorrhage need early recognition and prompt replacement of circulatory volume. About a third of patients with traumatic haemorrhage will present with a bradycardia; "normal

heart rate or even bradycardia can be associated with an acute reduction in blood volume.⁷ A retrospective study of patients with isolated extremity trauma or isolated penetrating abdominal trauma found that of the 71 patients presenting with systolic blood pressure of less than 100 mm Hg, 25 (35%) had a pulse rate of fewer than 100 beats/min.⁸ A retrospective study of patients at a trauma centre found that of 750 patients who presented with a systolic blood pressure of less than 90 mm Hg, 217 (28.9%) had a pulse rate of fewer than 90 beats/min.⁹ The authors of the study also suggested that bradycardia was associated with a better prognosis in certain subgroups of patients.

In the face of severe hypovolaemia, the bradycardia seen in the second phase of cardiovascular response to acute haemorrhage allows for a longer time for diastolic ventricular filling leading to an increased stroke volume.¹⁰ This bradycardia may result from the activation of a vagally mediated reflex arc originating from mechanoreceptors in the left ventricle.¹¹ The bradycardic response may also be the result of a parasympathetic reflex mediated via the vagus nerve triggered by blood in the peritoneum.¹² Vagus involvement is suggested by the fact that after cooling or cutting of the vagus nerves after rapid external haemorrhage in cats the fall in blood pressure was not affected, but the bradycardic response was absent.¹⁰ Why some patients do not show initial tachycardia is unknown.

Conclusion

Bradycardia in patients presenting with acute haemorrhage is common, either as part of a normal physiological response or as a relative bradycardia. The

assumption that initial tachycardia is the only haemodynamic response to acute central hypovolaemia is incorrect, and a failure to recognise bradycardia as a sign of acute haemorrhage is potentially dangerous. Increasing circulatory volume in these critically ill patients should not be delayed.

Contributors: IT and JD collected the data. IT wrote the paper. JD reviewed and redrafted the paper. JD is guarantor.

Funding: No additional funding.

Competing interests: None declared.

- 1 Jansen RPS. Relative bradycardia: a sign of acute intraperitoneal bleeding. *Aust NZ Obstet Gynaecol* 1978;18:206-8.
- 2 Secher NH, Sander Jensen K, Werner C, Warberg J, Bie P. Bradycardia during severe but reversible hypovolemic shock in man. *Circ Shock* 1984;14:267-74.
- 3 Adams SL, Greene JS. Absence of a tachycardic response to intraperitoneal haemorrhage. *J Emerg Med* 1986;4:383-9.
- 4 Snyder HS, Dresnick SJ. Lack of tachycardic response to hypotension in penetrating abdominal injuries. *J Emerg Med* 1989;7:335-9.
- 5 Vayer JS, Henderson JV, Bellamy RF, Galper AR. Absence of a tachycardic response to shock in penetrating intraperitoneal injury. *Ann Emerg Med* 1988;17:227-31.
- 6 Barriot P, Riou B. Haemorrhagic shock with paradoxical bradycardia. *Intens Care Med* 1987;13:203-7.
- 7 American College of Surgeons. Advanced Trauma Life Support for Doctors. (ATLS) Instructor Course Manual. Chicago: ACoS, 1997.
- 8 Thompson D, Adams SL, Barrett J. Relative bradycardia in patients with isolated penetrating abdominal trauma and isolated extremity trauma. *Ann Emerg Med* 1990;19:268-75.
- 9 Demetriades D, Chan LS, Bhasin P, Berne TV, Ramicone E, Huicochea F, et al. Relative bradycardia in patients with traumatic hypotension. *J Trauma* 1998;45:534-9.
- 10 Oberg P, Thoren P. Increased activity in vagal cardiac afferents correlated to the appearance of reflex bradycardia during severe haemorrhage in cats. *Acta Physiol Scand* 1970;80(suppl):22A-23A.
- 11 Oberg B, Thoren P. Increased activity in left ventricular receptors during haemorrhage or occlusion of caval veins in the cat. *Acta Physiol Scand* 1972;85:164-3.
- 12 Snyder HS. Lack of a tachycardic response to hypotension with ruptured ectopic pregnancy. *Am J Emerg Med* 1990;8:23-6.

(Accepted 23 July 2003)

A memorable patient

Happiness is ...

It has been observed in these pages that having a parent's health problem referred to a geriatrician is nature's way of telling you to grow up. The call to the psychogeriatrician is often the signal that a cherished relationship has undergone an irrevocable change and that a new order has taken its place as the prelude to an unwelcome full independence.

And so one day the time came to admit that my father's forgetfulness, irrational behaviour, and wandering were no longer just subjects for resigned family humour but were becoming a hazard to him and a nuisance to the local community. Thus, he and I came to sit in the psychogeriatrician's office—all very friendly, with me probably more nervous than anybody. The initial orientation challenges established the severity of Dad's problems—a receptive and expressive dysphasia behind which it was difficult to imagine a functioning intellect. There was a pause (was there any point continuing?), and then a little "comprehension" exercise: "What is happiness?"

Was this fair on a man in such a condition? I sometimes wonder if it wasn't just a gambit in the hope of passing a few minutes that loomed empty, since it's a question that I suspect would have many of us scrabbling for a useful response.

And what have other, working minds contributed? My *Concise Oxford Dictionary* has "happiness" merely as the noun associated with the word "happy," whose own entry seems to consist of synonyms: "lucky, fortunate, contented with one's lot." Those who

work with words have fitted the concept with diverse associations. Here are a few examples: something the pursuit of which is "an inalienable right of man" (Thomas Jefferson, *American Declaration of Independence*); that which when recalled leads to the greatest misery (Dante); the result of an annual income of twenty pounds with expenditure of nineteen pounds nineteen shillings and sixpence (Charles Dickens, *Pickwick Papers*); or simply the effect of "a good bank balance, a good cook, and a good appetite" (Thoreau).

Perhaps a waste of time, then, asking an old man who can't tell you what day of the week it is. But an answer came. "Happiness is a thing, like this," a wave of the hands. "You lose some and then you get some back." And that was that. The rest of the consultation was brief, and the offer of care at a day centre no surprise and, sadly, little help.

But my father's answer has stayed with me. Was it actually worth anything? Was it just random gibberish from an ischaemic temporoparietal zone? Or a final *haiku* from an old man withdrawing from the world, illustrated not with a calligrapher's flourish but with a hand jive? Whatever, he lived another nine months, and it stands as the last (possibly) sensible thing he said. I have always found it useful as a piece of philosophical advice, and I thought I'd share it with you.

Nick Jefferies *consultant anaesthetist, Fulstow*