- 4 Jarvis MJ, Feyerabend C, Bryant A, Hedges B, Primates P. Passive smoking in the home: plasma cotinine concentrations in non-smokers with smoking partners. *Tob Control* 2001;10:368-74.
- 5 Jarvis MJ, Foulds J, Feyerabend C. Exposure to passive smoking among bar staff. *Br J Addict* 1992;87:111-3.
- 6 Jarvis M, Tunstall-Pedoe H, Feyerabend C, Vesey C, Saloojee Y. Biochemical markers of smoke absorption and self reported exposure to passive smoking. *J Epidemiol Community Health* 1984;38:335-9.
- 7 Tunstall-Pedoe H, Brown CA, Woodward M, Tavendale R. Passive smoking by self report and serum cotinine and the prevalence of respiratory and coronary heart disease in the Scottish heart health study. *J Epidemiol Community Health* 1995;49:139-43.
- 8 Walker M, Shaper AG, Lennon L, Whincup PH. Twenty year follow-up of a cohort based in general practices in 24 British towns. *J Public Health Med* 2000;22:479-85.
- 9 Feyerabend C, Russell MA. A rapid gas-liquid chromatographic method for the determination of cotinine and nicotine in biological fluids. *J Pharm Pharmacol* 1990;42:450-2.
- 10 Jarvis MJ, Tunstall-Pedoe H, Feyerabend C, Vesey C, Saloojee Y. Comparison of tests used to distinguish smokers from nonsmokers. *Am J Public Health* 1987;77:1435-8.
- 11 Jarvis MJ, Goddard E, Higgins V, Feyerabend C, Bryant A, Cook DG. Children's exposure to passive smoking in England since the 1980s: cotinine evidence from population surveys. *BMJ* 2000;321:343-5.
- 12 Walker A, O'Brien M, Traynor J, Fox K, Goddard E, Foster K. *Living in Britain: results from the 2001 general household survey*. London: Stationery Office, 2002.
- 13 Sandler DP, Comstock GW, Helsing KJ, Shore DL. Deaths from all causes in non-smokers who lived with smokers. *Am J Public Health* 1989;79:163-7.
- 14 Hole DJ, Gillis CR, Chopra C, Hawthorne VM. Passive smoking and cardiorespiratory health in a general population in the west of Scotland. *BMJ* 1989;299:423-7.

(Accepted 6 May 2004)

doi 10.1136/bmj.38146.427188.55

Acquisition of *Helicobacter pylori* infection after outbreaks of gastroenteritis: prospective cohort survey in institutionalised young people

Rémi Laporte, Philippe Pernes, Pascale Pronni, Frédéric Gottrand, Pascal Vincent

Inserm
E9919-Université
de Lille II JE 2225,
Institut de Biologie
de Lille, France
Rémi Laporte
resident in paediatrics
Pascal Vincent
microbiologist

Centre Antoine de
Saint-Exupéry,
Vendin-le-Vieil,
France
Philippe Pernes
paediatrician
Pascale Pronnier
paediatrician
Clinique de
Pédiatrie, Hôpital
Jeanne de Flandre,
Lille, France
Frédéric Gottrand
professor

Correspondence to:
P Vincent,
Laboratoire de
Bactériologie-Hygiène,
Centre Hospitalier
Régional et
Universitaire,
Hôpital Calmette,
1 Boulevard J
Leclercq, 59037
Lille Cedex, France
p-vincent@chru-lille.fr

BMJ 2004;329:204-5

The exact mode of spread of *Helicobacter pylori* is still unknown. Transmission during transit disorders of the gastrointestinal tract has been suggested, although there is no evidence to date of transmission during outbreaks of gastroenteritis.¹⁻³ We determined whether gastroenteritis in young people infected with *H pylori* can lead to acquisition of the bacterium by peers.

Participants, methods, and results

Our study took place in a French institution for neurologically handicapped children and adolescents. The young people had been institutionalised for several years and resided across five housing sections (A to E). We included all 112 residents in May 2001. *H pylori* infection present at the outset of the study was

detected by using the non-invasive HpSA stool antigen test (Meridian; 91% sensitivity and 93% specificity).⁴ Stool samples were stored and transported at 4°C within 48 hours and then frozen. The residents were monitored for one year. Events and clinical data were recorded daily by nurses. Gastroenteritis was defined as a sudden outbreak of liquid stools in more than two residents concurrently. For each patient with *H pylori* infection, we defined one day of potentially infective diarrhoea or vomiting as that when at least one liquid stool or vomitus was emitted. Stool samples were collected every week for each resident who was free of infection at the study outset, and at the end of the study we compared the last sample with the first sample. When conversion was observed, we identified the oldest positive stool sample collected from that patient during follow up.

The prevalence of *H pylori* infection was high; 47 of the 112 residents (42%) were positive for *H pylori* at the study outset. Seven of the 65 residents who were initially negative for *H pylori* showed conversion during follow up (figure). Five of the seven young people lived in section E, and our records showed that the two other residents had frequent contacts with the infected patients from section E during physiotherapy and entertainment sessions. Vomitus was rare in all sections. The frequency of diarrhoeal stools from the infected patients varied across sections; residents of sections A and B had 475 and 338 person days of potentially infective diarrhoea over the year, respectively. Acute diarrhoea was rare in these sections, and no outbreaks of gastroenteritis were recorded. A lower frequency of diarrhoeal stools was observed in sections C, D, and E; 34, 104, and 164 person days of potentially infective diarrhoea over the year, respectively. The frequency was always low in sections C and D, where no outbreaks of gastroenteritis were

Relation between new cases of *Helicobacter pylori* infection and exposure to infective diarrhoea and vomitus in housing section E

What is already known on this topic

Diarrhoeal stools and vomitus from infected young people can contain viable *Helicobacter pylori*

What this study adds

New cases of *H pylori* infection can occur in cohabiting young people after acute diarrhoea or vomitus in infected peers

observed, whereas in section E, four peaks of potentially infective diarrhoea occurred during outbreaks of gastroenteritis (figure). In all cases, conversion occurred three to 11 weeks after a peak of potentially infective diarrhoea.

Comment

An association was found between gastroenteritis and subsequent acquisition of *H pylori* infection in cohabiting young people. Despite the limitations of the non-invasive test used to determine the presence of *H pylori*, time and space clustering shows that conversions were not observed by chance. This chronological link suggests that, under usual conditions of life, there might be a causal link between outbreaks of gastroenteritis and transmission of *H pylori*.⁵

We thank Marie-Jeanne Capron, Christine Guillemant, and the nursing staff of Centre Antoine de Saint-Exupéry, Vendin-le-Vieil, for help during the study, Danielle Samaille, Institut Pasteur de Lille, for transport of samples, and M Simonet for critical reading of the manuscript.

Contributors: RL performed the literature search, data collection, and biological and epidemiological analysis and interpretation, and drafted the report. PV was the academic supervisor of RL. PV initiated the project, was responsible for funding and design, organised the collection of samples, helped interpret the results, and wrote and revised the report; he will act as guarantor for the paper. PhP and PP were responsible for the care of the patients and management of the institution. FG was involved in the design of the study and advised on medical issues and discrepancies during the study. All authors critically analysed the data and wrote the paper.

Funding: RL received a scholarship from Laboratoire Gallia.

Competing interests: None declared.

Ethical approval: This study was approved by the local ethics committee, Vendin-le-Vieil.

- 1 Thomas JE, Gibson GR, Darboe MK, Dale A, Weaver LT. Isolation of *Helicobacter pylori* from human faeces. *Lancet* 1992;340:1194-5.
- 2 Parsonnet J, Shmueli H, Haggerty T. Fecal and oral shedding of *Helicobacter pylori* from healthy infected adults. *JAMA* 1999;282:2240-5.
- 3 Leung WK, Siu KL, Kwok CK, Chan SY, Sung R, Sung JJ. Isolation of *Helicobacter pylori* from vomitus in children and its implication in gastro-oral transmission. *Am J Gastroenterol* 1999;94:2881-4.
- 4 Gisbert JP, Pajares JM. Diagnosis of *Helicobacter pylori* infection by stool antigen determination: a systematic review. *Am J Gastroenterol* 2001;96:2829-38.
- 5 Hill AB. The environment and disease: association and causation. *Proc R Soc Med* 1965;58:295-300.

(Accepted 16 April 2004)

DRUG POINTS

Anterior uveitis is associated with travoprost

Manjula Kumarasamy, Shrivatsa P Desai

Uveitis and iritis are rare adverse effects of prostaglandin F2 α analogues¹ and are most common with latanoprost.² A 77 year old woman with no previous history of iritis started travoprost eye drops in November 2002 for advanced pseudoexfoliation glaucoma and cataract in both eyes. Two months later, owing to inadequate control of ocular pressure, she had phacotrabeculectomy in the right eye; she had stopped the travoprost the night before. A week later she complained of swelling and discomfort in the operated eye. We advised her to use her topical steroid more often. The eye settled down and pressure stabilised. In August 2003 she had phacotrabeculectomy in her left eye and again stopped travoprost. As intraocular pressure was still high two weeks later, she restarted taking travoprost. After three weeks she had severe iritis (3+ cells and flare) and corneal oedema and began to take timolol. However, intraocular pressure rose again and she chose to restart travoprost. Five days later acute iritis developed in the left eye, which resolved only after stopping travoprost and starting a combination of a β blocker and dorzolamide eyedrops to control the intraocular pressure. The patient's other drugs included bendrofluzide, diltiazem, ramipril, doxazosin, and chlorthalidopexide.

The close temporal association between starting travoprost and the onset of iritis, and recurrence after rechallenge suggests an adverse reaction to travoprost,

scoring eight with the Naranjo algorithm.³ In the only reported case of iritis associated with travoprost,⁴ the patient was not rechallenged with the drug. The manufacturers of travoprost and the Medicines and Healthcare Products Regulatory Agency are aware of only one other case of uveitis associated with travoprost in the United Kingdom. Similar to patients who had adverse reactions to latanoprost, this woman developed iritis after incisional surgery.⁵

Funding: None.

Competing interests: SPD has received partial reimbursement of travel expenses for attending sponsored symposia organized by various drug companies including Pfizer; Alcon; Allergan; and Merck, Sharp, and Dohme.

- 1 British Medical Association, Royal Pharmaceutical Society of Great Britain. *British national formulary*. London: BMA, RPS, 2003: 515 (No 46).
- 2 Warwar RE, Bullock JD, Ballal D. Cystoid macular edema and anterior uveitis associated with latanoprost use: experience and incidence in a retrospective review of 94 patients. *Ophthalmology* 1998;105:263-8.
- 3 Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA, et al. A method for estimating the probability of adverse drug reactions. *Clin Pharmacol Ther* 1981;30:239-45.
- 4 Faulkner WJ, Burk SE. Acute anterior uveitis and corneal edema associated with travoprost. *Arch Ophthalmol* 2003;121:1054-5.
- 5 Fechtner RD, Khouri AS, Zimmerman TJ, Bullock J, Feldman R, Kulkarni P, et al. Anterior uveitis associated with latanoprost. *Am J Ophthalmol* 1998;126:37-41.

(Accepted 2 March 2004)

Department of Ophthalmology, Doncaster Royal Infirmary, Doncaster DN2 5LT
Manjula Kumarasamy
staff grade ophthalmologist
Shrivatsa P Desai
consultant ophthalmologist

Correspondence to: S P Desai
desaisp@hotmail.com

BMJ 2004;329:205