but coverage was incomplete and uptake modest, and the screening ultimately seems to lead to wider inequalities in sexual health. Macleod and colleagues (p 940) invited 19 773 men and women aged 16-39 to participate in postal screening by providing a specimen collected at home.

With coverage of about 70% and uptake of about 30%, the screening found an overall prevalence of just below 3% for men and almost 4% for women. However, prevalence was higher in younger people, especially younger women who were harder to engage in screening.

POEM*

High dose statins lower C reactive protein

Question Is a lower C reactive protein level after aggressive treatment with statins associated with a lower risk of recurrent coronary events?

Synopsis This is a post hoc analysis of a randomised trial that compared atorvastatin (Lipitor) 80 mg with pravastatin (Pravachol) 40 mg in 4162 patients with acute coronary syndrome. Of those, 3745 had low density lipoprotein (LDL) and C reactive protein (CRP) levels measured 30 days after hospital discharge. There was a linear relationship between the reduction in LDL and the risk of recurrent coronary events (cardiovascular death or non-fatal myocardial infarction) and a similar relationship between the reduction in CRP and the risk of recurrent coronary events. The effects of LDL and CRP were statistically independent. Patients were divided into quartiles by their final CRP; the lowest quartile (<0.9 mg/l) had a relative risk of 1.0, compared with 1.5 for those with a CRP level between 0.9 and 4.2 mg/l, and 1.9 for those with a CRP level greater than 4.2 mg/l. The authors did a variety of subgroup analyses. The most relevant are those for patients who wouldn't otherwise be prescribed a statin; that is, those with LDL <70. In this group, the difference in recurrent coronary events was statistically significant, but clinically small: 3.1 v 2.4 events per 100 person years, corresponding to a number needed to treat of 140 per year. Another study in the same issue (*N Engl J Med* 2005;352:29-38) found that reduction in CRP levels by statins was associated with reductions in the extent of lesions as measured by endovascular ultrasonography. However, only a very high dose of statin (atorvastatin 80 mg) lowered the CRP at all.

Bottom line This post hoc analysis provides support to the hypothesis that statins have an effect on secondary prevention of cardiovascular disease that extends beyond their ability to lower low density lipoprotein levels. However, the absolute benefit is very small in patients with a low LDL who otherwise wouldn't be candidates for statins, even in this very high risk population of patients with acute coronary syndrome. We now need prospective randomised trials of primary prevention in patients who wouldn't otherwise be candidates for statins, comparing this approach to modifying inflammation with other candidate drugs. Such a trial (JUPITER) is underway. Until then, and until the balance of benefits and harms of screening for CRP is fully explored, we should not extend this policy of CRP screening to all of our patients, and certainly not to those without known coronary disease.

Level of evidence 2c (see www.infoPOEMs.com/levels.html). "Outcomes" research; ecological studies

Ridker PM, Cannon CP, Morrow D, et al. C-reactive protein levels and outcomes after statin therapy. *N Engl J Med* 2005;352:20-8.

©infoPOEMs 1992-2003 www.infoPOEMs.com/informationmastery.cfm

* Patient-Oriented Evidence that Matters. See editorial (*BMJ* 2002;325:983)

Editor's choice

Preventive medicine makes us miserable

The old adage—prevention is better than cure—is one we have heard so often that it's hard to shift from our minds. It is intuitively powerful. It just seems to make sense. But shift it we must, for it fuels what Iona Heath, in her cogent article this week (p 954), calls "the excessive self confidence of preventive medicine," which is making us ill and miserable.

Paradoxically, says Heath, the more people are exposed to doctors and contemporary health care, including the rhetoric of preventive care, the sicker they seem to feel. Meanwhile the developing world is starved of affordable treatments. Heath's solution? A tax on preventive drugs sold in rich countries to fund treatments in poor countries, helping both sides to a better balance.

More than two thirds of people in the UK now take medicines to treat or prevent ill health or to enhance wellbeing. Heath asks us to consider the opportunity cost even within a developed country of directing vast sums of money to health promotion rather than to providing better care for those who are actually ill. Of course, there's more money to be made from selling interventions for the healthy majority than for the sick minority, and it helps if you can turn a risk factor (high blood pressure, osteopenia) into a disease.

Whether or not preventive medicines work, they bring with them their share of iatrogenic harms. Heath asks, are we sure that the balance sheet of preventive activity offers more good than harm? Attempting to answer this question for women invited to attend mammography breast screening, Barratt and colleagues (p 936) have created a balance sheet of benefits and harms for women in different age groups.

As Paul Taylor explains in his accompanying editorial (p 915), the main potential harm is overdiagnosis—causing unnecessary anxiety and intervention for those who might otherwise have lived and died (of some other cause) without ever having had their cancer detected. Barratt et al estimate that between 2% and 30% of invasive cancers fall into this category. The benefits and harms for mammography are finely balanced. The decision to be screened is, they say, a gamble, with only a small chance of benefit but a great deal at stake. These kinds of data are being used in patient decision aids in Australia and should be tested more widely.

Because it is acted on healthy people, preventive medicine needs even stronger supporting evidence on benefits and harms than therapeutic interventions. And, for a new age, we need new adages. But will they have the same ring to them? Prevention may not be better than cure. A stitch in time may in some cases be unnecessary and even harmful. All better ideas to bmj.com.

Fiona Godlee *editor* (fgodlee@bmj.com)

To receive Editor's choice by email each week subscribe via our website: bmj.com/cgi/customalert