had no role in study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the paper for publication.

Competing interests: None declared.

Ethical approval: The Kenya Medical Research Institute national ethical and scientific review committees approved the study.

- 1 World Health Organization. *Management of the child with a serious infection or severe malnutrition: guidelines at the first referral level in developing countries*. Geneva: WHO, 2000.
- 2 English M, Berkley J, Mwangi I, Mohammed S, Ahmed M, Osier F, et al. Hypothetical performance of syndrome-based management of acute paediatric admissions of children aged more than 60 days in a Kenyan district hospital. *Bull WHO* 2003;81:166-73.
- 3 Mbogo CM, Mwangangi JM, Nzovu J, Gu W, Yan G, Gunter JT, et al. Spatial and temporal heterogeneity of Anopheles mosquitoes and Plasmodium falciparum transmission along the Kenyan coast. *Am J Trop Med Hyg* 2003;68:734-42.
- 4 Ministry of Health, Republic of Kenya. *AIDS in Kenya*. 6th ed. Nairobi: Government of Kenya, 2001.

- 5 Berkley JA, Mwangi I, Mwarumba S, Lowe B, Marsh K, Newton CRJC. Diagnosis of acute bacterial meningitis in children at a district hospital in sub-Saharan Africa. *Lancet* 2001;357:1753-7.
- 6 Falade AG, Mulholland EK, Adegbola RA, Greenwood BM. Bacterial isolates from blood and lung aspirate cultures in Gambian children with lobar pneumonia. *Ann Trop Paediatr* 1997;17:315-9.
- 7 Falade AG, Adegbola RA, Mulholland EK, Greenwood BM. Respiratory rate as a predictor of positive lung aspirates in young Gambian children with lobar pneumonia. *Ann Trop Paediatr* 2001;21:293-7.
- 8 Makani J, Matuja W, Liyombo E, Snow RW, Marsh K, Warrell DA. Admission diagnosis of cerebral malaria in adults in an endemic area of Tanzania: implications and clinical description. *Q J Med* 2003;96:355-62.
- 9 Reyburn H, Mbatia R, Drakeley C, Carneiro I, Mwakasungula E, Mwerinde O, et al. Overdiagnosis of malaria in patients with severe febrile illness in Tanzania: a prospective study. *BMJ* 2004;329:1212.
- 10 Berkley JA, Versteeg AC, Mwangi I, Lowe BS, Newton CR. Indicators of acute bacterial meningitis in children at a rural Kenyan district hospital. *Pediatrics* 2004;114:e713-9.
- 11 Duke T, Micheal A, Mokela D, Wal T, Reeder J. Chloramphenicol or ceftriaxone, or both, as treatment for meningitis in developing countries. *Arch Dis Child* 2003;88:536-9.

(Accepted 22 February 2005)

doi 10.1136/bmj.38408.471991.8F

Opportunity cost of antidepressant prescribing in England: analysis of routine data

Sandra Hollinghurst, David Kessler, Tim J Peters, David Gunnell

Recently, prescribing of antidepressant drugs has increased exceptionally.¹ At the same time, concerns have been raised about the medicalisation of human distress and, more recently, about the safety of antidepressants.¹

Many general practitioners would like to refer patients for psychological treatment, for which there is good evidence of effectiveness,² but are constrained by the lack of NHS therapists. We estimated the opportunity cost of the recent rise in antidepressant prescribing by valuing it in terms of an effective alternative treatment—cognitive behaviour therapy.

Methods and results

We used Department of Health data on the number and cost of antidepressant drugs dispensed in the community in England to quantify the changes between 1991 and 2002. The baseline year (1991) was chosen to ensure a meaningful timescale and to cover a period of consistent approaches to recording. We took population statistics from www.statistics.gov.uk/statbase, and applied an inflation rate of 32% (from www.statistics.gov.uk/rpi) to 1991 costs.

We estimated the number of patients that could have been treated using cognitive behaviour therapy in 2002, had the rise in prescribing not occurred and the associated costs been diverted to psychological treatment and therapists. We costed the time of a clinical psychologist, including supervision (total equivalent £40 168 (\$74 883; €57 738) full time a year).³ We estimated that each therapist could treat six patients a day for 40 weeks a year and that a treatment episode for mild or moderate depression would comprise six sessions.² We did a limited sensitivity analysis assuming that graduate mental health workers (£25 475 a year) rather than psychologists provided treatment and that treatment episodes consisted of 18 sessions in line with

Source: Department of Health

Prescriptions per head and total cost of antidepressants in England, 1991-2002

the National Institute for Clinical Excellence's recommendation for moderate or severe depression.²

Between 1991 and 2002, prescriptions per head for all antidepressants increased 2.8-fold and the total cost (adjusted for inflation) increased by £310m; the increase was almost entirely due to selective serotonin reuptake inhibitors (figure). These costs could have

This article was posted on bmj.com on 18 March 2005: <http://bmj.com/cgi/doi/10.1136/bmj.38377.715799.F7>

Academic Unit of Primary Health Care, Department of Community Based Medicine, University of Bristol, Bristol BS6 6JL
Sandra Hollinghurst
lecturer in health economics

David Kessler
general practitioner research fellow

Tim J Peters
professor of primary care health services research

Department of Social Medicine, University of Bristol, Bristol BS8 2PR

David Gunnell
professor of epidemiology

Correspondence to: S P Hollinghurst
s.p.hollinghurst@bristol.ac.uk

BMJ 2005;330:999-1000

been used to employ 7700 therapists (26 per primary care trust in England) providing 1.54 million treatment courses of six sessions each a year. This estimate increases to 2.43 million if the therapy is delivered by a graduate mental health worker and falls to 0.51 million if the course of treatment is lengthened to 18 sessions.

Comment

Resources associated with higher levels of NHS antidepressant prescribing in England in 2002 compared with 1991 could have been used to deliver cognitive behaviour therapy to 1.54 million patients, more than a third of adults with depression or mixed anxiety depression.⁴ The recent rise in antidepressant prescribing is likely to be due to increased awareness of depression by patients and professionals; reduced side effects associated with newer antidepressants; and the broadening range of indications for which antidepressants are prescribed (for example, panic disorder, seasonal affective disorder, premenstrual syndrome). Despite concern about the dangers of antidepressants,¹ evidence of ineffective and inefficient prescribing,⁵ and the effectiveness of alternative treatments,² drugs are overwhelmingly the mainstay of treatment for depression in general practice. Increases in the pharmacological treatment of depression have not been matched by the development of psychological services of proved effectiveness, which may reflect the absence of a powerful body, equivalent to the pharmaceutical industry, to promote their development and use.

Although cognitive behaviour therapy is relatively expensive and its population cost effectiveness has not been shown, other cheaper alternatives to both antidepressants and psychotherapy—for example, self help and exercise—may be of equal benefit to patients with mild to moderate depression.² Our analysis takes no account of the training costs of psychotherapists but we have also ignored the cumulative cost of drugs incurred in the 11 years. Despite these limitations, the analysis highlights the scale of resources expended in this area and the uncertainty around alternative

What is already known on this topic

The prescribing of antidepressant drugs has risen substantially in the United Kingdom since the early 1990s

Cognitive behaviour therapy is an effective alternative to antidepressant drugs

What this study adds

Opportunity costs indicate that development of psychological therapies is a feasible alternative to antidepressants

treatment for particular groups of patients; the results indicate that there is a clear need for further research to establish the most appropriate balance between drugs and non-pharmacological treatments for depression.

We thank Steve Pilling and Glyn Lewis for helpful comments and suggestions about an earlier draft of this paper.

Contributors: DG had the idea. SH did the analysis and drafted the paper. All authors contributed to the interpretation of the results and the editing of the paper. SH and DG are guarantors. Funding: None.

Competing interests: DG is a member of the Medicines and Healthcare Products Regulatory Agency expert working group on the safety of selective serotonin reuptake inhibitors. He acts as an independent adviser, receiving travel expenses and a small fee for meeting attendance and reading materials in preparation for the meeting.

Ethical approval: Not needed.

- 1 Gunnell D, Ashby D. Antidepressants and suicide: what is the balance of benefit and risk? *BMJ* 2004;329:34-8.
- 2 National Institute for Clinical Excellence. *Management of depression in primary and secondary care*. London: NICE, 2004. www.nice.org.uk/CG023NICEguideline (accessed 28 Feb 2005).
- 3 Netten A, Curtis L. Unit costs of health and social care 2003. Canterbury: Personal Social Services Research Unit, University of Kent, 2003.
- 4 Office for National Statistics. *Psychiatric morbidity among adults living in private households, 2000*. London: Stationery Office, 2001.
- 5 Lawrenson RA, Tyrer F, Newson RB, Farmer RDT. The treatment of depression in UK general practice: selective serotonin reuptake inhibitors and tricyclic antidepressants compared. *J Affective Disord* 2000;59:149-57.

(Accepted 10 January 2005)

doi 10.1136/bmj.38377.715799.F7

National survey of UK emergency endoscopy units

Andrew Douglass, Michael G Bramble, Ian Barrison

Friarage Hospital,
Northallerton
DL6 1JG

Andrew Douglass
consultant
gastroenterologist

The James Cook
University Hospital,
Middlesbrough
TS4 3BW

Michael G Bramble
consultant
gastroenterologist

continued over

Upper gastrointestinal bleeding is a common cause of hospital admission and is accompanied by considerable mortality. For patients to survive, the timing of endoscopy can be critical. Clinical scoring systems identify high risk patients who need prompt endoscopy after appropriate resuscitation.¹ Early endoscopic intervention to prevent rebleeding is effective in high risk patients.² A recent report indicated that patients are still dying as a consequence of delayed endoscopy,³ but no data exist on the provision of emergency endoscopy services in the United Kingdom. As part of a national census of endoscopy training units, we examined the extent of out of hours endoscopy provision, including volume of work and resources used.

Participants, methods, and results

We approached endoscopy units registered with the UK Joint Advisory Group. We developed a questionnaire from the British Society of Gastroenterology working party report,⁴ and distributed it to lead clinicians in 2002. We sent two reminders to centres that failed to reply. We finished collecting data by August 2002. The number of endoscopy rooms was a surrogate marker for the size of the unit. The response rate was 77% (150 centres).

This article was posted on bmj.com on 11 March 2005: <http://bmj.com/cgi/doi/10.1136/bmj.38379.662616.F7>

BMJ 2005;330:1000-1