Effect of supplementation during pregnancy with L-arginine and antioxidant vitamins in medical food on pre-eclampsia in high risk population: randomised controlled trial

Felipe Vadillo-Ortega, professor,¹ Otilia Perichart-Perera, titular researcher,² Salvador Espino, associate professor of obstetrics and gynaecology,² Marco Antonio Avila-Vergara, associate professor of obstetrics and gynaecology,³ Isabel Ibarra, associate professor,⁴ Roberto Ahued, professor of obstetrics and gynaecology,² Myrna Godines, associate professor of obstetrics and gynaecology,¹ Samuel Parry, associate professor and chief of division of maternal-fetal medicine,⁵ George Macones, Mitchell and Elaine Yanow professor and head of obstetrics and gynaecology,^{5,6} Jerome F Strauss, professor of obstetrics and gynaecology and dean of School of Medicine⁷

¹Department of Experimental Medicine, School of Medicine, Universidad Nacional, Autonomía de México, Torre de Investigación, 3er Piso, Ciudad Universitaria, México, D F 04510

²Instituto Nacional de Perinatología Isidro Espinosa de los Reyes, México

³Universidad Autónoma de Sinaloa, México

⁴Instituto de Investigaciones Biomédicas, UNAM, México

⁵Department of Obstetrics and Gynaecology, University of Pennsylvania, Philadelphia, PA, USA

⁶Department of Obstetrics and Gynaecology, Washington University, St Louis, MO, USA

⁷Department of Obstetrics and Gynaecology, Virginia Commonwealth University, Richmond, VA, USA

Correspondence to: F Vadillo-Ortega felipe.vadillo@gmail.com

Cite this as: *BMJ* 2011;342:d2901 doi:10.1136/bmj.d2901

ABSTRACT

Objective To test the hypothesis that a relative deficiency in L-arginine, the substrate for synthesis of the vasodilatory gas nitric oxide, may be associated with the development of pre-eclampsia in a population at high risk.

Design Randomised, blinded, placebo controlled clinical trial.

Setting Tertiary public hospital in Mexico City.

Participants Pregnant women with a history of a previous pregnancy complicated by pre-eclampsia, or pre-eclampsia in a first degree relative, and deemed to be at increased risk of recurrence of the disease were studied from week 14-32 of gestation and followed until delivery.

Interventions Supplementation with a medical food—bars containing L-arginine plus antioxidant vitamins, antioxidant vitamins alone, or placebo—during pregnancy.

Main outcome measure Development of pre-eclampsia/eclampsia.

Results 222 women were allocated to the placebo group, 228 received L-arginine plus antioxidant vitamins, and 222 received antioxidant vitamins alone. Women had 4-8 prenatal visits while receiving the bars. The incidence of pre-eclampsia was reduced significantly ($\chi^2=19.41$; $P<0.001$) in women randomised to L-arginine plus antioxidant vitamins compared with placebo (absolute risk reduction 0.17 (95% confidence interval 0.12 to 0.21). Antioxidant vitamins alone showed an observed benefit, but this effect was not statistically significant compared with placebo ($\chi^2=3.76$; $P=0.052$; absolute risk reduction 0.07, 0.005 to 0.15). L-arginine plus antioxidant vitamins compared with antioxidant vitamins alone resulted in a significant effect ($P=0.004$; absolute risk reduction 0.09, 0.05 to 0.14).

Conclusions Supplementation during pregnancy with a medical food containing L-arginine and antioxidant vitamins reduced the incidence of pre-eclampsia in a

population at high risk of the condition. Antioxidant vitamins alone did not have a protective effect for prevention of pre-eclampsia. Supplementation with L-arginine plus antioxidant vitamins needs to be evaluated in a low risk population to determine the generalisability of the protective effect, and the relative contributions of L-arginine and antioxidant vitamins to the observed effects of the combined treatment need to be determined.

Trial registration Clinical trials NCT00469846.

INTRODUCTION

Pre-eclampsia and eclampsia are among the leading causes of maternal and neonatal morbidity and mortality.¹ Despite growing knowledge of the pathophysiology of pregnancy induced hypertensive disorders, no preventive measures have been shown to be effective.² The underlying cause of pre-eclampsia/eclampsia is thought to be abnormal placentation, characterised by defective invasion of trophoblast cells and remodelling of the uterine vasculature,³ resulting in reduced utero-placental perfusion, which leads to activation of mechanisms promoting maternal vasoconstriction and activation or damage of endothelial cells.⁴ The endothelium is believed to be a primary target of mediators generated by the placenta. Damage is amplified by other factors such as reactive oxygen species.⁵

Nitric oxide is a potent endothelium derived vasodilator,⁶ and defective synthesis of nitric oxide has been documented in pre-eclampsia.⁷ The main site of production of nitric oxide is nitric oxide synthase in endothelial cells, which uses circulating L-arginine as a substrate. Hence, the local availability of this amino acid may be critical to the endothelial adaptive regulatory mechanisms opposing the vasoconstrictors in pre-eclampsia. L-arginine is considered to be a semi-essential amino acid because under increased demands endogenous synthesis is not sufficient to fulfil

requirements.⁸ Moreover, pregnancy has been reported to be a state of relative arginine deficiency,⁷ imposed by the increased formation of nitric oxide, supporting the adaptive vasodilatation of pregnancy, and use of L-arginine by the fetus.⁹ Pre-eclampsia is also associated with increased concentrations of factors that inhibit nitric oxide production. Concentrations of asymmetric dimethyl arginine, a competitive inhibitor of nitric oxide synthase, are raised in women with pre-eclampsia.¹⁰ Concentrations of soluble fms-like tyrosine kinase 1, which antagonises vascular endothelial growth factor dependent activation of nitric oxide synthase, have also been shown to be increased in pre-eclampsia.¹¹ Endoglin, which impairs activation of nitric oxide synthase mediated by transforming growth factor β , is also increased.¹²

In the past, the role of nutrition in the development of pre-eclampsia has been a subject of considerable discussion. Although little evidence exists to show that dietary manipulations can prevent pre-eclampsia, the notion that they might moderate the secondary features of the syndrome remains in favour.¹³ Substantial experimental data in animals and humans indicate that L-arginine could have a beneficial effect on haemodynamics.¹⁴⁻¹⁶ Of particular note, an expanding literature documents that administration of L-arginine improves vascular function in people with atherosclerosis and peripheral vascular disease.¹⁷⁻²⁰ The oral administration of L-arginine to patients with cardiovascular disease has not been associated with any significant adverse side effects. This includes previous reports in the literature of use of L-arginine in pregnant women. Facchinetti et al and Neri et al infused L-arginine into women whose pregnancies were complicated by intrauterine growth retardation and reported reduced myometrial activity.^{21,22} These observations raised the possibility that supplemental L-arginine in the diet could provide a source of substrate for nitric oxide synthesis during pregnancy, which could promote vasodilatation. More clinical studies are needed in this area because we have limited experience in the use of L-arginine and other nitric oxide donors for preventing pre-eclampsia, as stated in the Cochrane review by Meher and Duley.²³ On the other hand, evidence of endothelial damage mediated by reactive oxygen species has been proposed as another mechanism of endothelial damage in pre-eclampsia.²⁴⁻²⁶ Consequently, antioxidants have been proposed as prophylactic agents for pre-eclampsia and several trials with antioxidants including vitamin C and tocopherols in pre-eclampsia have been published.²⁷⁻²⁹ At the time this clinical trial was started, insufficient evidence existed for us to discard the use of antioxidants for prevention of pre-eclampsia. However, recent trials and Cochrane reviews do not support the use of antioxidants for this purpose.^{30,31} We did a three arm clinical trial to test the hypothesis that a combination of L-arginine and antioxidant supplementation would reduce the risk of pre-eclampsia in a high risk population in Mexico City.

METHODS

Study participants

We enrolled pregnant women between 14 and 32 weeks of gestation at high risk of pre-eclampsia who were receiving prenatal care at the Instituto Nacional de Perinatología Isidro Espinosa de los Reyes in Mexico City between January 2001 and December 2005. We included patients at increased risk of pre-eclampsia, which we defined as either a personal history of pre-eclampsia or pre-eclampsia in a first degree relative. Eligible participants agreed to have their prenatal care and delivery at the institution and provide informed consent. We excluded patients with multiple gestation, known major fetal anomalies (as defined by ultrasound studies by the fetal medicine department), diabetes mellitus or gestational diabetes, pre-existing hypertension, pre-existing renal disease, collagen vascular disease, cancer or strong family history of cancer in first degree relatives, and pre-existing maternal disease needing drug treatment. We excluded women with type 2 diabetes, cancer, or a strong family history of cancer because the angiogenic actions of vascular endothelial growth factor are thought to be mediated, in part, by nitric oxide.³² All women were screened for gestational diabetes at week 14 and again at week 24 of gestation, according to the institutional protocol. We decided in advance that if a woman was diagnosed as having gestational diabetes after randomisation she would discontinue taking bars because of the aforementioned safety concerns. However, we included such women in the data analysis. We excluded patients with autoimmune disease because peroxynitrites have been implicated in the pathogenesis of tissue damage in autoimmune disease.

Recruitment and randomisation

Staff of the department of obstetrics reviewed prenatal records to identify patients who might qualify for the study. A history of pre-eclampsia was confirmed by review of the patient's records, and a family history of pre-eclampsia was based on patients' self report. Gestational age was assessed by last menstrual period and confirmed by a first trimester ultrasound evaluation. The protocol for recruitment included an initial visit in which suitability for randomisation was evaluated, an invitation to participate was tendered, a complete obstetric and nutritional history was taken, and written informed consent was obtained. Participants were randomly assigned to receive one of the three treatments. The principal investigator made the assignment centrally after the patient had given informed consent, by using a computer generated code in random size blocks with concealment of allocation by sealed envelopes.

Each participant in the L-arginine plus antioxidant vitamins group received two bars a day. The average consumption of L-arginine in the United States is around 5.4 g a day,⁸ so we decided to supplement at least 100% of this amount in the form of an available medical food (Heart Bars, Nellson Nutraceutical, CA, USA). Two Heart Bars a day deliver 6.6 g of L-arginine

and antioxidant vitamins.³³ Table 1 shows the composition of the bars used in the study. Participants in the antioxidant vitamins alone group received two bars a day devoid of L-arginine but containing antioxidant vitamins. Participants in the placebo group received two placebo bars a day devoid of L-arginine and antioxidant vitamins.

The bars were packaged in similar envelopes that made them indistinguishable by appearance, and they were flavoured such that they had the same taste irrespective of composition. The bars were packed in boxes containing a sufficient supply for five weeks of treatment. Only the principal investigator knew the group codes. We asked participants to consume bars until the day of delivery. A basal sample of blood was taken at the randomisation visit, and protein was measured in a 24 hour urine sample before participants started the consumption of bars.

Clinical follow-up

Participants were scheduled for clinical follow-up every three to four weeks. Each visit from this point included the following: (1) Arterial pressure measurement. Measurements of the arterial pressure were made according to the recommendations of the American Heart Association (AHA) using the auscultatory method with mercury sphygmomanometers.³⁴ All staff collecting arterial pressure measurements were certified by external monitors from the AHA, who made site visits to the hospital. (2) A sample of 10 mL of venous blood was collected for determination of L-arginine concentration. Plasma was separated and maintained at -80°C until analysis. (3) Participants were instructed to obtain a 24 hour urine sample the

day before the next visit. We assessed urine protein by quantification in 24 hour urine samples with an automated method (DiaSys Diagnostic Systems, Holsheim, Germany).³⁵ (4) To measure compliance, we gave each patient a personal diary to record consumption of bars, which they brought to every appointment. We also instructed them to keep the empty envelopes of the consumed bars, which they also brought back at each appointment. (5) We offered nutritional education and assessment of nutritional status during pregnancy to all participants. We instructed them to avoid the consumption of over the counter prenatal antioxidant vitamins and asked about non-study related vitamin ingestion at each visit.

End points

We defined pre-eclampsia as hypertension (systolic blood pressure ≥ 140 mm Hg, diastolic blood pressure ≥ 90 mm Hg, or both) and proteinuria (>300 mg/24 hours) presenting after 20 weeks of gestation in women known to be previously normotensive. We defined eclampsia as non-epileptic convulsions. We defined mild pre-eclampsia as when hypertension and proteinuria were present but no evidence of systemic organ damage was detectable. Severe pre-eclampsia was detected when proteinuria was above 2.0 g/24 hours, blood pressure was $\geq 160/110$ mm Hg, or both. We also assessed several neonatal end points, including preterm birth (born before 37 weeks of gestation), birth weight, small for gestational age (according to institutional charts), and Apgar scores.

We based the sample size calculation on consideration of all three two way comparisons (placebo *v* antioxidant vitamins alone, placebo *v* antioxidant vitamins plus L-arginine, and antioxidant vitamins plus L-arginine *v* antioxidant vitamins alone), assuming a 50% proportional reduction in pre-eclampsia in any group compared with placebo, an α error of 0.05, a β error of 0.2, and a 1:1:1 allocation ratio. We used a Bonferroni correction so that a test specific error (α) of 0.016 (0.05/3) was used for significance testing of all primary two way comparisons. We also used the α error of 0.016 in our sample size estimation. We assumed a prevalence of pre-eclampsia in the placebo group of 30% (on the basis of pilot data from our institution).

L-arginine assays

We measured L-arginine concentrations by high performance liquid chromatography after derivatisation with the o-phthalaldehyde-ethanethiol reagent as described by Lundsjo et al.³⁶ The derivatised samples were separated by using a reversed phase Kingsorb 3μ C18 column (Phenomenex, Torrance, CA) according to the standard method. We used peak areas for quantification of L-arginine concentrations, using an external standard as reference. We report values as μM in plasma. We monitored within assay and between assay coefficients of variation and maintained them under 3%.

Table 1 | Composition of Heart Bar

Compound	Content
Total fat	2.0 g
Saturated fat	0.5 g
Polyunsaturated fat	1.0 g
Monounsaturated fat	0.5 g
Trans fat	0 g
Cholesterol	10 mg
Sodium	65 mg
Potassium	100 mg
Total carbohydrate	19 g
Dietary fibre	1.0 g
Sugars	12.0 g
Protein	9.0 g
L-arginine	3.3 g
Vitamin C	250 mg
Vitamin E	200 UI
Niacin	25 mg
Vitamin B-6	2.0 mg
Vitamin B-12	4.8 μg
Folate	200 μg

Each bar weighed 35 g and contained 534 kJ (130 kcal); bars for placebo group were devoid of L-arginine and antioxidant vitamins; bars for antioxidant vitamins group were devoid of L-arginine.

Table 2 | Baseline characteristics of participants. Values are means (SD) unless stated otherwise

Characteristic	Placebo (n=222)	L-arginine + vitamins (n=228)	Vitamins alone (n=222)	P value
Age (years)	28.2 (5.1)	28.0 (6.1)	27.6 (5.5)	0.47*
Gestational age at enrolment (weeks)	21.1 (4.7)	20.9 (4.6)	21.1 (5.3)	0.84*
Previous gestation with pre-eclampsia/first degree relative with pre-eclampsia	211/11	218/10	209/13	0.65†
Median (range) parity	2 (1-9)	2 (1-6)	2 (1-6)	0.09†
Median (range) vaginal deliveries	0 (0-5)	0 (0-3)	0 (0-6)	0.42†
Median (range) previous caesarean sections	1 (0-2)	1 (0-3)	1 (0-3)	0.48†
Median (range) abortions	0 (0-5)	0 (0-3)	0 (0-2)	0.13†
Systolic blood pressure at enrolment (mm Hg)	105.1 (9.8)	105.4 (8.8)	105.6 (8.8)	0.86*
Diastolic blood pressure at enrolment (mm Hg)	68.3 (9.1)	67.2 (8.7)	67.6 (7.9)	0.40*
Plasma L-arginine (µM)	21.9 (8.5)	22.7 (8.3)	21.9 (9.2)	0.52*
Pre-gestational body mass index	26.8 (4.9)	27.5 (5.3)	26.1 (4.1)	0.28*
Smoked during pregnancy	5	4	6	0.87†

*Analysis of variance.

†Kruskal-Wallis test.

Statistical analysis

The Universidad Autonoma de Sinaloa (MAA-V) and our biostatistics group (SE and MG) did an independent analysis of the data. The analysis was based on an intention to treat principle. We compared baseline measures among the treatment groups by using analysis of variance methods and Kruskal-Wallis tests. We calculated the proportion of participants who developed pre-eclampsia in each treatment group and did an overall test of equivalence in these proportions by using a Pearson χ^2 statistic test. We calculated absolute risk reduction and relative risk, including 95% confidence intervals, and statistical significance.³⁷

RESULTS

We approached 696 women (figure). We excluded 24 women because they failed to meet inclusion criteria (10 patients) or refused to participate (14 women). Six hundred and seventy-two women met our eligibility criteria and agreed to participate. Forty-four (6.5%)

women had a direct relative who had developed pre-eclampsia, and they were equally distributed among the three arms in this study. Six hundred and twenty-eight women had a previous pregnancy complicated by pre-eclampsia. We allocated 222 women to the placebo group, 228 women received the L-arginine plus antioxidant vitamins bars, and 222 women received the bars with antioxidant vitamins alone. The treatment groups were well balanced with regards to baseline characteristics (table 2).

Clinical follow-up was similar among the women included in the analysis. All groups had a median of 5 (range 2-9) visits during the study (Kruskal-Wallis, $P=0.21$). Less than 5% of study visits were missed for all three groups. In terms of compliance, the placebo group consumed a mean of 1.1 (SD 0.5) bars a day, the L-arginine plus antioxidant vitamins group consumed 1.2 (0.4) bars a day, and the vitamins alone group consumed 1.1 (0.43) bars a day (analysis of variance, $P=0.23$). The nutritionists developed several strategies to promote compliance. Briefly, patients were instructed to mix bars with other foods such as yoghurt, gelatine, milk shakes, or fruit.

Reported side effects were significantly more frequent in the group consuming L-arginine plus antioxidant vitamins bars than in the placebo group, including nausea ($P=0.019$; absolute risk reduction 0.05, 95% confidence interval 0.02 to 0.08; relative risk 1.25, 1.04 to 1.51); symptoms of dyspepsia ($P=0.04$; 0.03, 0.01 to 0.06; 1.34, 1.02 to 1.77); dizziness ($P=0.039$; 0.03, 0.01 to 0.05; 1.42, 1.02 to 1.97); palpitations ($P=0.019$; 0.04, 0.01 to 0.07; 1.36, 1.05 to 1.76); and headache ($P=0.01$; 0.06, 0.03 to 0.09; 1.26, 1.06 to 1.51). None of these side effects led to a participant dropping out of the study.

One hundred and twenty-five patients discontinued their assigned treatment. However, we followed them up until the end of pregnancy and included them analytically in the group to which they were randomised. Fifty-six women decided to drop out of the study voluntarily: 18 in the placebo group, 17 in the L-arginine plus antioxidant vitamins group, and 21 in the

Flow diagram of trial

Table 3 | Clinical outcomes

	No (%)*			Absolute risk reduction (95%CI); relative risk (95% CI)*		
	Placebo (n=222)	L-arginine + vitamins (n=228)	Vitamins alone (n= 222)	L-arginine + vitamins v placebo	Vitamins alone v placebo	L-arginine + vitamins v vitamins alone
Primary outcome						
Pre-eclampsia or eclampsia	67 (30)	29 (13)	50 (23)	0.17 (0.12 to 0.21); 0.42 (0.28 to 0.62) (χ^2 : P<0.001)	0.07 (0.005 to 0.15); 0.74 (0.54 to 1.02) (χ^2 : P= 0.052)	0.09 (0.05 to 0.14); 0.56 (0.37 to 0.85) (χ^2 : P=0.004)
Secondary outcomes						
Total preterm delivery	44 (20)	24 (11)	52 (23)	0.09 (0.05 to 0.13); 0.53 (0.33 to 0.84) (χ^2 : P=0.003)	0.03 (-0.02 to 0.09); 1.18 (0.82 to 1.68) (χ^2 : P=0.419)	0.12 (-0.08 to 0.17); 0.44 (0.28 to 0.70) (χ^2 : P<0.001)
Spontaneous preterm delivery	13 (6)	12 (5)	15 (7)	0.005 (-0.03 to 0.02); 0.89 (0.41 to 1.92) (χ^2 : P=0.631)	0.009 (-0.02 to 0.04); 1.15 (0.56 to 2.36) (χ^2 : P=0.845)	0.01 (-0.04 to 0.14); 0.77 (0.37 to 1.62) (χ^2 : P=0.386)
Caesarean section	152 (68.4)	155 (67.9)	148 (66.6)	-0.004 (-0.67 to 0.57); 0.99 (0.87 to 1.12) (χ^2 : P=0.832)	-0.01 (-0.08 to 0.04); 0.97 (0.85 to 1.10) (χ^2 : P= 0.612)	0.01 (-0.04 to 0.75); 1.01 (0.89 to 1.15) (χ^2 : P=0.843)
Mean (SD) gestational age at delivery	38.4 (2.2)	39.0 (1.9)	38.6 (1.9)	0.65 (0.27 to 1.04)† (P=0.001)	0.29 (-0.68 to 0.09)† (P=0.13)	0.35 (-0.7 to 0.004)† (P=0.051)
Sex of baby (F/M)	115/107	117/111	110/112	χ^2 : P=0.993	χ^2 : P=0.704	χ^2 : P=0.780
Mean (SD) Apgar score:						
1 minute	7.8 (1.1)	8.2 (0.8)	8.0 (0.8)	0.44 (0.24 to 0.64)† (P=0.000)	0.25 (0.04 to 0.47)† (P=0.015)	0.18 (0.01 to 0.36)† (P=0.04)
5 minutes	8.7 (0.7)	8.9 (0.3)	8.9 (0.3)	0.21 (0.10 to 0.33)† (P=0.000)	0.16 (0.05 to 0.46)† (P=0.007)	0.05 (0.02 to 0.12)† (P=0.18)
Mean (SD) neonatal weight (g)	3070.2 (470)	2988.6 (506)	2990.7 (451)	81, (-15 to 179)† (P=0.1)	2 (-102 to 98)† (P=0.96)	79 (-17 to 175)† (P=0.1)
Mean (SD) neonatal length (cm)	48.5 (3.0)	47.9 (3.5)	47.8 (2.9)	0.6 (-1.2 to 0.1)† (P=0.10)	0.1 (-0.6 to 0.8)† (P=0.78)	0.7 (0.03 to 1.3)† (P=0.04)

*Unless stated otherwise.

†Mean difference (95% CI).

antioxidant vitamins alone group. Sixty-nine participants discontinued the use of bars because of the development of gestational diabetes: 24 in the placebo group, 23 in the L-arginine plus antioxidant vitamins group, and 22 in the antioxidant vitamins alone group.

Primary end point

Table 3 shows the findings for the primary end point. The incidence of pre-eclampsia/eclampsia was significantly lower ($\chi^2=19.41$; $P<0.001$) in women randomised to L-arginine plus antioxidant vitamins compared with placebo (absolute risk reduction 0.17, 0.12 to 0.21). The number needed to treat (NNT) was 5.73 (95% confidence interval 4.0 to 10.0). Women receiving antioxidant vitamins alone had an observed benefit, but this effect was not statistically significant compared with placebo ($\chi^2=3.76$; $P=0.052$; absolute risk reduction 0.07, 0.005 to 0.15; NNT 13.06, 6 to 200). The incidence of hypertensive disease was also reduced in the L-arginine plus vitamins group compared with the vitamins alone group ($\chi^2=8.16$; $P=0.004$). Adding L-arginine to antioxidant vitamins compared with vitamins alone resulted in a significant effect on absolute risk reduction (0.09, 0.05 to 0.14; NNT 10.20, 6 to 36).

As enrolment took place over a wide interval from 14 to 32 weeks of gestation, we assessed the relation between gestational age at enrolment and development of pre-eclampsia in women who received L-arginine and antioxidant vitamins compared with placebo. Treatment before 24 weeks reduced the incidence of pre-eclampsia (relative risk 0.37, 0.23 to 0.58), whereas treatment after 24 weeks did not (0.64, 0.30 to 1.37). However, when we estimated the interaction term

between treatment effect and time of first treatment, we did not find it to be significant ($B=0.002$; $P=0.96$)

Secondary end points

The overall rate of preterm birth was also reduced in women randomised to L-arginine plus antioxidant vitamins, although this was not because of a reduction in spontaneous preterm births (preterm premature rupture of membranes or preterm labour) (table 3). We found no difference in mean birth weight or in the proportion of small for gestational age infants. Other obstetrical complications, including placental abruption and postpartum haemorrhage, were similar among treatment groups. No maternal deaths occurred. Five neonatal deaths occurred in the study, two in the placebo group and three in the vitamins group. These deaths were associated with extreme prematurity. We found no differences in plasma concentrations of L-arginine between treatment groups at the beginning of the study.

The mean plasma concentration of L-arginine was lower at the first visit in women who later developed pre-eclampsia (20.5, SD 8.3) compared with those who did not develop pre-eclampsia (22.6, SD 8.8) (mean difference 2.01, 95% confidence interval 0.49 to 3.68; $P<0.01$). The figures in the web appendix show the concentrations of L-arginine (fig A) and the diastolic and systolic blood pressures (fig B) in the different treatment groups. L-arginine concentrations were highest in the group receiving bars containing L-arginine, and the diastolic and systolic pressures were also significantly lower after initial treatment and negatively correlated with L-arginine plasma concentrations.

DISCUSSION

We observed a significant reduction in the incidence of pre-eclampsia/eclampsia in pregnant women who consumed bars containing L-arginine plus antioxidant vitamins. The results of this trial support the proposed hypothesis that supplemental L-arginine can reduce the risk of pre-eclampsia. Consistent with the main study outcome, a secondary finding of the study was that the L-arginine plus antioxidant vitamin supplementation resulted in a significant reduction in the risk of indicated preterm birth compared with placebo.

Methodological considerations and strengths

Endothelial nitric oxide synthase is constitutively expressed in endothelial cells, and its activity depends on the level of expression of the enzyme, availability of the substrate, and concentrations of at least three different inhibitors that have been associated with pre-eclampsia. Synthesis of nitric oxide thus depends on the circulating concentration of L-arginine, and adaptive synthesis of nitric oxide in the endothelium is linked to dietary consumption of this amino acid. Our central hypothesis was that impaired vasodilatation in pregnancies complicated by pre-eclampsia could be prevented by supplementation with L-arginine in the diet. The existing literature suggests that L-arginine has direct effects on blood pressure in experimental animal models, normal humans, hypertensive patients, women with pre-eclampsia, and healthy pregnant women.¹⁴⁻²²

Consistent with recent research, antioxidant vitamins alone did not show a statistically significant reduction in the incidence of pre-eclampsia/eclampsia in our study. Adding L-arginine to antioxidant vitamins resulted in significant protection against the development of pre-eclampsia/eclampsia. Our trial was not powered to detect the small difference we found between the group consuming L-arginine plus antioxidant vitamins and the group consuming bars containing only vitamins. An intriguing result is that supplementing antioxidant vitamins had a statistically significant effect on reducing mild pre-eclampsia. This finding is not consistent with the pooled results of the other clinical trials of antioxidant vitamins, which do not suggest a benefit from treatment.³⁰ Recently, the National Institutes of Health's Maternal-Fetal Medicine Units completed a large trial including several institutions in the United States, which reached a similar conclusion.³¹ In addition, detrimental effects of supplemental vitamin C and vitamin E during gestation have been reported.^{30 38 39}

Limitations of study

The intervention used in this study must be evaluated in a population of pregnant women at low risk of pre-eclampsia. This is particularly relevant because we found a high prevalence of recurrence of pre-eclampsia in the studied population. Unfortunately, no information is available from other Mexican centres that could help to identify characteristics associated with this high recurrence rate compared with centres in other

countries that have reported lower recurrence of pre-eclampsia.^{40 41}

At the time we started this clinical trial, no other commercial presentation for L-arginine was available other than supplemented bars. However, new products containing the amino acid L-arginine in a more palatable presentation are emerging. This makes it feasible to propose a population based study of L-arginine supplementation in larger groups of women with different levels of risk for pre-eclampsia. We did not include a study arm that consumed only L-arginine, as the producer could not supply such bars. In addition, we were aware that L-arginine metabolites may produce deleterious free radicals,⁴² so we considered the inclusion of antioxidant vitamins in the same bars to be beneficial. Unfortunately, our study design could not define the relative contributions of L-arginine and antioxidant vitamins to the beneficial effect of the combination on risk of pre-eclampsia. These ingredients may have had additive effects, or the interaction may have been synergistic. Further studies are needed to clarify these relations. Additionally, our trial was powered to detect a 50% reduction in pre-eclampsia and was thus not designed to detect smaller differences in risk reduction between L-arginine plus antioxidant vitamins and antioxidant vitamins alone.

We did not find concerning side effects during the supplementation of bars. However, caution is needed in women with peptic ulcer disease, as administration of L-arginine may worsen their symptoms.

The protective effect for pre-eclampsia may be greatest if L-arginine and antioxidant vitamins are supplemented before 24 weeks of gestation. Although our study design did not allow a rigorous test of time of supplementation and outcome, the above noted trend should be considered in the design of future clinical trials to evaluate the efficacy of L-arginine supplementation.

Study in context of previous studies

Previous attempts to use L-arginine during pregnancy were designed to modify the course of established disease, using the amino acid as a hypotensive agent.⁴³ Staff et al showed no effect of L-arginine on clinical course when used in women with pre-eclampsia beyond 28 weeks of gestation.⁴⁴ This is in concert with our findings, in which women who started treatment beyond 24 weeks of gestation did not seem to benefit from treatment. Another clinical trial showed no benefits of arginine supplementation once the disease was established.⁴⁵ However, preliminary evidence indicates that supplementing L-arginine during pregnancy may reduce the risk of pre-eclampsia. Germain et al disclosed some initial findings in women at high risk of pre-eclampsia who received L-arginine from week 10 of gestation.⁴⁶ A small clinical trial of L-arginine supplementation in women with chronic hypertension that showed some maternal benefit was published after we submitted our work for publication, supporting the potential benefits of this intervention.⁴⁷

WHAT IS ALREADY KNOWN ON THIS TOPIC

Pre-eclampsia has been shown to be associated with defective vasodilatation

Defective synthesis of nitric oxide, a key vasodilator, has been documented in pre-eclampsia

Experimental data in animals and humans indicate that supplementation with L-arginine, the substrate for nitric oxide synthase, has a beneficial effect on arterial pressure

WHAT THIS STUDY ADDS

In a population of women at high risk of pre-eclampsia, dietary supplementation with L-arginine and antioxidant vitamins reduced occurrence of the disease

Supplementation with antioxidant vitamins alone did not reduce occurrence of pre-eclampsia, consistent with previous studies

Further study is needed to determine whether these results can be replicated and to identify whether they are due to L-arginine alone or the combination of L-arginine and antioxidant vitamins

Conclusions

Our observations are consistent with the notion that availability of the substrate for nitric oxide synthesis (L-arginine) prolongs the latency to development of pre-eclampsia in a high risk population of women taking the amino acid supplement in the presence of antioxidant vitamins in a medical food. This relatively simple and low cost intervention may have value in reducing the risk of pre-eclampsia and associated pre-term birth.

We gratefully acknowledge the technical assistance and contributions of Felipe Peraza (UAS), Ricardo Adame, Margie Balas, Jorge Beltran Montoya, Gerardo Buendia, Esther Casanueva, Rosena Furusho, Rocío Gallardo, Antonia Hernandez, Jose Antonio Hernandez, Flor Paredes, Landy Rivera, Sarahi Roldan, and Marcela Serrano (INPerIER). We also thank Adriana Dabaghi, Diana Gutierrez, Gabriela Robles-Gil, Valeria Ortiz, Esther Schiffman, and Araceli Suverza (Universidad Iberoamericana Santa Fe).

Contributors: FV-O obtained funding, acted as study coordinator, and co-wrote the manuscript. OP-P coordinated clinical nutrition follow-up. SE and MG participated in prenatal follow-up and data analysis. MAA-V participated in data analysis and interpretation. It was responsible for laboratory techniques. RA supervised clinical activities. SP was involved in study design. GM was involved in study design and co-wrote the manuscript. JFS was involved in study design, obtained funding, and co-wrote the manuscript. FV-O is the guarantor.

Funding: Funding was provided by the Bill and Melinda Gates Foundation, National Institutes of Health (P60MD002256, UL1RR031990), and Fondo de Salud, CONACYT (MO-303). The funding agencies had no involvement in any phase of this study, including design; collection, analysis, and interpretation of data; and writing and publication of the manuscript.

Competing interests: All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: funding for the study from the Bill and Melinda Gates Foundation, National Institutes of Health, and CONACYT; no relationships with any companies that might have an interest in the submitted work in the previous three years; and no non-financial interests that may be relevant to the submitted work.

Ethical approval: The project was approved by the local institutional review board (IRB authorisation number 212250-02071). All participants gave informed consent.

Data sharing: The original study protocol, the dataset used for the analysis, and the computer code used to produce the results are available from the corresponding author.

- 1 Ronsmans C, Graham W. Maternal mortality: who, when, where, and why. *Lancet* 2006;368:1189-200.
- 2 Meher S, Duley L. Interventions for preventing pre-eclampsia and its consequences: generic protocol. *Cochrane Database Syst Rev* 2005;2:CD005301.

- 3 Mignini LE, Villar J, Khan KS. Mapping the theories of pre-eclampsia: the need for systematic reviews of mechanisms of the disease. *Am J Obstet Gynecol* 2006;194:317-21.
- 4 Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *Lancet* 2005;365:785-99.
- 5 Davidge S. Oxidative stress and altered endothelial cell function in pre-eclampsia. *Semin Reprod Endocrinol* 1998;16:65-73.
- 6 Rees DD, Palmer RM, Moncada S. Role of endothelium-derived nitric oxide in the regulation of blood pressure. *Proc Natl Acad Sci U S A* 1989;86:3375-8.
- 7 Morris N, Eaton BM. Nitric oxide, the endothelium, pregnancy and pre-eclampsia. *Br J Obstet Gynaecol* 1996;103:4-15.
- 8 Visek WJ. Arginine needs, physiological state and usual diets: a reevaluation. *J Nutr* 1986;116:36-46.
- 9 Conrad KP, Joffe GM, Kruszyna H, Kruszyna R, Rochelle LG, Smith RP, et al. Identification of increased nitric oxide biosynthesis during pregnancy in rats. *FASEB J* 1993;7:566-71.
- 10 Savvidou MD, Hingorani AD, Tsikas D, Frolich JC, Vallance P, Nicolaides KH. Endothelial dysfunction and raised plasma concentrations of asymmetric dimethylarginine in pregnant women who subsequently develop pre-eclampsia. *Lancet* 2003;361:1511-7.
- 11 Maynard SE, Min JY, Merchan J, Lim KH, Li J, Mondal S, et al. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in pre-eclampsia. *J Clin Invest* 2003;111:649-58.
- 12 Venkatesha S, Toporsian M, Lam C, Hanai J, Mammoto T, Kim YM, et al. Soluble endoglin contributes to the pathogenesis of pre-eclampsia. *Nature Med* 2006;12:642-9.
- 13 Fraser R. Pre-eclampsia and diet. In: Strain JJ, Caballero B, Sadler MJ, eds *The encyclopedia of human nutrition*. Academic Press, 1998:1620-6.
- 14 Hishikawa K, Nakaki T, Tsuda M, Esumi H, Ohshima H, Suzuki H, et al. Effects of systemic L-arginine administration on hemodynamics and nitric oxide release in man. *Jpn Heart J* 1992;33:41-8.
- 15 Campese VM, Amar M, Anjali C, Medhat T, Wurgraft A. Effects of L-arginine on systemic and renal haemodynamics in salt-sensitive patients with essential hypertension. *J Hum Hypertens* 1997;11:527-32.
- 16 Rosano GMC, Panina G, Cerquetani E, Leonardo F, Pelliccia F, Bonfigli B, et al. L-arginine improves endothelial function in newly diagnosed hypertensives. *J Am Coll Cardiol* 1998;31:262A.
- 17 Creager MA, Gallagher SJ, Gierd XJ, Coleman SM, Dzau VJ, Cooke JP. L-arginine improves endothelium-dependent vasodilation in hypercholesterolemic humans. *J Clin Invest* 1992;90:1248-53.
- 18 Adams MR, McCredie R, Jessup W, Robinson J, Sullivan D, Celermajer DS. Oral L-arginine improves endothelium-dependent dilation and reduces monocyte adhesion to endothelial cells in young men with coronary artery disease. *Atherosclerosis* 1997;129:261-9.
- 19 Rector TS, Bank AJ, Mullen KA, Tschumperlin LK, Sih R, Pillai K, et al. Randomized, double-blind placebo-controlled study of supplemental oral L-arginine in patients with heart failure. *Circulation* 1996;93:2135-41.
- 20 Bode-Boger SM, Boger RH, Galland A, Tsikas D, Frolich JC. L-arginine-induced vasodilatation in healthy humans: pharmacokinetic pharmacodynamic relationship. *Br J Clin Pharmacol* 1998;46:489-97.
- 21 Facchinetti F, Neri I, Genazzani AR. L-arginine infusion reduces preterm uterine contractions. *J Perinat Med* 1996;24:383-5.
- 22 Neri I, Mazza V, Galassi MC, Volpe A, Facchinetti F. Effects of L-arginine on utero-placental circulation in growth-retarded fetuses. *Acta Obstet Gynecol Scand* 1996;75:208-12.
- 23 Meher S, Duley L. Nitric oxide for preventing pre-eclampsia and its complications. *Cochrane Database Syst Rev* 2007;2:CD006490.
- 24 Hubel CA, McLaughlin MK, Evans RW, Hauth BA, Sims CJ, Roberts JM. Fasting serum triglycerides, free fatty acids, and malondialdehyde are increased in pre-eclampsia, are positively correlated and decrease within 48 hours post partum. *Am J Obstet Gynecol* 1996;174:975-82.
- 25 Barden A, Beilin LJ, Ritchie J, Croft KD, Walters BN, Michael CA. Plasma and urinary 8-isoprostane as an indicator of lipid peroxidation in pre-eclampsia and normal pregnancy. *Clin Sci* 1996;91:711-8.
- 26 Sagol S, Ozkinay E, Ozsener S. Impaired antioxidant activity in women with pre-eclampsia. *Int J Obstet Gynecol* 1999;64:121-7.
- 27 Stratta P, Canavese C, Porcu M, Dogliani M, Todros T, Garbo E, et al. Vitamin E supplementation in pre-eclampsia. *Gynecol Obstet Invest* 1994;37:246-9.
- 28 Gulmezoglu AM, Hofmeyr GJ, Oosthuizen MM. Antioxidants in the treatment of severe pre-eclampsia: an explanatory randomised controlled trial. *Br J Obstet Gynaecol* 1997;104:689-96.
- 29 Chappell L, Seed PT, Briley AL, Kelly FJ, Lee R, Hunt BJ, et al. Effect of antioxidants on the occurrence of pre-eclampsia in women at increased risk: a randomized trial. *Lancet* 1999;354:810-6.

- 30 Rumbold A, Duley L, Crowther CA, Haslam RR. Antioxidants for preventing pre-eclampsia. *Cochrane Database Syst Rev* 2008;1:CD004227.
- 31 Roberts JM, Myatt L, Spong CY, Thom EA, Hauth JC, Leveno KJ, et al. Vitamins C and E to prevent complications of pregnancy-associated hypertension. *N Engl J Med* 2010;362:1282-91.
- 32 Papapetropoulos A, Garcia-Cardena G, Madri JA, Sessa WC. Nitric oxide production contributes to the angiogenic properties of vascular endothelial growth factor in human endothelial cells. *J Clin Invest* 1997;100:3131-9.
- 33 Maxwell AJ, Anderson BE, Cooke JP. Nutritional therapy for peripheral arterial disease: a double-blind, placebo-controlled, randomized trial of HeartBar. *Vasc Med* 2000;5:11-9.
- 34 Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al. Recommendations for blood pressure measurement in humans and experimental animals. Part 1: blood pressure measurement in humans: a statement for professionals and public education of the American Heart Association Council on High Blood Pressure Research. *Hypertension* 2005;45:142-61.
- 35 Orsonneau J, Douet P, Massoubre C, Lustenberger P, Bernard S. An improved pirogallol re-molybdate method for determining total urine protein. *Clin Chem* 1989;35:2233-6.
- 36 Lundsjo A, Hagelberg S, Palmer K, Lindblad BS. Amino acid profiles by HPLC after filter paper sampling: "appropriate technology" for monitoring of nutritional status. *Clin Chim Acta* 1990;191:201-10.
- 37 Levine M, Walter S, Lee H, Haines T, Holbrook A, Pharm D, et al. User's guides to the medical literature. IV: how to use an article about harm. *JAMA* 1994;271:1615-9.
- 38 Podmore ID, Griffiths HR, Herbert KE, Mistry N, Mistry P, Lunec J. Vitamin C exhibits pro-oxidant properties. *Nature* 1998;392:559.
- 39 Banerjee S, Chambers A, Campbell S. Is vitamin E a safe prophylaxis for pre-eclampsia? *Am J Obstet Gynecol* 2006;194:1228-33.
- 40 Hnat MD, Sibai BM, Caritis S, Hauht J, Lindheimer MD, MacPherson C, et al. Perinatal outcome in women with recurrent pre-eclampsia compared with women who develop pre-eclampsia as nulliparas *Am J Obstet Gynecol* 2002;186:422-6.
- 41 Hjartardottir S, Leifsson B, Geirsson R, Steinhorsdottir V. Recurrence of hypertensive disorder in second pregnancy. *Am J Obstet Gynecol* 2006;194:916-20.
- 42 Rubbo H, Radi R, Trujillo M, Telleri R, Kalyanaraman B, Barnes S, et al. Nitric oxide regulation of superoxide and peroxynitrite-dependent lipid peroxidation: formation of novel nitrogen-containing oxidized lipid derivatives. *J Biol Chem* 1994;269:26066-75.
- 43 Facchinetti F, Longo M, Piccinni F, Neri I, Volpe A. L-arginine infusion reduces blood pressure in pre-eclamptic women through nitric oxide release. *J Soc Gynecol Investig* 1999;4:202-7.
- 44 Staff AC, Berge L, Haugen G, Lorenzen B, Mikkelsen B, Henriksen T. Dietary supplementation with L-arginine or placebo in women with pre-eclampsia. *Acta Obstet Gynecol Scand* 2004;83:103-7.
- 45 Rytlewski K, Olszanecki R, Korbut R, Zdebski Z. Effects of prolonged oral supplementation with L-arginine on blood pressure and nitric oxide synthesis in pre-eclampsia. *Eur J Clin Invest* 2005;35:32-7.
- 46 Germain AM, Valdez G, Romanik MC, Reyes S. Letter to the editor: evidence supporting a beneficial role for long term L-arginine supplementation in high-risk pregnancies. *Hypertension* 2004;44:e1.
- 47 Neri I, Monari F, Sgarbi L, Berardi A, Masellis G, Facchinetti F. L-arginine supplementation in women with chronic hypertension: impact on blood pressure and maternal and neonatal complications. *J Matern Fetal Neonatal Med* 2010;23:1456-60.

Accepted: 28 March 2011