Diabetes mellitus as a compelling indication for use of renin angiotensin system blockers: systematic review and meta-analysis of randomized trials

Sripal Bangalore,¹ Robert Fakheri,¹ Bora Toklu,² Franz H Messerli³

¹New York University School of Medicine, New York, NY, USA

²Mount Sinai Beth Israel Medical Center, New York, NY, USA

³Mount Sinai Health Medical Center, Icahn School of Medicine, New York, NY, USA

Correspondence to: S Bangalore
sripalbangalore@gmail.com

Additional material is published online only. To view please visit the journal online.

Cite this as: *BMJ* 2016;352:i438
<http://dx.doi.org/10.1136/bmj.i438>

Accepted: 08 January 2016

ABSTRACT

OBJECTIVE

To evaluate the outcomes with use of renin angiotensin system (RAS) blockers compared with other antihypertensive agents in people with diabetes.

DESIGN

Meta-analysis.

DATA SOURCES AND STUDY SELECTION

PubMed, Embase, and the Cochrane central register of controlled trials databases for randomized trials of RAS blockers versus other antihypertensive agents in people with diabetes mellitus. Outcomes were death, cardiovascular death, myocardial infarction, angina, stroke, heart failure, revascularization, and end stage renal disease.

RESULTS

The search yielded 19 randomized controlled trials that enrolled 25 414 participants with diabetes for a total of 95 910 patient years of follow-up. When compared with other antihypertensive agents, RAS blockers were associated with a similar risk of death (relative risk 0.99, 95% confidence interval 0.93 to 1.05), cardiovascular death (1.02, 0.83 to 1.24), myocardial infarction (0.87, 0.64 to 1.18), angina pectoris (0.80, 0.58 to 1.11), stroke (1.04, 0.92 to 1.17), heart failure (0.90, 0.76 to 1.07), and revascularization (0.97, 0.77 to 1.22). There was also no difference in the hard renal outcome of end stage renal disease (0.99, 0.78 to 1.28) (power of 94% to show a 23% reduction in end stage renal disease).

CONCLUSIONS

In people with diabetes, RAS blockers are not superior to other antihypertensive drug classes such as thiazides, calcium channel blockers, and β blockers at reducing the risk of hard cardiovascular and renal

endpoints. These findings support the recommendations of the guidelines of the European Society of Cardiology/European Society of Hypertension and eighth Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure to also use other antihypertensive agents in people with diabetes but without kidney disease.

Introduction

People with diabetes are at increased risk of cardiovascular and renal events.¹ Early placebo controlled trials (such as the Heart Outcomes Prevention Evaluation and European Trial on Reduction of Cardiac Events With Perindopril in Stable Coronary Artery Disease) have shown significant benefits from use of renin angiotensin system (RAS) blockers on cardiovascular and renal events in people with diabetes, benefits touted to be independent of the drugs blood pressure lowering efficacy. As such, the 2015 American Diabetes Association guidelines recommend RAS blockers (angiotensin converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs)) as first line treatment for people with diabetes and hypertension.² Similarly, the 2013 American Society of Hypertension/International Society of Hypertension guidelines favor RAS blockers as a first line treatment in people with diabetes.³ The National Kidney Foundation-Kidney Disease Outcomes Quality Initiative clinical practice guidelines state in its executive summary that “Hypertensive people with diabetes and chronic kidney disease stages 1-4 should be treated with an ACE inhibitor or an ARB, usually in combination with a diuretic.”⁴ In contrast, the 2013 European Society of Cardiology/European Society of Hypertension guidelines⁵ and the 2014 evidence based guidelines from the panel members of the eighth Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure⁶ recommend any class of antihypertensive agents in people with diabetes, with a preference for RAS blockers only in the presence of proteinuria or microalbuminuria. This seemingly discordant set of recommendations begs the questions about the evidence base to support superior cardioprotective and renoprotective effects of RAS blockers in people with diabetes.

We explored whether RAS blockers are superior to other antihypertensive agents for the prevention of hard cardiovascular and renal events in people with diabetes.

Methods

Eligibility criteria

We searched PubMed, Embase, and the Cochrane central register of controlled trials until December 2015 (week 1)

WHAT IS ALREADY KNOWN ON THIS TOPIC

Various guidelines recommend renin angiotensin system (RAS) blockers as first line treatment for people with diabetes, predominantly based on placebo controlled trials done 20 years ago

However, other guidelines recommend RAS blockers on a par with other antihypertensives based on more recent trials comparing RAS blockers versus active comparators

WHAT THIS STUDY ADDS

Our study suggests that in people with diabetes, RAS blockers are similar to other antihypertensives at reducing the risk of hard cardiovascular and renal endpoints. These findings support the European Society of Cardiology/European Society of Hypertension and eighth Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure guideline recommendations. Those guidelines recommend using any antihypertensives in people with diabetes but without kidney disease.

for randomized controlled trials of RAS blockers (ACE inhibitor or ARB) (see supplementary table S1 for MeSH terms) in people with diabetes or impaired fasting glucose. There were no language restrictions for the search. In addition, we searched the bibliography of identified original trials, meta-analyses, and review articles to find other eligible trials ("snowball search"). Weekly reminders from PubMed kept the search up to date.

Eligible trials had to fulfill two criteria: randomized controlled trials comparing RAS blockers with other antihypertensive agents in participants with diabetes or impaired fasting glucose, and a sample size of at least 100 participants with diabetes with follow-up of at least one year (to minimize small study effect). We excluded studies conducted in cohorts with heart failure given the known efficacy of RAS blockers in this patient group. In addition, we excluded studies that had been redacted for any reason, compared ACE inhibitors with ARBs, RAS blockers with placebo, or randomized participants to an ACE inhibitor plus ARB.

Trial selection and bias assessment

Three authors (RF, BT, SB) independently assessed trial eligibility, trial bias risk, and data extraction, with disagreements resolved by consensus. The bias risk of trials was assessed using the components for randomized trials recommended by the Cochrane Collaboration⁷: allocation sequence generation, allocation concealment, and blinding of outcome assessors. For each component, we categorized trials as being at low, high, or unclear risk of bias. We considered trials with high or unclear risk of bias for any one of the above components as trials with high risk of bias.

Outcomes

Outcomes were death, cardiovascular death, myocardial infarction, angina, stroke, heart failure, revascularization, end stage renal disease, major adverse cardiovascular events, and drug withdrawal owing to adverse events.

Statistical analyses

Statistical analyses were performed using an intention to treat approach and in line with recommendations from the Cochrane Collaboration and the preferred reporting items for systematic reviews and meta-analyses statement.^{7,8} We carried out analyses to compare RAS blockers (ACE inhibitor or ARB) with other agents. Subgroup analyses compared RAS blockers with each class of comparative agent (calcium channel blockers, diuretics, and β blockers). The meta-analytic summary estimates (relative risk) were calculated using the fixed effect model and the random-effects model of DerSimonian and Laird.⁹ Continuity correction was used for trials with zero events. We used the I^2 statistic to assess heterogeneity—the proportion of total variation observed between the trials attributable to differences between trials rather than to sampling error (chance),¹⁰ with an I^2 value less than 25% considered low and more than 75% high. We assessed small study effect using the Begg's and the Egger's test and by visual evaluation of the funnel plots for asymmetry.

A metaregression analysis was performed to evaluate the relation of percent of participants with nephropathy at baseline on the outcomes. We used a residual maximum likelihood to estimate the additive (between study) component of variance τ^2 for the metaregression analysis. Bootstrap analyses were performed using a Monte Carlo permutation test for metaregression, using 10 000 random permutations.¹¹ Sensitivity analyses were done after excluding trials that included participants with impaired fasting glucose. Analyses were performed using standard statistical software (Stata 12.1, Stata, TX),¹² with $P < 0.05$ used to denote statistical significance.

Patient involvement

No patients were involved in setting the research question or the outcome measures, nor were they involved in developing plans for design or implementation of the study. No patients were asked to advise on interpretation or writing up of results. There are no plans to disseminate the results of the research to study participants or the relevant patient community.

Results

Study selection and baseline characteristics

Our search yielded 19 randomized controlled trials (see supplementary figure S1), enrolling 25 414 people with diabetes. The participants were followed for a mean of 3.8 years, totalling 95 910 patient years of follow-up. Fifteen trials compared RAS blockers with a calcium channel blocker, three with a thiazide diuretic, and two with a β blocker. In 14 trials the RAS blockers were an ACE inhibitor and in six an ARB. All trials enrolled people with type 2 diabetes. In addition, the majority of the trials ($n=17$) enrolled people with diabetes and hypertension.

Table 1 outlines the baseline characteristics of the included trials. In the studies that reported race/ethnicity, black people were a minority of the enrolled patients. The primary endpoint for the trials was heterogeneous (see supplementary table S2). Only three trials enrolled patients with microalbuminuria or proteinuria.

Outcomes

RAS blockers versus other antihypertensives

When compared with other antihypertensive agents, RAS blockers were associated with a similar risk of death (relative risk 0.99, 95% confidence interval 0.93 to 1.05, fig 1), cardiovascular death (1.02, 0.83 to 1.24, fig 2), myocardial infarction (0.87, 0.64 to 1.18, fig 3), angina pectoris (0.80, 0.58 to 1.11, fig 4), stroke (1.04, 0.92 to 1.17, fig 5), heart failure (0.90, 0.76 to 1.07, fig 6), and revascularization (0.97, 0.77 to 1.22, fig 7). The outcome of major adverse cardiovascular events (0.97, 0.89 to 1.06) did not differ between the two groups. In addition, there was no difference in end stage renal disease (0.99, 0.78 to 1.28, fig 8) or drug withdrawal owing to adverse effects (fig 9). The results were consistent when using a fixed effect model (figs 1-9). Heterogeneity was low to moderate, with no evidence of small study effect/publication bias (see supplementary figures S2-S10).

Table 1 | Baseline characteristics and risk of bias assessment of included trials

Trials	Year	Sample size	Follow-up (years)	Cohort	Age (years)	Black people (%)	Risk of bias*
RAS blockers versus calcium channel blockers:							
ABCD (hypertensive) ^{13, 14}	1998	470	5.6	Diabetes mellitus and hypertension	58	14	+++
ABCD (normotensive) ¹⁵	2002	354	5.3	Diabetes mellitus and normotensive	59	7	+++
ALLHAT ^{16, 17} (diabetes mellitus)	2002	7107	4.9	Diabetes mellitus and hypertension	67	39	+++
ALLHAT ^{16, 17, 18} (impaired fasting glucose)	2002	771	4.9	Impaired fasting glucose and hypertension	67	30	+++
BENEDICT ¹⁹	2004	604	3.6	Diabetes mellitus and hypertension	62	NR	±±±
CAMELOT ²⁰ (diabetes mellitus)	2004	233	2	Diabetes mellitus and coronary artery disease	58	NR	+++
CAMELOT ²⁰ (impaired fasting glucose)	2004	233	2	Impaired fasting glucose and coronary artery disease	58	NR	+++
CASE-J ^{21, 22, 23} (diabetes mellitus)	2008	1195	3.2	Diabetes mellitus and hypertension	67	NR	+++
FACET ²⁴	1998	380	2.9	Diabetes mellitus and hypertension	63	NR	±±±
Fogari et al ²⁵	2002	205	4	Diabetes mellitus with proteinuria and hypertension	63	NR	±±±
IDNT ^{26, 27, 28}	2001	1146	2.6	Diabetes mellitus with nephropathy and hypertension	60	13	±±±
JMIC-B ^{29, 30} (diabetes mellitus)	2004	372	3	Diabetes mellitus, hypertension, and coronary artery disease	64	NR	+++
J-MIND ³¹	2001	436	2	Diabetes mellitus and hypertension	60	NR	±±±
MITEC ³²	2009	209	2	Diabetes mellitus and hypertension	60	NR	±±±
MOSES ^{33, 34} (diabetes mellitus)	2005	498	2.5	Diabetes mellitus, hypertension, and cerebrovascular accident	70	NR	+++
NAGOYA HEART ³⁵	2012	1150	3.2	Diabetes mellitus and hypertension	63	NR	±±±
STOP-Hypertension-2 ³⁶ (diabetes mellitus)	1999	466	5	Diabetes mellitus and elderly hypertension	76	NR	±±±
RAS blockers versus diuretic:							
ALLHAT ^{16, 17} (diabetes mellitus)	2002	9504	4.9	Diabetes mellitus and hypertension	67	39	+++
ALLHAT ^{16, 17, 18} (impaired fasting glucose)	2002	1035	4.9	Impaired fasting glucose and hypertension	67	30	+++
ANBP2 ^{37, 38} (diabetes mellitus)	2003	441	4.1	Diabetes mellitus and elderly hypertension	72	NR	±±±
NESTOR ³⁹	2003	569	1	Diabetes mellitus with microalbuminuria and hypertension	60	5	±±±
RAS blockers versus β blockers:							
UKPDS 39 ⁴⁰	1998	758	8.4	Diabetes mellitus and hypertension	56	8	+++
LIFE ^{41, 42, 43, 44} (diabetes mellitus)	2002	1195	4.8	Diabetes mellitus and hypertension with left ventricular hypertrophy	67	12	+++

ABCD=Appropriate Blood Pressure Control in Diabetes; ALLHAT=Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial; ANBP2=Second Australian National Blood Pressure Study; BENEDICT=Bergamo Nephrologic Diabetes Complications Trial; CAMELOT=Comparison of Amlodipine vs Enalapril to Limit Occurrences of Thrombosis; CASE-J=Candesartan Antihypertensive Survival Evaluation in Japan; FACET=Fosinopril Versus Amlodipine Cardiovascular Events Randomized Trial; IDNT=Irbesartan Type II Diabetic Nephropathy Trial; JMIC-B=Japan Multicenter Investigation for Cardiovascular Diseases-B; J-MIND=Japan Multicenter Investigation of Antihypertensive Treatment for Nephropathy in Diabetics; LIFE=Losartan Intervention For Endpoint reduction; MITEC=Media Intima Thickness Evaluation with Candesartan cilexetil; MOSES=Morbidity and Mortality After Stroke, Eprosartan Compared With Nitrendipine for Secondary Prevention; NAGOYA HEART=Comparison between valsartan and amlodipine regarding morbidity and mortality in patients with hypertension and glucose intolerance; NESTOR=Natrilix SR versus Enalapril Study in Type 2 diabetic hypertensives with microalbuminuria; NR=not reported; RAS-inh=Renin-Angiotensin System inhibitor; STOP-Hypertension=Swedish Trial in Old Patients with Hypertension; UKPDS=UK Prospective Diabetes Study Group.

*Represents risk of bias based on: sequence generation of allocation; allocation concealment and blinding. + represents low bias risk and ± unclear bias risk.

RAS blockers versus calcium channel blockers

When compared with calcium channel blockers, RAS blockers were associated with a similar risk of death (1.01, 0.92 to 1.10, fig 1), cardiovascular death (1.17, 0.90 to 1.50, fig 2), myocardial infarction (0.84, 0.54 to 1.30, fig 3), angina pectoris (0.69, 0.33 to 1.42, fig 4), stroke (1.08, 0.90 to 1.28, fig 5), and revascularization (1.01, 0.74 to 1.39, figs 7 and 10). However, RAS blockers were associated with a significant reduction in the risk of heart failure compared with calcium channel blockers (0.78, 0.70 to 0.88, figs 6 and 10). There was no difference in drug withdrawal owing to adverse effects (fig 9) or end stage renal disease (figs 8 and 10). Heterogeneity was low to moderate (fig 10), with no evidence of small study effect/publication bias (see supplementary figures S2-10).

RAS blockers versus thiazide diuretics

Compared with thiazide diuretics, RAS blockers were associated with a similar risk of death (0.99, 0.90 to 1.08, fig 1), cardiovascular death (0.50, 0.05 to 5.46, fig 2), and other outcomes (figs 3-9 and 11). Only three trials compared RAS blockers with diuretics and hence

the confidence interval for most outcomes was wide. Heterogeneity was low to moderate (fig 11), with no evidence of small study effect/publication bias (see supplementary figures S2-10).

RAS blockers versus β blockers

Compared with β blockers, RAS blockers were associated with a similar risk of death (0.84, 0.47 to 1.51, fig 1), cardiovascular death (0.87, 0.47 to 1.60, fig 2), and other outcomes tested (figs 3-9 and 12). Only two trials compared RAS blockers with β blockers and hence the confidence interval for most outcomes was wide. Heterogeneity was high for the outcome of death but low to moderate for other outcomes (fig 12), with no evidence of small study effect/publication bias (see supplementary figures S2-10).

Influence of nephropathy

Metaregression analysis, aimed to assess the relation of percent of patients with nephropathy at baseline on the outcomes, showed no significant relation ($P>0.05$) for all outcomes.

Fig 1 | Outcomes of death with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

Sensitivity analysis

Results were largely similar in sensitivity analysis excluding trials of participants with impaired fasting glucose (see supplementary table S3).

Discussion

This analysis of patients with diabetes (largely without microalbuminuria or proteinuria) with 95 910 patient years of follow-up from randomized trials failed to show a superiority of renin angiotensin system (RAS) blockade over other antihypertensive agents for reduction of cardiovascular and renal outcomes. More importantly, compared with other agents RAS blockers showed no benefit in reducing the risk of death or myocardial infarction and end stage renal disease. The results were consistent in comparisons of RAS blockers with all controls and with individual antihypertensive agents.

RAS blockers for diabetes

People with diabetes mellitus are at increased risk of hypertension, and the concomitant presence of diabetes and hypertension is associated with an exponential increase in the risk of cardiovascular, cerebrovascular,

and renal events.⁴⁵ Previous trials have shown that blood pressure reduction in such patients leads to a significant reduction in cardiovascular events, emphasizing the need for aggressive management of hypertension in this cohort.⁴⁶ However, whether one antihypertensive drug class is superior to the other is controversial. Early studies of RAS blockade in patients with diabetes and microalbuminuria showed a significant “renoprotective” effect (mainly by slowing progression to clinical proteinuria) compared with placebo,⁴⁷ leading to the recommendation of RAS blockers for this indication. This recommendation was later extended to all people with diabetes. In addition, small head to head comparison trials of RAS blockers (fosinopril) versus calcium channel blockers (amlodipine) such as the Fosinopril Versus Amlodipine Cardiovascular Events Randomized Trial (380 participants) showed a significant reduction in cardiovascular events (secondary endpoint of combined outcome of myocardial infarction, stroke, or hospital admission for angina) with RAS blockers compared with calcium channel blockers.²⁴ However, hard endpoints of death or myocardial infarction did not differ in this trial.

Fig 2 | Outcome of cardiovascular death with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

Fig 3 | Outcome of myocardial infarction with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

Fig 4 | Outcome of angina pectoris with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

Fig 5 | Outcome of stroke with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

Fig 6 | Outcome of heart failure with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

Fig 7 | Outcome of revascularization with renin angiotensin system (RAS) blockers compared with antihypertensives in people with diabetes

Subsequent guidelines, including the seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure promoted diabetes as a compelling indication for RAS blockade. The American Diabetic Association guidelines states “In people with diabetes, inhibitors of the renin-angiotensin system (RAS) may have unique advantages for initial or early treatment of hypertension.”

However, more recent trials have failed to show superiority of RAS blockers compared with other antihypertensive agents for hard cardiovascular outcomes.³⁵ In addition, studies with larger sample size (for example, the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT) with 13 101 patients with diabetes) also failed to show the superiority of RAS blockers compared with other

Fig 8 | Outcome of end stage renal disease with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

Fig 9 | Outcome of drug withdrawal owing to adverse effects with renin angiotensin system (RAS) blockers compared with other antihypertensives in people with diabetes

antihypertensive agents for cardiovascular outcomes.¹⁸ Indeed, two decades ago one author stated about the progression of chronic kidney disease that “despite some discrepancies in experimental studies, recent controlled clinical trials show a similar slowing of progression with either ACEi [ACE inhibitor] or CCB

[calcium channel blocker].”⁴⁸ Consequently opinion now diverges among various guidelines about the role of RAS blockade in people with diabetes, with some guidelines still recommending RAS blockers as preferred drugs and some relegating them to be on a par with other antihypertensive drug classes.

Fig 10 | Outcomes with renin angiotensin system (RAS) blockers compared with calcium channel blockers in people with diabetes

Fig 11 | Outcomes with renin angiotensin system (RAS) blockers compared with diuretics in people with diabetes

Fig 12 | Outcomes with renin angiotensin system (RAS) blockers compared with β blockers in people with diabetes

Given the controversy and the discordance in various guideline recommendations about the role of RAS blockers in people with diabetes we evaluated the comparative effectiveness of RAS blockers compared with other antihypertensive agents in people with diabetes, excluding cohorts where RAS blockers are

shown to provide benefit (that is, heart failure). We also excluded placebo controlled trials since placebo is not the standard of care in such patients. The results of this study with over 95 000 patient years of follow-up from head to head randomized trials of RAS blockers versus other antihypertensive agents failed to

show a superiority of RAS blockers over other antihypertensive agents for the prevention of hard cardiovascular or renal outcomes. The notable exception in our analysis was that the RAS blockers were superior to calcium channel blockers for the prevention of heart failure. This outcome (heart failure) was mainly driven by the ALLHAT trial, the findings of which have been criticized.⁴⁹ Our findings of similar outcomes with RAS blockers compared with other antihypertensive agents therefore supports both the 2013 Society of Cardiology/European Society of Hypertension guidelines⁵ and the 2014 eighth Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure⁶ recommendation that any class of antihypertensive agents can be used in people with diabetes.

Diabetes with proteinuria or microalbuminuria

Both the 2013 Society of Cardiology/European Society of Hypertension guidelines⁵ and the 2014 eighth Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure⁶ recommend RAS blockers in the presence of proteinuria or microalbuminuria. The guideline recommendations favoring RAS blockers in those with diabetes and chronic kidney disease are driven mainly by placebo controlled trials of RAS blockers where there was a reduction in doubling of serum creatinine concentration (the Microalbuminuria, Cardiovascular, and Renal Outcomes substudy of the Heart Outcomes Prevention Evaluation study (MICRO-HOPE) and the Reduction of Endpoints in NIDDM with the Angiotensin II Antagonist Losartan study (RENAAL) and end stage renal disease (RENAAL) compared with placebo.^{50 51} Casas and colleagues showed a greater benefit of RAS blockers on renal outcomes in placebo controlled trials than active comparison trials and concluded that the benefits of RAS blockers on renal outcomes probably results from a blood pressure lowering effect.⁵² Palmer and colleagues, in a meta-analysis of trials enrolling patients with diabetes and kidney disease, found that no drug regimen was more effective than placebo for reducing all cause mortality.⁵³ However, end stage renal disease and a doubling of creatinine concentration was less likely and regression of albuminuria was more likely with ACE inhibitors or ARBs compared with placebo.⁵³ Limited trials have shown this superiority in active comparator trials. In the Irbesartan Type II Diabetic Nephropathy Trial (1715 patients with diabetic nephropathy) the RAS blocker irbesartan compared with the calcium channel blocker amlodipine was associated with a significant reduction in the risk of the primary composite end point (32.6% v 41.1%; $P=0.006$) of a doubling of serum creatinine concentration, the development of end stage renal disease, or death from any cause, driven by differences in doubling of serum creatinine concentration (16.9% v 25.4%; $P<0.001$), with numerically lower end stage renal disease (14.2% v 18.3%; $P=0.07$) but with no difference in death (15.0% v 14.6%; $P>0.05$).²⁸ Finally, Wu and

colleagues in a meta-analysis of RAS blockers in patients with diabetes showed no statistically significant difference among treatments for the hard endpoint of end stage renal disease even in the placebo comparisons.⁵⁴ However, an ACE inhibitor reduced the risk of doubling of serum creatinine compared with placebo.⁵⁴ Our study excluded placebo controlled trials and failed to show a benefit for the outcome of end stage renal disease with RAS blockers compared with other antihypertensive agents. Our analysis with 17 626 patients (for the outcome of end stage renal disease) had a power of 94% to show a 23% reduction in end stage renal disease with RAS blockers compared with controls and is thus sufficiently powered to show a difference if one existed.

Limitations of this study

We used trial level data only for the analyses and hence were unable to control for differences between trials. Although a separate analysis in the cohort of patients with nephropathy is desirable, trials did not report outcomes separately for this cohort. Moreover, the definition of nephropathy was variable. In addition, most of the studies in patients with diabetes and nephropathy were placebo controlled trials and excluded in the current analysis. There were only a few trials for the RAS blocker compared with diuretics and RAS blockers compared with β blocker, and the analyses are likely underpowered. For the renal outcomes, we only evaluated end stage renal disease as an outcome as this is considered a "hard" renal endpoint. While one can argue that a doubling of serum creatinine concentration is a stringent renal endpoint, only one trial reported this outcome. Most patients with diabetes and hypertension require on an average two antihypertensive agents and thus which agent should be first may not be critical. The results were partly driven by the ALLHAT trial, given the sample size. However, this trial did not monitor urine protein.

Conclusions

This analysis of head to head comparison trials of RAS blockers versus other antihypertensive agents in people with diabetes (and largely without microalbuminuria or proteinuria) failed to show a superiority of RAS blockers compared with other antihypertensive agents for the prevention of hard outcomes. The results support the recommendation of both the 2013 European Society of Cardiology/European Society of Hypertension guidelines⁵ and the 2014 eighth report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure⁶ that any class of antihypertensive agents can be used in people with diabetes especially in those without renal impairment.

Contributors: SB conceived, designed, and supervised the study, carried out the statistical analysis, and drafted the manuscript. RF and BT acquired the data. All authors analyses and interpreted the data and critically revised the manuscript for important intellectual content. SB is the guarantor. He had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Funding: None.

Competing interests: All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: the following financial relationships with organisations that might have an interest in the submitted work in the previous three years: SB receives honorariums from Abbott, Boehringer Ingelheim, Daiichi Sankyo, Merck, Gilead, and Pfizer. FHM is a consultant for Daiichi, Sankyo, Pfizer, Takeda, Abbott, AbbVie, Servier, Medtronic, and Ipca Laboratories.

Ethical approval: Not required.

Data sharing: No additional data available.

Transparency: The guarantor (SB) affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned have been explained.

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>.

- Wingard DL, Barrett-Connor EL, Scheidt-Nave C, McPhillips JB. Prevalence of cardiovascular and renal complications in older adults with normal or impaired glucose tolerance or NIDDM. A population-based study. *Diabetes Care* 1993;16:1022-5. doi:10.2337/diacare.16.7.1022.
- Standards of medical care in diabetes—2015: summary of revisions. *Diabetes Care* 2015;38(Suppl 1):S4doi:10.2337/dc15-S003.
- Weber MA, Schiffrin EL, White WB, et al. Clinical practice guidelines for the management of hypertension in the community a statement by the American Society of Hypertension and the International Society of Hypertension. *J Hypertens* 2014;32:3-15. doi:10.1097/HJH.0000000000000065.
- KDOQI. KDOQI clinical practice guidelines and clinical practice recommendations for diabetes and chronic kidney disease. *Am J Kidney Dis* 2007;49(Suppl 2):S12-154. doi:10.1053/j.ajkd.2006.12.005.
- Mancia G, Fagard R, Narkiewicz K, et al. Task Force Members. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2013;31:1281-357. doi:10.1097/01.hjh.0000431740.32696.cc.
- James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014;311:507-20. doi:10.1001/jama.2013.284427.
- Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.0.0 ed: The Cochrane Collaboration 2008. www.cochrane-handbook.org. Accessed Dec 12, 2013.
- Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. Quality of Reporting of Meta-analyses. *Lancet* 1999;354:1896-900. doi:10.1016/S0140-6736(99)04149-5.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88. doi:10.1016/0197-2456(86)90046-2.
- Galbraith RF. A note on graphical presentation of estimated odds ratios from several clinical trials. *Stat Med* 1988;7:889-94. doi:10.1002/sim.4780070807.
- Higgins JP, Thompson SG. Controlling the risk of spurious findings from meta-regression. *Stat Med* 2004;23:1663-82. doi:10.1002/sim.1752.
- Bradburn MJ, Deeks JJ, Altman DG. Sbe24: metan - an alternative meta-analysis command. *Stata Tech Bull Reprints* 1998;8:86-100.
- Estacio RO, Jeffers BW, Hiatt WR, Biggstaff SL, Gifford N, Schrier RW. The effect of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin-dependent diabetes and hypertension. *N Engl J Med* 1998;338:645-52. doi:10.1056/NEJM199803053381003.
- Schrier RW, Estacio RO. Additional follow-up from the ABCD trial in patients with type 2 diabetes and hypertension. *N Engl J Med* 2000;343:1969. doi:10.1056/NEJM200012283432614.
- Schrier RW, Estacio RO, Esler A, Mehler P. Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes. *Kidney Int* 2002;61:1086-97. doi:10.1046/j.1523-1755.2002.00213.x.
- Barzilay JI, Davis BR, Cutler JA, et al. ALLHAT Collaborative Research Group. Fasting glucose levels and incident diabetes mellitus in older nondiabetic adults randomized to receive 3 different classes of antihypertensive treatment: a report from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Arch Intern Med* 2006;166:2191-201. doi:10.1001/archinte.166.20.2191.
- ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2002;288:2981-97. doi:10.1001/jama.288.23.2981.
- Whelton PK, Barzilay J, Cushman WC, et al. ALLHAT Collaborative Research Group. Clinical outcomes in antihypertensive treatment of type 2 diabetes, impaired fasting glucose concentration, and normoglycemia: Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Arch Intern Med* 2005;165:1401-9. doi:10.1001/archinte.165.12.1401.
- Ruggenti P, Fassi A, Ilieva AP, et al. Bergamo Nephrologic Diabetes Complications Trial (BENEDICT) Investigators. Preventing microalbuminuria in type 2 diabetes. *N Engl J Med* 2004;351:1941-51. doi:10.1056/NEJMoa042167.
- Nissen SE, Tuzcu EM, Libby P, et al. CAMELOT Investigators. Effect of antihypertensive agents on cardiovascular events in patients with coronary disease and normal blood pressure: the CAMELOT study: a randomized controlled trial. *JAMA* 2004;292:2217-25. doi:10.1001/jama.292.18.2217.
- Fukui T, Rahman M, Hayashi K, et al. CASE-J Study Group. Candesartan Antihypertensive Survival Evaluation in Japan (CASE-J) trial of cardiovascular events in high-risk hypertensive patients: rationale, design, and methods. *Hypertens Res* 2003;26:979-90. doi:10.1291/hypres.26.979.
- Ogihara T, Nakao K, Fukui T, et al. Candesartan Antihypertensive Survival Evaluation in Japan Trial Group. Effects of candesartan compared with amlodipine in hypertensive patients with high cardiovascular risks: candesartan antihypertensive survival evaluation in Japan trial. *Hypertension* 2008;51:393-8. doi:10.1161/HYPERTENSIONAHA.107.098475.
- Nakao K, Hirata M, Oba K, et al. CASE-J Trial Group. Role of diabetes and obesity in outcomes of the candesartan antihypertensive survival evaluation in Japan (CASE-J) trial. *Hypertens Res* 2010;33:600-6. doi:10.1038/hr.2010.38.
- Tatti P, Pahor M, Byington RP, et al. Outcome results of the Fosinopril Versus Amlodipine Cardiovascular Events Randomized Trial (FACET) in patients with hypertension and NIDDM. *Diabetes Care* 1998;21:597-603. doi:10.2337/diacare.21.4.597.
- Fogari R, Preti P, Zoppi A, et al. Effects of amlodipine fosinopril combination on microalbuminuria in hypertensive type 2 diabetic patients. *Am J Hypertens* 2002;15:1042-9. doi:10.1016/S0895-7061(02)03017-0.
- Atkins RC, Briganti EM, Lewis JB, et al. Proteinuria reduction and progression to renal failure in patients with type 2 diabetes mellitus and overt nephropathy. *Am J Kidney Dis* 2005;45:281-7. doi:10.1053/j.ajkd.2004.10.019.
- Berl T, Hunsicker LG, Lewis JB, et al. Irbesartan Diabetic Nephropathy Trial. Collaborative Study Group. Cardiovascular outcomes in the Irbesartan Diabetic Nephropathy Trial of patients with type 2 diabetes and overt nephropathy. *Ann Intern Med* 2003;138:542-9. doi:10.7326/0003-4819-138-7-200304010-00010.
- Lewis EJ, Hunsicker LG, Clarke WR, et al. Collaborative Study Group. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851-60. doi:10.1056/NEJMoa011303.
- Yui Y, Sumiyoshi T, Kodama K, et al. Japan Multicenter Investigation for Cardiovascular Diseases-B Study Group. Comparison of nifedipine retard with angiotensin converting enzyme inhibitors in Japanese hypertensive patients with coronary artery disease: the Japan Multicenter Investigation for Cardiovascular Diseases-B (JMIC-B) randomized trial. *Hypertens Res* 2004;27:181-91. doi:10.1291/hypres.27.181.
- Yui Y, Sumiyoshi T, Kodama K, et al. Nifedipine retard was as effective as angiotensin converting enzyme inhibitors in preventing cardiac events in high-risk hypertensive patients with diabetes and coronary artery disease: the Japan Multicenter Investigation for Cardiovascular Diseases-B (JMIC-B) subgroup analysis. *Hypertens Res* 2004;27:449-56. doi:10.1291/hypres.27.449.
- Baba S. J-MIND Study Group. Nifedipine and enalapril equally reduce the progression of nephropathy in hypertensive type 2 diabetics. *Diabetes Res Clin Pract* 2001;54:191-201. doi:10.1016/S0168-8227(01)00288-1.
- Baguet JP, Asmar R, Valensi P, Nisse-Durgeat S, Mallion JM. Effects of candesartan cilexetil on carotid remodeling in hypertensive diabetic patients: the MITEC study. *Vasc Health Risk Manag* 2009;5:175-83. doi:10.2147/VHRM.S3409.
- Schrader J, Lüders S, Kulschewski A, et al. MOSES Study Group. Morbidity and Mortality After Stroke, Eprosartan Compared with Nitrendipine for Secondary Prevention: principal results of a prospective randomized controlled study (MOSES). *Stroke* 2005;36:1218-24. doi:10.1161/01.STR.0000166048.35740.a9.

- 34 Boulanger JM, Hill MD. Morbidity and mortality after stroke--eprosartan compared with nitrendipine for secondary prevention: principal results of a prospective randomized controlled study (MOSES). *Stroke* 2006;37:335-6. author reply 338. doi:10.1161/01.STR.0000199659.31103.d2.
- 35 Muramatsu T, Matsushita K, Yamashita K, et al. NAGOYA HEART Study Investigators. Comparison between valsartan and amlodipine regarding cardiovascular morbidity and mortality in hypertensive patients with glucose intolerance: NAGOYA HEART Study. *Hypertension* 2012;59:580-6. doi:10.1161/HYPERTENSIONAHA.111.184226.
- 36 Hansson L, Lindholm LH, Ekblom T, et al. Randomised trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 study. *Lancet* 1999;354:1751-6. doi:10.1016/S0140-6736(99)10327-1.
- 37 Chowdhury EK, Owen A, Ademi Z, et al. Second Australian National Blood Pressure Study Management Committee. Short- and long-term survival in treated elderly hypertensive patients with or without diabetes: findings from the Second Australian National Blood Pressure study. *Am J Hypertens* 2014;27:199-206. doi:10.1093/ajh/hpt212.
- 38 Wing LM, Reid CM, Ryan P, et al. Second Australian National Blood Pressure Study Group. A comparison of outcomes with angiotensin-converting--enzyme inhibitors and diuretics for hypertension in the elderly. *N Engl J Med* 2003;348:583-92. doi:10.1056/NEJMoa021716.
- 39 Marre M, Puig JG, Kokot F, et al. Equivalence of indapamide SR and enalapril on microalbuminuria reduction in hypertensive patients with type 2 diabetes: the NESTOR Study. *J Hypertens* 2004;22:1613-22. doi:10.1097/01.hjh.0000133733.32125.09.
- 40 UK Prospective Diabetes Study Group. Efficacy of atenolol and captopril in reducing risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 39. *BMJ* 1998;317:713-20. doi:10.1136/bmj.317.7160.713.
- 41 Dahlöf B, Devereux RB, Kjeldsen SE, et al. LIFE Study Group. Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002;359:995-1003. doi:10.1016/S0140-6736(02)08089-3.
- 42 Kizer JR, Dahlöf B, Kjeldsen SE, et al. Stroke reduction in hypertensive adults with cardiac hypertrophy randomized to losartan versus atenolol: the Losartan Intervention For Endpoint reduction in hypertension study. *Hypertension* 2005;45:46-52. doi:10.1161/01.HYP.0000151324.05355.1c.
- 43 Lindholm LH, Ibsen H, Borch-Johnsen K, et al. LIFE study group. Risk of new-onset diabetes in the Losartan Intervention For Endpoint reduction in hypertension study. *J Hypertens* 2002;20:1879-86. doi:10.1097/00004872-200209000-00035.
- 44 Lindholm LH, Ibsen H, Dahlöf B, et al. LIFE Study Group. Cardiovascular morbidity and mortality in patients with diabetes in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002;359:1004-10. doi:10.1016/S0140-6736(02)08090-X.
- 45 Epstein M, Sowers JR. Diabetes mellitus and hypertension. *Hypertension* 1992;19:403-18. doi:10.1161/01.HYP.19.5.403.
- 46 Parving HH, Andersen AR, Smidt UM, Svendsen PA. Early aggressive antihypertensive treatment reduces rate of decline in kidney function in diabetic nephropathy. *Lancet* 1983;321:1175-9. doi:10.1016/S0140-6736(83)92462-5.
- 47 Viberti G, Mogensen CE, Groop LC, Pauls JF. European Microalbuminuria Captopril Study Group. Effect of captopril on progression to clinical proteinuria in patients with insulin-dependent diabetes mellitus and microalbuminuria. *JAMA* 1994;271:275-9. doi:10.1001/jama.1994.03510280037029.
- 48 Ritz E. Are antihypertensive drugs similar in protecting the kidney? *Am J Hypertens* 1995;8:53S-8S. doi:10.1016/0895-7061(95)00189-1.
- 49 Messerli FH. ALLHAT, or the soft science of the secondary end point. *Ann Intern Med* 2003;139:777-80. doi:10.7326/0003-4819-139-9-200311040-00012.
- 50 Heart Outcomes Prevention Evaluation Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. *Lancet* 2000;355:253-9. doi:10.1016/S0140-6736(99)12323-7.
- 51 Brenner BM, Cooper ME, de Zeeuw D, et al. RENAAL Study Investigators. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861-9. doi:10.1056/NEJMoa011161.
- 52 Casas JP, Chua W, Loukogeorgakis S, et al. Effect of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: systematic review and meta-analysis. *Lancet* 2005;366:2026-33. doi:10.1016/S0140-6736(05)67814-2.
- 53 Palmer SC, Mavridis D, Navarese E, et al. Comparative efficacy and safety of blood pressure-lowering agents in adults with diabetes and kidney disease: a network meta-analysis. *Lancet* 2015;385:2047-56. doi:10.1016/S0140-6736(14)62459-4.
- 54 Wu HY, Huang JW, Lin HJ, et al. Comparative effectiveness of renin-angiotensin system blockers and other antihypertensive drugs in patients with diabetes: systematic review and bayesian network meta-analysis. *BMJ* 2013;347:f6008. doi:10.1136/bmj.f6008.

Web extra: supplementary appendix