



Department of Biology, University of Oxford, Oxford, UK

christopher.dye@zoo.ox.ac.uk

Cite this as: *BMJ* 2022;377:o867

<http://dx.doi.org/10.1136/bmj.o867>

Published: 27 April 2022

## The benefits of large scale covid-19 vaccination

### New evidence confirms that fewer people die in better vaccinated communities

Christopher Dye *professor of epidemiology*

The first covid-19 vaccines were administered under emergency use authorisation in December 2020, just one year into the pandemic, a “miracle” of pharmaceutical innovation that has saved an estimated million lives or more in the US alone.<sup>1,2</sup> The authorisation was given on the basis of safety and efficacy in randomised controlled trials, which found that immunisation with Pfizer-BioNTech and Moderna mRNA vaccines protected a remarkably high percentage (>90%) of recipients from developing symptomatic infection and, to a lesser extent, from asymptomatic infection too. In other words, when tested against the SARS-CoV-2 variants prevailing in 2020 and early 2021, these novel covid-19 vaccines could stop the great majority of infections from causing illness and help to prevent transmission of SARS-CoV-2. But could vaccination prevent infection and illness on a large scale, outside the controlled environment of clinical trials? A linked study by Suthar and colleagues (doi:10.1136/bmj-2021-069317) adds to the evidence that it can, across the US.<sup>3</sup>

Given the practical challenges of scaling up immunisation programmes—maintaining cold chains, carrying out mass inoculation in busy or makeshift clinics, and accurately reporting both numbers vaccinated and health outcomes—real world vaccine effectiveness is typically less than the efficacy achieved in clinical trials. Following reports that effectiveness has remained high for a variety of outcomes (infection, symptomatic illness, visits to emergency departments, hospital admissions, severe illness, and death) in diverse settings,<sup>4-10</sup> Suthar and colleagues have now investigated the impact of covid-19 vaccination, largely with Pfizer and Moderna vaccines, across 2558 counties in 48 US states, covering nearly 80% of the US population. Their evaluation is based on more than 30 million cases of covid-19 and more than 400 000 deaths linked to covid-19, which were reported during the second year of the pandemic, between December 2020 and December 2021.<sup>3</sup>

They measured effectiveness by comparing reported covid-19 incidence and mortality rates in counties with very low (0-9%), low (10-39%), medium (40-69%), and high ( $\geq$ 70%) percentages of adults ( $\geq$ 18 years) who had received at least one dose of vaccine. During the first half of 2021, when the alpha variant of SARS-CoV-2 was dominant, the covid-19 mortality rate was reduced by 60%, 75%, and 81% in counties with low, medium, and high vaccination coverage, compared with counties that had very low coverage. The corresponding figures for the reduction in case incidence were 57%, 70%, and 80%. The impact on mortality was similar during the second half of 2021 when the delta variant became dominant in the US, with smaller effects on incidence.<sup>3</sup>

Clearly, counties with higher vaccination coverage had fewer covid-19 cases and deaths per head of population, and the measured effectiveness in counties with high vaccine coverage was reassuringly large. More than this, vaccination had a disproportionately large effect in counties with low and medium coverage. For instance, an incremental increase in coverage of only 20% (from very low to low) and 50% (from very low to medium) led to reductions in mortality of 60% and 75%, respectively.

Suthar and colleagues argue that vaccination benefits whole communities, and indeed it does when coverage is high.<sup>3</sup> But they did not seek, and their data do not show, any extra effect of herd immunity, whereby vaccinated people prevented the transmission of infection to others in their communities.<sup>11</sup> A more likely explanation for the disproportionately beneficial effect in counties with low and medium coverage is that vaccination campaigns first targeted older people who are at greatest risk of severe illness and death from covid-19. Vaccine rollout in most countries began with older and otherwise vulnerable people and progressively included younger and less vulnerable people. Hong Kong is a notable exception, where high covid-19 death rates in March 2022 were associated with low vaccination rates particularly among older people.<sup>12</sup> Unlike Hong Kong, the US has followed the norm: in states that have achieved relatively low vaccination coverage overall, the percentage of older people vaccinated is invariably higher than the population average.<sup>13</sup> Suthar and colleagues did not investigate the effect of vaccination by age, but doing this should be possible with existing data available to the US Centers for Disease Control and Prevention.

The findings of this study also make clear that many more lives could have been saved, and will be saved, by encouraging people to keep up to date with vaccination in the face of waning immunity and new SARS-CoV-2 variants and by achieving even higher population coverage. How many lives is a matter for others to explore. Meanwhile, this new study is another confidence booster for covid-19 vaccines.

Competing interests: The BMJ has judged that there are no disqualifying financial ties to commercial companies. The authors declare the following other interests: CD has done consultancy work for Qiagen unrelated to the topic of this editorial. Further details of The BMJ policy on financial interests is here: <https://www.bmj.com/sites/default/files/attachments/resources/2016/03/16-current-bmj-education-coi-form.pdf>.

Provenance and peer review: Commissioned; not peer reviewed.

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide licence to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s) based on

the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) licence any third party to do any or all of the above. **[please confirm]**

- 1 Branswell H. Why Covid-19 vaccines are a freaking miracle. 2022. <https://www.stat-news.com/2022/02/14/why-covid-19-vaccines-are-a-freaking-miracle/>.
- 2 Schneider EC, Shah A, Sah P, Moghadas SM, Vilches T, Galvani A. The U.S. COVID-19 vaccination program at one year: How many deaths and hospitalizations were averted? 2021. <https://www.commonwealthfund.org/publications/issue-briefs/2021/dec/us-covid-19-vaccination-program-one-year-how-many-deaths-and-hospitalizations-were-averted>.
- 3 Suthar AB, Wang J, Seffren V, Wiegand RE, Griffing S, Zell E. Public health impact of covid-19 vaccines in the US: observational study. *BMJ* 2022;377:e069317.
- 4 Dagan N, Barda N, Kepten E, et al. BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *N Engl J Med* 2021;384:1412-23. doi: 10.1056/NEJMoa2101765 pmid: 33626250
- 5 Haas EJ, Angulo FJ, McLaughlin JM, et al. Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data. *Lancet* 2021;397:1819-29. doi: 10.1016/S0140-6736(21)00947-8 pmid: 33964222
- 6 Self WH, Tenforde MW, Rhoads JP, et al IVY Network. Comparative effectiveness of Moderna, Pfizer-BioNTech, and Janssen (Johnson & Johnson) vaccines in preventing covid-19 hospitalizations among adults without immunocompromising conditions - United States, March-August 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:1337-43. doi: 10.15585/mmwr.mm7038e1 pmid: 34555004
- 7 Hall VJ, Foulkes S, Saei A, et al SIREN Study Group. COVID-19 vaccine coverage in health-care workers in England and effectiveness of BNT162b2 mRNA vaccine against infection (SIREN): a prospective, multicentre, cohort study. *Lancet* 2021;397:1725-35. doi: 10.1016/S0140-6736(21)00790-X pmid: 33901423
- 8 Lin D-Y, Gu Y, Wheeler B, et al. Effectiveness of Covid-19 Vaccines over a 9-Month Period in North Carolina. *N Engl J Med* 2022;386:933-41. doi: 10.1056/NEJMoa2117128 pmid: 35020982
- 9 Christie A, Henley SJ, Mattocks L, et al. Decreases in COVID-19 cases, emergency department visits, hospital admissions, and deaths among older adults following the introduction of COVID-19 vaccine - United States, September 6, 2020-May 1, 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:858-64. doi: 10.15585/mmwr.mm7023e2 pmid: 34111059
- 10 McNamara LA, Wiegand RE, Burke RM, et al. Estimating the early impact of the US COVID-19 vaccination programme on COVID-19 cases, emergency department visits, hospital admissions, and deaths among adults aged 65 years and older: an ecological analysis of national surveillance data. *Lancet* 2022;399:152-60. doi: 10.1016/S0140-6736(21)02226-1 pmid: 34741818
- 11 Dean NE, Halloran ME. Protecting the herd with vaccination. *Science* 2022;375:1088-9. doi: 10.1126/science.abo2959 pmid: 35271331
- 12 The Government of Hong Kong Special Administrative Region. Hong Kong Vaccination Dashboard. 2022. <https://www.covidvaccine.gov.hk/en/dashboard>.
- 13 Mayo Foundation for Medical Education and Research. U.S. COVID-19 vaccine tracker. 2022. <https://www.mayoclinic.org/coronavirus-covid-19/vaccine-tracker>.

This article is made freely available for personal use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.