Papers and Originals

Natural History of Cerebral Angiomas

W. R. HENDERSON,* F.R.C.S.; R. DE R. L. GOMEZ,† M.D.

[WITH SPECIAL PLATE FACING PAGE 572]

Brit. med. J., 1967, 4, 571-574

This paper is based on 70 patients who were seen during 1946-61 and were followed-up in 1966. The main purpose is to assess the natural history of angiomas, with particular reference to haemorrhage; also to emphasize some aspects of management. Only six angiomas were removed.

The first comprehensive studies of angiomas were by Cushing and Bailey (1928) (Special Plate, Fig. 1) and by Dandy (1928), who described a few cases of epilepsy in which large angiomas were found during exploratory operation, usually for suspected tumour. The possibility of surgical removal, however, had to await the introduction of angiography by Egas Moniz in 1927; and later the widespread use of the percutaneous method in the investigation of subarachnoid haemorrhage revealed many smaller, previously silent, angiomas, and led to a pronounced increase in the numbers diagnosed during the past 20 years. Angiography enabled operations to be planned to ensure the essential early division of all feeding arteries, and it became possible to remove completely many small and medium-sized angiomas, and sometimes even the more challenging large angiomas (Norlen, 1949; Paterson and McKissock, 1956; Olivecrona and Ladenheim, 1957); fits, however, were not always abolished, and previous neurological disability often remained. Prevention of recurrent haemorrhage became the chief indication for operation, but the risk of haemorrhage had to be assessed against the possibility of operative damage.

More recently it became recognized that haemorrhage sometimes occurs from very small angiomas, designated "cryptic" hamartomas by Crawford and Russell (1956). They are often situated in the depth of a sulcus (Special Plate, Fig. 2) and are apt to produce a large intracerebral haematoma, sometimes without subarachnoid bleeding, that may be rapidly fatal (Margolis *et al.*, 1951; Krakenbühl and Siebenmann, 1965). It seems that minute angiomas are sometimes destroyed by the haemorrhage and consequently may not show on angiograms. They are an important cause of so-called spontaneous cerebral haematomas of unknown origin, especially in children and young adults, which, when diagnosed in time, are curable by urgent evacuation; haematomas caused most of the preventable deaths in this series.

Clinical Groups

Of the 70 angiomas 63 were found by angiography, 4 at necropsy, and 3 at operation. They varied in size from a few millimetres to 10 cm., and may be grouped into three sizes (Table I). The size of the angioma is important in relation to the incidence of epilepsy (Mackenzie, 1953), of cranial bruit, and of intracerebral haematoma, and to the possibility of surgical removal.

* Neurosurgeon, General Infirmary at Leeds.

† Neurosurgeon, Mexico City.

The first manifestations in 68 patients were: (1) haemorrhage in 53 patients (78%) from angiomas of all sizes but mainly smaller angiomas without cranial bruit; (2) epileptic fits in 11 patients (16%) with large angiomas and bruit; (3) progressive hemiparesis in two patients with basal angiomas; and (4) high intracranial pressure suggesting tumour in two patients.

TABLE I—Size of 70 Angiomas in Relation to First Symptoms

Size of Angioma	Haemorrhage	Epilepsy	Slow Hemiparesis	High Intracranial Pressure	No Symptoms
16 small (2 cm. or less) ..	16	—	—	—	—
35 medium (> 2-6 cm.) ..	30	2	—	1	2
19 large (6 to 10 cm.) ..	7	9	2	1	—
70	53	11	2	2	2

The ages at onset varied from 3 to 60 years; 21 patients were in the twenties and 16 were under 20. Silent angiomas were present in two patients with ruptured aneurysm.

Haemorrhage

Haemorrhage (subarachnoid, intracerebral, or intraventricular) occurred in 58 patients; in 53 it was the first manifestation, and five patients with large angiomas had late haemorrhage, many (10, 19, 23, 30, and 37) years after the onset of epilepsy or slow hemiparesis. After the first haemorrhage five patients died (three from intracerebral haematoma and two from late haemorrhage) and five angiomas were removed or clipped. There were 23 mild attacks of sudden headache without (or very brief) loss of consciousness or focal signs, and 21 moderate attacks with either loss of consciousness or focal signs, or both, that disappeared, so that 44 patients recovered completely; and there were nine severe attacks (four with known haematoma) followed by incomplete recovery.

A second haemorrhage occurred in 28 patients out of 48; five died from the haemorrhage (after intervals of five weeks, 15 months, two, seven, and nine years) and one angioma was removed. Further haemorrhages, making altogether three to six attacks, occurred in nine patients, one of whom died after the fourth haemorrhage; and another patient had nine almost certain haemorrhages (five proved by lumbar puncture), all followed by good recovery. Thus six patients (all with large or deep inoperable angiomas) died from recurrent haemorrhage. There were only seven short intervals, of less than six weeks, before 54 recurrent haemorrhages; 38 intervals were one year or longer, and 16 were 5 to 26 years. Only three of the nine patients who had severe first attacks had further haemorrhages. Seven patients had full recovery after all of three to nine

haemorrhages from medium-size inoperable angiomas (mostly supplied by choroidal arteries) and remained well 17 to 30 years after the first haemorrhage.

As regards permanent disability, 35 patients (60%) recovered completely from all attacks (Table II); and 12 had permanent residual signs, mainly hemiparesis or hemianopia, or occasionally cortical sensory loss or mild dysphasia, that were often slight and caused little or no disability. There was impairment of memory in one patient, and mild personality change in another. Eleven patients later developed occasional epileptic attacks. Two patients were not working.

TABLE II.—Severity of Haemorrhage, Based on the Most Severe Attack, in 58 Patients

Size of Angioma	Severity of Haemorrhage			Died	Total
	Mild	Moderate	Severe		
Small	5 (1)	7 (4)	3 (2)	2 (2)	17 (9)
Medium	10	9 (1)	7 (2)	4 (2)	30 (5)
Large	2	2	2	5	11
	17 (1)	18 (5)	12 (4)	11 (4)	58 (14)

The figures in parentheses are of known intracerebral haematomas.

It is well known that haemorrhages from angiomas are less fatal and cause less morbidity than haemorrhages from saccular aneurysms, that they occur at an earlier age, that recurrent attacks occur after longer intervals, often of many years, and that angiomas do not cause arterial spasm and cerebral infarction that sometimes produces severe permanent disability or fatality after rupture of an aneurysm. Haemorrhages from very small angiomas tend to occur in early life, as might be expected with a congenital lesion, but severe bleeding from large angiomas that contain so many abnormal vessels, surprisingly, often does not occur until later life, suggesting its dependence on degenerative changes; necropsies, however, often show evidence of previous clinically unrecognized slight leakages of blood (Russell and Rubinstein, 1963) that probably are more common than is known, especially from large angiomas. It seems from this small series that medium-sized surface angiomas often bleed into cerebrospinal fluid without causing permanent disability, and that their removal need not be attempted if the brain is likely to be damaged.

Intracerebral Haematoma

Haematomas were found at operation in 14 patients (Table II); four died. In five patients the cerebrospinal fluid was normal, which made early diagnosis more difficult, and in eight it contained blood; none had cranial bruit. Nine angiomas were small and five were of medium size (the largest was 5 cm.). There were no known haematomas with large angiomas that often extend from a wide base on the brain surface deeply to the ventricle and could bleed easily into cerebrospinal fluid, either directly or by tracking along the potential space between a dense angioma and brain. Blood may collect inside this space, forming a small haematoma that is discovered during removal of the angioma, or it may extend widely into the brain. A large haematoma occupies the white matter up to the cortex (Special Plate, Fig. 2), and the subependymal glia also seems to form a barrier; it is likely that fibre tracts at the site of bleeding are torn while nearby compressed tracts are capable of recovery, so that permanent focal signs are usually localized but may be severe—for example, monoparesis or quadrantic hemianopia.

Clinically, there is great variation in the rapidity of onset. Usually sudden severe headache occurs with vomiting and perhaps neck stiffness, and, when the motor area is involved, rapid hemiparesis. Death may occur within a few hours, usually from ventricular haemorrhage (Crawford and Russell, 1956), and no effective treatment is possible. More commonly, and when operation is urgently necessary within a day or two, the sudden headache is followed after a delay of some hours

by progressive deterioration, with drowsiness deepening to coma, and with spreading signs and perhaps a dilating pupil which, when it occurs, is an important indication of haematoma; two such patients died in two and four days after late aspiration of 25 and 30 ml. of old blood (Special Plate, Fig. 2). With smaller haematomas improvement starts with return to consciousness within two days but focal signs often persist.

Large haematomas, up to 10 cm. long, are mostly in the central (sulcus) region, and there is early hemiparesis (nine haematomas arose from small convexity angiomas in the middle cerebral area), whereas haematomas nearer the poles of the hemisphere are usually smaller and have milder symptoms without focal signs. Diagnosis may be confused by a history of mild antecedent trauma to the head (as in two patients) that occasionally seems to precede haemorrhage (Margolis *et al.*, 1951; Crawford and Russell, 1956), and is probably coincidence, but if relevant it suggests that trauma may start bleeding from an angioma that is about to rupture spontaneously; negative burr-hole exploration for suspected extradural haemorrhage should include cerebral aspiration. In two patients with only drowsiness, inconspicuous signs, and normal lumbar puncture the symptoms appeared to have been gradual in onset, over a week or two, and were at first attributed to a rapidly growing cerebral tumour. Angioma should be suspected, first, when blood is obtained during burr-hole aspiration of a mass that produced symptoms of recent apparently gradual onset, because bleeding into a tumour is very rare; and, secondly, as a likely cause of cerebral stroke in children (six haematomas were in children aged 17 or younger). Permanent isolated focal signs—for example, hemianopia—in young patients after an uninvestigated acute cerebral illness, sometimes called meningitis, may have resulted from angioma haemorrhage.

Radiologically, there may be a laterally displaced pineal shadow. Angiography shows slight lateral displacement of the midline, which may be the only radiological evidence of haematoma, especially in the frontal or occipital region (a large angioma alone may produce a similar shift); but a haematoma in the central region, the common site, produces typical displacement of the middle cerebral artery, either downwards with parallel stretching of the ascending branches (Special Plate, Fig. 3) or upwards (Special Plate, Fig. 4).

Seven haematomas, sometimes unsuspected, were evacuated during craniotomy, and in seven patients up to 50 ml. of old blood was aspirated through a burr hole. Aspiration was sometimes late, or was inadequate, and probably all four deaths should have been prevented (two angiomas on the convexity were small and removable, and two were deep and inoperable). The patients who recovered remained well without recurrence

W. R. HENDERSON AND R. DE R. L. GOMEZ: CEREBRAL ANGIOMAS

FIG. 1.—Diagrammatic representation of medium-size superficial angioma showing enlarged feeding arteries and draining veins. Reproduced from Cushing and Bailey's *Tumours Arising from the Blood Vessels of the Brain*, by permission of the authors and of the publishers, Charles C. Thomas, Springfield, Illinois, U.S.A.

FIG. 2.—Intracerebral haematoma from small angioma (arrow) in a sulcus in a child aged 7 years who died 42 hours after haemorrhage (lumbar puncture normal).

FIG. 3.—Angiogram in boy aged 15 years showing feeding artery (arrow) to small angioma, characteristic early filling of draining vein during arterial phase, and stretching by intracerebral haematoma of ascending branches of depressed middle cerebral artery.

FIG. 4.—Angiogram in girl aged 10 years showing upward displacement of middle cerebral artery by large intracerebral haematoma without visible angioma which, at operation, was found histologically in adherent blood clot at anterior wall of haematoma.

FIG. 5.—Angioma in basal ganglia and internal capsule produced gradual hemiparesis for 19 years before fatal haemorrhage at age 45. The huge venous channel drained to the vein of Galen.

FIG. 6.—Angiogram (venous phase) of large 10-cm. angioma that caused focal epilepsy, showing drainage deeply to the vein of Galen and superficially to the sagittal sinus. Repeat angiography seven years later showed no change in size of angioma, but drainage through straight sinus was replaced by distended veins to skull base and behind orbit (associated with recent orbital congestion and proptosis).

W. R. HENDERSON AND R. DE R. L. GOMEZ: NATURAL HISTORY OF CEREBRAL ANGIOMAS

FIG. 1

FIG. 2

FIG. 3

FIG. 4

FIG. 5

FIG. 6

of bleeding for 7 to 14 years. Burr-hole aspiration is sometimes satisfactory, but evacuation by craniotomy is usually better, especially when the angioma is removable; very small angiomas that are hidden in a sulcus may be traced by adherent clot inside the haematoma, as was proved histologically in one patient with a large haematoma but no angioma radiologically (Special Plate, Fig. 4). In very ill children with hemiplegia, dilated pupil, and possibly a history of recent mild head injury, it is probably safest to exclude a large central intracerebral (or extradural) haematoma by aspiration through two burr holes without angiography; an angioma can be removed after angiography later.

Epilepsy

Five of the 11 patients had no other complaint (except two with occasional migraine) during 10 to 40 years; four had late haemorrhage (two fatal) after fits for 10, 23, 30, and 37 years; and two patients died, one after epilepsy for 26 years (with, during the last nine years, gradual hemiparesis, dysphasia, and eventually mental deterioration), and the other (following rapid deterioration immediately after repeat angiography and subsequent radiotherapy) after mild epilepsy for about 10 years. The angiomas were superficial and large, 5 to 10 cm. in size (Special Plate, Fig. 6). Nine patients had cranial bruit. The classical combination of epilepsy and cranial bruit is strongly suggestive of angioma, but it may occur with a meningioma and is not diagnostic without subarachnoid haemorrhage. The fits, with or without loss of consciousness, were often Jacksonian; they were usually separated by intervals varying from months to several years, and were well controlled by anti-convulsant drugs. Removal of angioma was not attempted (occasionally the risk of producing increased disability may be justified when fits are frequent and disabling).

Eleven other patients had fits that started after, and presumably were caused by, haemorrhage, often within 12 months but occasionally after several years; usually, persisting signs of cerebral damage were present, and in five patients there had been a haematoma. In contrast to the epilepsy group, most of the angiomas were smaller, 4 cm. or less, and without a bruit. Posthaemorrhagic fits also were infrequent and were well controlled by drugs.

E.E.G. was of no value. In a few patients after epilepsy for many years it was normal; slow waves in two patients with gradual hemiparesis merely supported the obvious presence of a destructive cerebral lesion, but of course gave no indication of pathology.

Progressive Hemiparesis

Two patients with large angiomas in the basal ganglia had slowly progressive severe hemiparesis and sensory loss, without fits, for 20 and 19 years, when the latter patient died from haemorrhage; necropsy showed that the angioma was confined to and had destroyed the basal ganglia and internal capsule (Special Plate, Fig. 5). Slow hemiparesis was the initial symptom in five out of Paterson's (1955) nine cases of angioma in the basal ganglia.

N. L. BROWSE ET AL.: MANAGEMENT OF PULMONARY EMBOLI

FIG. 1.—Case 1. Ascending phlebogram of the clinically normal left leg showing typical recent thrombus. A loose thrombus of this size would probably be lethal if it broke free. The thrombus, easily removed, is shown below.

FIG. 2.—Case 2. Left, phlebogram showing a complete left iliac vein thrombosis with irregular fresh thrombus jutting into the vena cava. Right, phlebogram one year after plication of the inferior vena cava. The plication at the upper border of L3 is patent and the left iliac vein has fully recanalized.

FIG. 3.—Case 3. Percutaneous femoral puncture phlebogram of patient who had a pulmonary embolus six months after plication of the inferior vena cava without preoperative phlebography. The right inferior vena cava shows thrombosis and recanalization below the plication at the level of the body of L3, but there is a patent left inferior vena cava providing an unguarded route for emboli from the left leg.

High Intracranial Pressure

Two patients had high intracranial pressure with severe papilloedema and headaches of apparently gradual onset, without a history of fits or haemorrhage, that was suggestive of tumour. One very large space-occupying angioma was above the tentorium, and the other, of medium size, in the posterior fossa, obstructed the cerebrospinal fluid pathways.

Cranial Bruit

The presence or absence of a bruit, heard most easily over the eyeball, was recorded in 37 cases. It was nearly always present with large angiomas (the smallest was 4 cm.) but not with small ones (the largest without bruit was 5 cm.). Only one patient was aware of the noise.

Posterior Fossa Angiomas

Logue and Monckton (1954) described nine posterior fossa angiomas which usually produced cerebellar or lower cranial nerve signs; two in the cerebellum were removed and seven were in the brain stem. The clinical signs (without adequate history) may be wrongly attributed to tumour; similarly, intermittent symptoms from repeated small haemorrhages may be attributed to disseminated sclerosis. Our three patients with cerebellar angiomas remained well 7, 13, and 21 years after one or more mild haemorrhages; nevertheless their removal probably should have been attempted.

Angiography

Ideally, early management of subarachnoid haemorrhage rests on surgical observation from the day of haemorrhage, which is really the first investigation; the indication for and timing of angiography are based on the patient's (general and neurological) fitness primarily for possible operation and secondarily for angiography, which is potentially as risky as operation. Angiography should be regarded as a first-stage operation rather than as a diagnostic investigation.

Aneurysm is the commonest source of subarachnoid haemorrhage, and before angiography it is assumed that early operation may be required to prevent possible recurrent bleeding; on the other hand, a ruptured aneurysm often produces arterial spasm that may be dangerously increased by angiography during the first two or three weeks with resulting cerebral infarction (spasm sometimes spontaneously causes recurrent loss of consciousness and focal signs that are mistakenly attributed to recurrent haemorrhage). The third danger after haemorrhage is the formation of a large intracerebral haematoma (from angioma or aneurysm), which is very important because death from this cause can be prevented by prompt evacuation within a day or two; it is the most important complication of angioma haemorrhage. The suspicion of haematoma or of spasm, and therefore the timing of angiography, is based mainly on careful clinical observation for signs suggestive of progressive cerebral compression or transient increased spasm; the possibility of haematoma requires either immediate angiography or, when the patient is very ill, burr-hole exploration without angiography.

Angiography is not at any time without risk, and it is sometimes necessary to restrict it, or perhaps stop it after one injection, without completion of the series, which should never be routine. When an angioma is found on the first injected side, further injections are rarely required, either to demonstrate the complete blood supply of a large angioma, which is unnecessary except when the surgeon thinks it may be removable, or, unless there have been no focal manifestations, to exclude a coincidental ruptured aneurysm, which is rare. Such an aneurysm may be seen on the feeding artery to the angioma,

as in one of the two patients with silent middle cerebral angioma; the other patient had a ruptured posterior cerebral aneurysm that had produced partial oculomotor nerve paresis. Vertebral angiography was used conservatively because it seemed that the rare finding of a removable angioma or aneurysm in the posterior fossa (or occipital lobe) did not justify the added risk of routine complete four-vessel angiography.

The conventional three-exposure angiography seems to be adequate; rapid serial angiography is unnecessary even in the hope of showing with certainty a minute (not thrombosed) angioma, which, in any event, may be difficult to find at operation except in the wall of an obvious haematoma (Special Plate, Fig. 4). Repeat angiography after several years is also unnecessary and is inadvisable; in three patients the angioma showed no change in size after five to seven years. (In a 24-year-old patient with a 10-cm. angioma (Special Plate, Fig. 6) that had produced occasional mild focal fits for about 10 years, repeat angiography produced immediate severe hemiparesis and aphasia, followed by further rapid deterioration after radiotherapy, and death within a few months.) Post-operative angiography (a single lateral injection) was routine after excision or clipping of the angioma.

Operation

Removal of most deep (basal ganglia, choroidal, or brain-stem) angiomas is impossible, and there is no certainty that occlusion of a feeding artery that may give a branch to normal brain will not cause serious disability. Many superficial small and medium-size angiomas can safely be removed, but it is inadvisable to risk producing disability by attempting removal of large angiomas except perhaps when situated near a pole of the hemisphere.

There were 22 operations. Four angiomas up to 4 cm. in size were removed. The feeding arteries to six angiomas were clipped; two disappeared but four partially remained (without further bleeding during 9 to 13 years). In eight patients a haematoma was aspirated, usually through a burr hole. Three angiomas were found during exploratory operation, and one patient had flap decompression for high intracranial pressure but without improvement.

Radiotherapy was tried in five patients with large angiomas but produced no improvement, and four have died. McWhirter and Dott (1955) pointed out that the angioma blood vessels, though abnormal, are not neoplastic and that there is no reason to expect them to be more affected than the surrounding normal vessels. They stated that radiotherapy should not be employed, as have Paterson and McKissock (1956) and Olivecrona and Ladenheim (1957).

Follow-up

Twenty-one of the 69 patients followed up have died, 15 (22%) from the effects of the angioma and 6 from unrelated causes (four lived 7 to 11 years after subarachnoid haemorrhage and two with silent angiomas had ruptured aneurysm); one untraced patient was known to be alive eight years after haemorrhage. Most of the 15 patients died from haemorrhage: four from intracerebral haematoma, three from intraventricular haemorrhage, two from late haemorrhage (after 23 years' epilepsy and 19 years' hemiparesis), and one death followed vertebral angiography after full recovery from the third subarachnoid haemorrhage in eight years from a mid-brain angioma (previously shown by carotid angiography); all were proved at necropsy. Two sudden deaths without necropsy were assumed to have resulted from haemorrhage from large angiomas—one after epilepsy for 39 years and the other eight years after previous haemorrhage. The remaining three deaths (no necropsy) occurred in patients with very large supra-

tentorial angiomas after epilepsy for 10 and 26 years, and after a six-months' history of high intracranial pressure.

All 15 angiomas were thought to be either too large or too deep for removal except two small middle cerebral convexity angiomas with haematoma that should have been cured; and death after delayed evacuation of haematoma should not have occurred in two patients with deep angiomas. The five avoidable deaths were thus due to intracerebral haematoma or angiography. As regards intervals between first symptom and death, only 3 (4.4%) patients died, all from haematoma, within the first two months, and seven within 15 months; the remaining eight deaths occurred after 7 to 39 years. The earlier deaths were due mainly to haematomas from small angiomas, while the later deaths were in patients with large angiomas.

Conclusion

The natural history of angioma haemorrhage seems to be related to the size of the angioma. Haemorrhages from the common medium-size angiomas are usually mild and may recur at long intervals over many years. Small angiomas, on the other hand, are apt to bleed in children and rapidly produce a dangerous intracerebral haematoma in the first attack, whereas fatal haemorrhages from large (inoperable) angiomas often do not occur until later life, sometimes after fits or progressive cerebral disability for many years.

Forty-nine (84%) of the 58 patients who had haemorrhage were alive five years after the first bleeding; only two patients died from the effects of a safely removable angioma (with haematoma). These results probably justify a conservative surgical policy, but it is nevertheless believed that more angiomas should have been removed to prevent fatal haemorrhages that will no doubt occur in future years. Svien and McRae (1965) found that only 16% of 68 unoperated patients (out of a total of 95) seen at the Mayo clinic during 1920–61 had died from the angioma. We agree with their conclusion that, in retrospect, conservative treatment is appropriate for 80 to 85% of patients, but when patients are first seen the possibility of fatal recurrent haemorrhage is unpredictable, and some deaths will be prevented by excising the considerable number of angiomas that are removable without risk of producing serious neurological damage. It is, however, by the prompt evacuation of intracerebral haematoma that many avoidable deaths can be prevented.

It is emphasized that angiography should be used with restraint, especially in patients with large angiomas (and in the investigation of all subarachnoid haemorrhages), to avoid producing increased disability or fatality.

It is a pleasure to thank all the physicians who referred patients, and also the patients' general practitioners, for assisting with the follow-up.

REFERENCES

- Crawford, J. V., and Russell, D. S. (1956). *J. Neurol. Neurosurg. Psychiat.*, 19, 1.
- Cushing, H., and Bailey, P. (1928). *Tumours Arising from the Blood Vessels of the Brain*. Springfield, Illinois.
- Dandy, W. E. (1928). *Arch. Surg.*, 17, 190.
- Egas Moniz (1927). *Rev. Neurol.*, 2, 72. (See also Lima, 1950.)
- Krayenbühl, H., and Siebenmann, R. (1965). *J. Neurosurg.*, 22, 7.
- Lima, P. M. de A. (1950). *Cerebral Angiography*. London.
- Logue, V., and Monckton, G. (1954). *Brain*, 77, 252.
- Mackenzie, I. (1953). *Ibid.*, 76, 184.
- McWhirter, R., and Dott, N. M. (1955). *British Practice in Radiotherapy*, edited by E. R. Carling, B. W. Windeyer, and D. W. Smithers, p. 343. London.
- Margolis, G., Odum G. L., Woodhall B., and Bloor, B. M. (1951). *J. Neurosurg.*, 8, 564.
- Norlen, G. (1949). *Ibid.*, 6, 475.
- Olivecrona, H., and Ladenheim, J. (1957). *Congenital Arteriovenous Aneurysms of the Carotid and Vertebral Arterial Systems*. Berlin.
- Paterson, J. H. (1955). *Schweiz. Arch. Neurol. Psychiat.*, 75, 392.
- and McKissock, W. (1956). *Brain*, 79, 233.
- Russell, D. S., and Rubinstein, L. J. (1963). *Pathology of Tumours of the Nervous System*, 2nd ed. London.
- Svien, H. J., and McRae, J. A. (1965). *J. Neurosurg.*, 23, 23.