The ratio of the expected deaths to the observed deaths was not constant in successive time periods, but showed a fall from 1:7.3 in 1931-40 to 1:3.3 in 1961-65. As the magnitude of this ratio can be regarded as a measure of the increased risk of developing intrathoracic malignancy when asbestosis is present, it would appear that this risk had been more than halved between 1931-40 and 1961-5, though the risk of an asbestotic developing an intrathoracic neoplasm in 1961-5 would appear to be about three times that when no asbestosis is present. Conclusions based solely on information contained in death certificates must be suspect because of incompleteness and inaccuracy of the information contained in an unknown number of the certificates. There is no doubt that the number of deaths, recorded with asbestosis as a cause of death in the decade 1931-40, fell short of the number of actual deaths to which asbestosis contributed, but there is good reason to believe that the gap between the actual and the recorded deaths with asbestosis has narrowed considerably in recent years. It is probable that fewer cases of intrathoracic malignancy have escaped detection, before death and at necropsy, in recent years than in 1931-40. The ratio of expected to observed deaths with intrathoracic malignancy in 1931-40 might have been higher than the calculated ratio of 1:7.3.

The evidence provided by the study of data on death certificates and that produced by Knox and his colleagues² indicate that the increased risk of developing intrathoracic malignancy, either because of the presence of asbestosis or because of a prolonged exposure to asbestos dust, has diminished since 1933, when measures of dust control became effective in at least some working areas of asbestos factories.

I wish to thank Dr. T. Lloyd-Davies and Dr. W. D. Buchanan, of the Factory Department of the Ministry of Labour, for giving permission for me to examine, and for making readily accessible, the copies of death certificates of cases of asbestosis in the possession of the Department. I also wish to thank Dr. J. F. Knox for allowing me to see the data included in the paper in press by Knox, Doll, and Hill.³—I am, etc.,

J. BEATTIE.

Queens' College,
Cambridge

REFERENCES

- ¹ Buchanan, W. D., *Ann. N.Y. Acad. Sci.*, 1965, 132, 507.
- ² General Register Office, *The Registrar General's Statistical Review of England and Wales*, Part 1, Tables, Medical, 1931 and yearly to 1965. H.M.S.O., London.
- ³ Knox, J. F., Doll, R. S., and Hill, I. D., *Brit. J. industr. Med.*, in press.

SIR,—Your leading article on cancer and asbestos (24 August, p. 448) refers to the detection of asbestos bodies in the sputum of about 50% of males in certain industrial cities. To the best of our knowledge nobody has actually reported on this, and we think therefore that you have accidentally substituted the word "sputum" for "lung smear," since Anjilvel and Thurlbeck,¹ on whose work you base your statement, were reporting on lung smears from necropsy material.

In view of your interest in the subject, may we mention, while writing, that the same authors state in their article that no particular association was noted between asbestos bodies in the lungs and the presence of cancer

in the 33 patients in the series with malignant disease? They also state: "It is clear that asbestos bodies are not more common in patients with malignant disease in the autopsy populations we have studied, and thus casual exposure to asbestos is not an important cause of malignancy in the general population."—I am, etc.,

W. P. HOWARD,
Secretary,
Asbestosis Information Committee.

London W.1.

REFERENCE

- ¹ Anjilvel, L., and Thurlbeck, W. M. *Canad. med. Ass. J.*, 1966, 95, 1179.

Elderly in the Wrong Unit

SIR,—We were most interested here in the paper "The Elderly in the Wrong Unit," by Dr. A. G. Mezey and others (6 July, p. 16), and in the subsequent letter from Dr. Bernard Isaacs (10 August, p. 373).

We agree wholeheartedly with the need for the comprehensive psychogeriatric unit as stressed in the paper and previously by Kay, Roth, and Hall.¹ We have been impressed by the large area of overlap in management problems between the ostensibly "psychogeriatric" and the ostensibly "pure geriatric" patients, and are currently investigating this problem in the Leicester area. Readers may be interested that in this mental hospital the geriatric work has been gathered into a self-contained unit (as has the work of other psychiatric subspecialties). As a step towards full integration with the area geriatric service a consultant geriatrician, who is not a psychiatrist, has been appointed to the staff and will commence his appointment on 1 November 1968. He will spend more than half his time working in our geriatric unit and the remainder of his time in the area geriatric service. A new assessment ward has been opened for him, and it will be most interesting to see how the pattern of admission develops in the various units in his charge.—I am, etc.,

Carlton Hayes Hospital, NORMAN KAYE.
Narborough, Nr. Leicester.

REFERENCE

- ¹ Kay, D. W. K., Roth, M., and Hall, M. R. P., *Brit. med. J.*, 1966, 2, 967.

Research into Mental Retardation

SIR,—The establishment of an Institute for Research into Mental Retardation in this country has been a welcome advance in a field of study which is now receiving more attention than ever before.

In recent years the number of specialists dealing with mental subnormality (mental deficiency in Scotland) has increased. There is as yet no national professional association entirely devoted to mental retardation, although there are regional societies for the study of the subject and relatively small sections of some professional organizations concerned with the specialty. The time may be ripe for the formation of a British Association on Mental Retardation, comparable perhaps to the American Association on Mental Deficiency.

At present hospitals and specialists working in mental retardation in different areas of the country tend to work in isolation although

they have generally similar clinical and administrative problems to meet. Today, as the approach to mental retardation becomes increasingly multi-disciplinary, a large number of specialists, geneticists, biochemists, neurologists, paediatricians, orthopaedic surgeons, psychologists, and others have a contribution to make in the diagnosis and management of mental retardation. A British Association on Mental Retardation open to all the specialists with an interest in it would appear to be a natural, logical, and essential development in the evolution of provision for the mentally retarded.—I am, etc.,

Stansfield View Hospital, D. A. SPENCER.
Todmorden, Lancs.

Obstructive Airways Disease

SIR,—The findings reported by Drs. D. J. Lane, J. B. L. Howell, and Mr. B. Giblin (21 September, p. 707) confirm previous reports by Park¹ and by others of reduced sensitivity to carbon dioxide in patients with chronic airways obstruction. However, this phenomenon does not exclude Park's suggestion that the initiation of CO₂ retention is by mechanical impairment of ventilation. Although the PCO₂ does not correlate negatively with the magnitude of airway resistance,² there is evidence that superadded restriction of ventilation is important in the development of chronic hypercapnia. Thus raised PCO₂ levels have been found to be related to reduced lung compliance,³ spirometric evidence of ventilatory restriction,^{4,5} and an inefficient rapid and shallow breathing pattern.⁶—I am, etc.,

J. J. SEGALL.

Islington Chest Clinic,
London N.19.

REFERENCES

- ¹ Park, S. S., *Amer. Rev. resp. Dis.*, 1965, 92, 245.
- ² Astin, T. W., and Penman, R. W. B., *Amer. Rev. resp. Dis.*, 1967, 95, 567.
- ³ Kahana, L. M., Aronovitch, M., and Place, R., *Amer. Rev. resp. Dis.*, 1963, 87, 699.
- ⁴ Segall, J. J., and Butterworth, B. A., *Scand. J. resp. Dis.*, 1966, 47, 215.
- ⁵ Palmer, K. N. V., and Diamant, M. L., *Lancet*, 1968, 1, 1233.
- ⁶ Burrows, B., Saksena, F. B., and Diener, C. F., *Ann. intern. med.*, 1966, 65, 685.

Computer Analysis of Dental Surveys

SIR,—As a footnote to your recent series of articles on "Medicine and the Computer,"¹ I should also like to point out that during the past few years the computer has played an increasingly important part in dental research. This is especially true in the field of dental epidemiology, where with the aid of these machines detailed results of surveys can be obtained in a matter of days as compared with several months, if not years, if the results are processed by hand.

Dental clinical data may be prepared by collecting the information on a clinical chart in the usual way, coding this information on to an abstract sheet, and finally punching cards from this sheet. Nevertheless, this method is time-consuming and necessitates considerable storage space to house the various documents. To speed up this process, increase the accuracy, and reduce the number of documents involved, a special set of punch cards has been designed.

This set consists of five cards the front of which are used for recording the information as it is collected. The first card is for general information about the individual, while the remaining four cards are for recording the findings of the clinical examinations (including the state of each surface of each tooth; the results of a brief orthodontic assessment; the presence, degree, and distribution of opacities, hypoplasia, and attrition; the state of dental cleanliness and the presence or absence of stain, calculus, and gingivitis). On the reverse of the last card a space is provided in which the observer may record clinical notes. It is also possible to indicate whether any additional record has been made, such as a clinical photograph, intra-oral x-ray, or cephalometric radiograph. Certain parts of the record such as the name and date of birth can be entered directly into the appropriate boxes; others have to be coded first. When the coding is complex provision is made for entering the information using more conventional methods.

The recorder enters the details of general information on the first card before the clinical examination begins, and then as the examiner dictates his findings for entry on the following four cards. Subsequently the information which was not coded at the time of examination is entered in the appropriate box in its coded form. The cards are now ready for punching; they are designed for use with an automatic punch and are so arranged that the information to be punched can be seen while it is punched into the same card. After punching and verifying the cards are sorted back into their original order and they are then ready for processing by the computer.

The set has been used for recording information from individuals of a wide age range. Several examiners have found that it was easy to use, and it has been filled in accurately by a number of different recorders, including student nurses and clerical staff, all of whom required only two or three minutes' instruction to become familiar with the layout of the design. Coding and punching can be done rapidly, especially by trained data-processing operators, and it has been found possible to produce detailed results of a survey of a group of children within days of the examinations.

The cards have been designed to record all the original information and not abstracted information. For example, the date of birth and date of examination are recorded and not the age of the patient. Similarly the state of teeth is recorded by the details of the state of each surface and not by using one of the prevalence indices. It has been emphasized¹ that the name, together with the sex and date of birth if entered on a record, identifies it to a particular individual. If these facts are collected according to defined rules all the records relating to one individual can be linked together, even if they are collected by different observers and have different identification numbers. The name was included in this record for this reason, but it has since proved useful for producing lists of the subject examined in both numerical and alphabetical order, and thus reducing some of the clerical work associated with surveys.—I am, etc.,

R. J. ANDERSON.

Department of Dental
Health,
The Dental School,
University of Birmingham,
Birmingham 4.

REFERENCES

- ¹ *Brit. med. J.*, 1968, 2, 823, 3, 51, 116, 180, 247, 309, 367, and 426.
- ² *Computers in the Service of Medicine*, 1968, Vol. 1. Oxford University Press.

Recognition for Seniority

SIR,—It is a sad reflection on the value that the Minister of Health places on the opinion of our professional representatives that he has overruled their claim that seniority in general practice should be recognized for what it is. In our surgeries has arrived a document (E.C.N.690) in which the Minister admits that he has decided, contrary to the representations of the profession, to insist on a minimum number of post-graduate training sessions to qualify for continuation of a payment which at present recognizes seniority *per se*.

The profession has had time to consider the joint discussions which led to the statement that seniority would be tied to continuing education, and has rejected it; it is Association policy that compulsory post-graduate training should not be a prerequisite for receiving payment for seniority. Younger principals in general practice who, like myself, have senior partners who have worked incredibly hard to build up a practice over a number of years, but who are still giving a first-class service, keeping in touch with modern medicine by discussion with their colleagues, and regularly reading medical journals, are the first to recognize and respect the true worth of seniority. The practitioner who has reached the stage of qualifying for a seniority award has years of experience to offer his patients, which no amount of post-graduate education will replace or enhance, and this the younger doctor cannot offer, however many courses he may attend.

By all means encourage all principals to continue with postgraduate education, and finance such activity as much as possible; but may our representatives do all in their power to fight this iniquitous Ministerial decision to deny senior practitioners the award which should be their reward for years of service to their patients.—I am, etc.,

Ipswich, Suffolk.

FRANK WELLS.

Records System for General Practice

SIR,—The question of general practice records is too important to be clouded by supposed misunderstandings and their refutations. The facts are clear: "... we think a new folder might be introduced gradually" (Dr. I. S. L. Loudon and others, 24 August, p. 501). Thus the need for a transitional period is clearly supported.

The questions now for this transitional stage are whether we have two markedly different-sized records to file: the new folder and the E.C. 5 and 6, or do we have one filing system which accommodates both the present and the future, until a radical change can eventually be effected with the help of mechanical handling.

The Record Wallet design opts for the latter by folding an approximate hospital folder size in three parts to fit the present filing requirements in shelves, drawers, or boxes. The factual experiment reported (18 May, p. 420) showed that this latter proposition was possible and practicable. The 7½ in. (18 cm.) tall dimension of the wallet was deliberately chosen as a compromise to retain as serviceable the English and Scottish present-day filing equipment.

Dr. Loudon and colleagues are perfectly entitled to decide that they prefer to accommodate two sizes of records during the transition period.—I am, etc.,

E. V. KUENSBERG.

Edinburgh.

** This correspondence is now closed.—
ED., B.M.J.

Responsibilities of Consultants

SIR,—The consultant staff of Bedford General Hospital (21 September, p. 746) might be interested to know that the Regional Hospitals' Consultants and Specialists Association is an independent association, created to further the interests of consultants and specialists; 20 years ago it was the only association to fight for the non-teaching members of the profession. While we have no direct negotiating power with the Ministry of Health we, as an association, are able to exert more influence than a consultant on his own, and therefore can act as the ombudsman for regional consultants and specialists who are having difficulties.

We would welcome any consultants and specialists who are "angry and dismayed" to join us; the more members we have, the stronger our voice.—I am, etc.,

DONALD YOUNG,

President, Regional Hospitals'
Consultants and Specialists Association.

Royal College of Surgeons,
London W.C.2.

Points from Letters

G.P. Obstetric Units

Dr. J. SHACKLETON BAILEY (Eye, Suffolk) writes: Well staffed and run general-practitioner maternity units offer mothers a service which in many respects is superior to that of the large central hospital ward, especially in remote areas. . . . No doubt with more staff and modernization the large obstetric unit will commend itself more both to patients and their doctors, but it must always have disadvantages from the point of view of both. . . . Perhaps in these days it is too late to refer to the rapport that general practitioner and patient establish through confinements in a family, as the loss of this sort of element, whatever the patients may still feel, does not seem to be so much a matter of regret among general practitioners as it would have been years ago. In relation to the individual episode I think it must still be recognized that personal relationships in or near the accustomed background are as relevant to obstetrics as geriatrics. To many general practitioners who take a positive view of their vocation it must seem that we should therefore plan many good maternity units, seeking to improve their use and safety rather than watch the obstetric pundits run them down and attempt to "educate" the public into the fashion of their brave new world of mass reproduction. The superiority claimed for the large central obstetric unit mainly rests on accessibility of blood and consultants. I don't think a consultant has ever been into our local maternity unit, and, though this may seldom be really indicated, one would feel that a more "domiciliary" attitude to such units by consultant obstetricians would be of advantage. To achieve this we should undoubtedly require an increase in the number of consultants and a greater recognition of the permanence of their role in relation to the general practitioner. The availability of blood for emergency transfusion could be even more easily achieved.