the onset of neurological signs her gait had worsened. The left lower limb now showed a spastic paresis with an extensor plantar response. Impairment of all forms of cutaneous sensibility below the D 7 level on the right side completed the picture of a partial Brown-Séquard syndrome. The sensory loss was most dense over the L 3 dermatome, and there was a band of hyperaesthesia at D 6 of the left side. There were no motor signs in the right leg. The C.S.F. protein had risen to 500 mg./100 ml. and the cell count was 121/cu. mm., mainly lymphocytes.

Spinal cord compression at the D 6 level was diagnosed and myelography undertaken by the lumbar route. The spinal block was partial and myelography was repeated by the cisternal route. Again the block was partial. There was flocculation of the dye into pockets throughout the mid-dorsal, lower dorsal, and lumbar regions. Though the patient was not febrile and the agglutination tests remained normal, myelitis and arachnoiditis due to brucellosis were suspected. Treatment with tetracycline, streptomycin, and prednisolone (60 mg./day) was instituted. The response was dramatic and she was soon able to walk. The C.S.F. was normal six weeks later, and she was able to walk out of the hospital in another fortnight with no residual neurological signs.

A month later she was seen as an outpatient and complained of some numbness over the lateral border of the left foot. Hypoalgesia and hypoaesthesia could be detected in the L 5 root area. Agglutination test for brucellosis remained negative and she responded rapidly to prednisolone and tetracycline. Two years later she remained well and had no neurological signs.

We did not have full laboratory facilities to do complement fixation tests.

COMMENT

There was no doubt about the diagnosis of brucellosis in this case. The organism was grown in culture, and the titre of agglutination tests was significant (Spink *et al.*, 1952). We could not demonstrate the organism in the C.S.F. or get a positive agglutination test from the fluid.

Neurological complications may appear at the onset of the illness, after a few febrile episodes, during convalescence, or long after the fever, and may at times overshadow the other manifestations of brucellosis (Nelson-Jones, 1951). Any of the common strains—*melitensis*, *abortus*, or *suis*—can involve the nervous system.

Leptomeningitis is an early feature and may lead to adhesive arachnoiditis in the cranium or the spinal canal. The dorsal region is most frequently affected. A pseudopsychiatric form of the disease is well known (Spink, 1959). Nelson-Jones (1951) stressed an encephalitic element with mental symptoms. In the absence of cranial nerve signs it is doubtful whether our patient had cranial meningitis. A mild encephalitis could explain her earlier nausea, vomiting, and giddiness.

Meningoencephalitis secondary to brucellosis has been described by Fincham *et al.* (1963) and by Nichols (1951).

Contralateral Pneumothorax with Congenital Diaphragmatic Hernia

Brit. med. J., 1968, 4, 433-434

The importance of congenital diaphragmatic hernia as the cause of acute cardiorespiratory difficulties in the perinatal period has been better appreciated since Butler and Claireaux (1962) found that it was present in 8% of perinatal deaths. As a result congenital diaphragmatic hernia is now usually diagnosed before death, and in the past 15 years at the Hospital for Sick Children there has been a sixfold increase in the number of babies with this anomaly who have been admitted within 24 hours of birth (Young, 1968).

E

The former have also reported meningomyelitis. Their case presented with ataxia and weakness of the legs, and had C.S.F. changes—protein 238 mg./100 ml. and 73 lymphocytes/cu. mm. Other forms described by Fincham *et al.* include subarachnoid haemorrhage, neuritis, and myelopathy. The meningo-encephalitic type can mimic an intracranial mass. DeJong (1936) gave a good account of neurological features of brucellosis. Steindler (1940) and Lowe and Lipscombe (1947) dealt mainly with neurological features secondary to skeletal disease, not seen in our patient.

Adhesive arachnoiditis is a frequent feature seen on laminectomy for suspected spinal cord compression in many tropical countries. It seems likely that brucellosis may account for a few of these cases. In view of the extremely gratifying response usually seen after specific therapy, it is desirable to look for evidence of brucellosis when faced with a case of arachnoiditis or meningomyelitis of obscure origin. The C.S.F. may show a growth of brucella in culture or be positive for agglutinins. These are, however, notoriously difficult to demonstrate (Fincham *et al.*, 1963).

We are grateful to the Director-General, Armed Forces Medical Services (India), for permitting us to report this case.

M. G. SAHADEVAN, M.B., M.R.C.P.,
Major.

MAHINDER SINGH, M.B., M.S.,
Lieutenant-Colonel.

P. P. JOSEPH, M.B., M.R.C.P.ED., M.R.C.P.GLASG.,
Major.

R. S. HOON, M.B., F.R.C.P.ED.,
Brigadier.

Armed Forces Medical College, Poona, India.

REFERENCES

- DeJong, R. N. (1936). *J. Nerv. ment. Dis.*, **83**, 430.
Fincham, R. W., Sahs, A. L., and Joynt, R. J. (1963). *J. Amer. med. Ass.*, **184**, 269.
Hughes (1897). *Mediterranean, Malta, or Undulant Fever*. Quoted by Nelson-Jones (1951).
Lowe, G. H., jun., and Lipscombe, P. R. (1947). *Surgery*, **22**, 525.
Mathur, T. N. (1954). *Indian med. Gaz.*, **89**, 292.
Mathur, T. N. (1955). *Indian J. med. Sci.*, **9**, 692.
Nelson-Jones, A. (1951). *Lancet*, **1**, 495.
Nichols, E. (1951). *Ann. intern. Med.*, **35**, 673.
Pathania, N. S., Sachdeva, J. R., and Chopra, J. S. (1966). *J. Indian med. Ass.*, **47**, 290.
Spink, W. W. (1959). In *Textbook of Medicine*, 10th ed., edited by R. L. Cecil and R. F. Loeb, p. 226. Philadelphia.
Spink, W. W., McCullough, N. B., Hutchings, L. M., and Mingle, C. K. (1952). *J. Amer. med. Ass.*, **149**, 805.
Steindler, A. (1940). Quoted by Pathania *et al.* (1966).
Tomaselli (1879). Cited by Nelson-Jones (1951).

These very young infants now present some problems not seen before, when surgeons saw only those babies who had survived for several days before admission. This report concerns one postoperative complication, the development of pneumothorax on the opposite side to the hernia. Contralateral pneumothorax is particularly important in the infant with early cardiorespiratory distress, in whom the diagnosis of the diaphragmatic hernia is made within hours of birth. The case histories of four infants admitted under Professor A. W. Wilkinson are given and some aspects of congenital diaphragmatic hernia discussed.

CASE REPORTS

Case 1.—A male infant had difficulty in establishing respiration at birth, which necessitated intubation, aspiration, and assisted

ventilation. He did not breathe adequately, and x-ray examination of the chest showed left diaphragmatic hernia. He was intubated and assisted ventilation was given during transfer. At operation five and a half hours after birth a left Bochdalek hernia was repaired with difficulty and a left intercostal drain inserted. An x-ray film taken after operation showed a right-sided pneumothorax, so underwater-seal drain was inserted into the right pleural cavity. This brought considerable improvement, but assisted ventilation had to be continued and he died 12 hours later. Necropsy findings confirmed that the left lung was hypoplastic (see Table), but no other defect was found.

Case 2.—This male infant, born at home, was pink at first but after three breaths became cyanosed, and he was transferred to hospital. An x-ray film of chest showed left diaphragmatic hernia. At operation six hours after birth a left Bochdalek hernia was repaired and left pleural drainage and tracheostomy were carried out. Postoperative x-ray film showed right pneumothorax, which was drained; assisted ventilation was continued until death at 36 hours. Necropsy showed very large patent ductus arteriosus and hypoplastic left lung (see Table).

Weight of Each Lung of the Four Infants With Left Diaphragmatic Hernia

Case No.	Infant's Weight	Right Lung	Left Lung	Age at Death
1	3,550 g.	16 g.	5 g.	1 day
2	2,870 g.	24 g.	8 g.	2 days
3	3,300 g.	56 g.	20 g.	5 days
4	2,700 g.	18 g.	7 g.	1 day

Case 3.—This infant, a male, required intubation at birth and spontaneous ventilation was then established. Dextrocardia was noticed, and an x-ray film of the chest showed left diaphragmatic hernia. On admission he was in acute distress and immediate intubation was necessary. At operation six hours after birth a left Bochdalek hernia was repaired and the left pleural cavity drained. He was ventilated postoperatively for 12 hours, and after extubation was in good condition for a further 24 hours, when x-ray examination showed good expansion of the left lung, but cardiac contour was unusual. He deteriorated and required tracheostomy, and 48 hours postoperatively further deterioration occurred owing to right pneumothorax, which was drained. He continued on assisted ventilation for a further two days until death. Necropsy showed large patent ductus arteriosus, preductal coarctation (diameter 2 mm.), right ventricular hypertrophy, and bilateral bronchopneumonia with the left lung less than half the weight of the right.

Case 4.—This infant required intubation at birth. Satisfactory spontaneous ventilation was not established, and he was transferred with intermittent positive-pressure ventilation. Temperature on admission was 94° F. (34.4° C.). At operation six hours after birth a left Bochdalek hernia was repaired, left pleural cavity was aspirated as the diaphragm was closed, but an intercostal drain was not inserted. Postoperative x-ray picture showed bilateral pneumothorax; bilateral intercostal drains were inserted. Postoperatively he required assisted ventilation until death 24 hours later. The only other finding at necropsy was pneumomediastinum and hypoplasia of the left lung.

DISCUSSION

Early diagnosis and prompt operation on an infant with congenital diaphragmatic hernia which has given rise to cardio-respiratory difficulties in the first few hours of life is the only form of treatment which offers a reasonable chance of success. Such infants, even when early intubation and assisted ventilation are necessary, may withstand journeys lasting an hour or two, during which manual ventilation is necessary, and may

subsequently survive surgical repair of the hernia (Eckstein and Glover, 1964).

Diagnosis may be difficult because the clinical signs are different from those in older infants. The most valuable sign is dextrocardia, which is found in 90% of infants with left-sided hernia, and it was present in the four cases reported here. Moreover, 95% of infants admitted within 24 hours of birth have a left-sided hernia. A plain radiograph of the chest is the most useful aid to diagnosis, but if this entails delay it is probably better to transfer the baby immediately to a surgical unit.

Pneumothorax on the side of the hernia is inevitably produced at the time of operation and is treated appropriately.

Contralateral pneumothorax, which is a particular hazard in infants admitted for correction of diaphragmatic hernia within 24 hours of birth, has seldom been recorded (McCaughy, 1963; Wolf *et al.*, 1968). Each of the four infants described was admitted within six hours of birth; the severe disturbance caused by the right-sided pneumothorax contributed to the death of two. It would therefore seem justifiable, in the infant admitted within 24 hours of birth, to institute bilateral intercostal drainage at the time of operation.

The normal neonatal lung is relatively resistant to damage during assisted ventilation, but the infant with diaphragmatic hernia has proved to be highly susceptible to such damage. The hypoplasia of both lungs, the more affected being the lung on the side of the hernia, has now been recorded in both the human patient (Carter *et al.*, 1962; Areechon and Reid, 1963; Gross, 1964) and the experimental animal (deLorimier *et al.*, 1967). The opening pressure in the small hypoplastic air spaces is considerably higher than in the normal, so that the increased liability to pneumothorax in infants with congenital diaphragmatic hernia is probably due to surfactant deficiency resulting in the opening pressure in some parts of the lung being in excess of the rupturing pressure in other parts of that lung. The knowledge of the hypoplasia of the lungs and the consequent dangers make it of the utmost importance that the possibility of pneumothorax should be kept in the clinician's mind when he is treating a newborn infant with congenital diaphragmatic hernia.

I wish to record my thanks to Professor A. W. Wilkinson for permission to report these cases and for his help in the preparation of this article.

D. G. YOUNG, M.B., CH.B., F.R.C.S.ED.,
Senior Lecturer in Paediatric Surgery, Institute of
Child Health, University of London, London
W.C.1, and Hospital for Sick Children, Great
Ormond Street, London W.C.1.

REFERENCES

- Areechon, W., and Reid, L. (1963). *Brit. med. J.*, **1**, 230.
- Butler, N., and Claireaux, A. E. (1962). *Lancet*, **1**, 659.
- Carter, R. E. B., Waterston, D. J., and Aberdeen, E. (1962). *Lancet*, **1**, 656.
- deLorimier, A. A., Tierney, D. F., and Parker, H. R. (1967). *Surgery*, **62**, 12.
- Eckstein, H. B., and Glover, W. J. (1964). *Lancet*, **1**, 427.
- Gross, R. E. (1964). *J. thorac. cardiovasc. Surg.*, **48**, 152.
- McCaughy, T. J. (1963). *Canad. Anaesth. Soc. J.*, **10**, 616.
- Wolf, S. I., Hassan, K. J., and Randolph, J. (1968). *Clin. Proc. Child. Hosp. (Wash.)*, **24**, 12.
- Young, D. G. (1968). *Recent Advances in Paediatric Surgery*, 2nd ed., edited by A. W. Wilkinson. In press.