does not provide a forum for the discussion which is essential for managing complex conditions." There is increasing professional and public interest in this subject, and it is to be hoped that my questions will stimulate open discussion by other practitioners of what has hitherto been an esoteric field.—I am, etc.,

BASIL A. STOLL

London N.W.11.

Labelling of Poisonous Commercial Preparations

SIR,—I was pleased to read (*Supplement*, 4 October, p. 1) that the B.M.A. Public Health Committee had taken up the question of labelling of commercial preparations with the Home Office, and that despite its adverse reply they were going to pursue the matter further with the Home Office.

The three factors which were put forward by the Home Office as influencing its decision not to carry out the B.M.A.'s recommendations are quite indefensible. Some answers to these factors are as follows. The extent to which labelling is already required under statutory schemes is extremely limited. There are many poisonous substances present in household preparations which are exempted from the Pharmacy and Poisons Acts, particularly by Schedule III but also by Schedule II of these Acts. In addition the majority of the common poisoning agents such as ethylene glycol, carbon tetrachloride, turpentine, paraffin, strong sodium hypochlorite solutions, and methyl ethyl ketone peroxide, to mention only a few, are not even mentioned in these Acts. For example, ethylene glycol (L.D. adults 3–4 fl. oz. (100 ml.) and therefore much less for a child) is a common constituent of motor-car anti-freeze preparations and brake fluids, and these must be present in millions of homes throughout this country with nothing on the labels to warn people that these preparations are dangerous. The same goes for many other poisonous substances present in common commercial preparations which are routinely used in millions of homes in this country. Voluntary schemes of labelling are useless. There was recently¹ a report even in the non-medical journal *Which* highlighting the lack of implementation of the so-called voluntary scheme of labelling which had been agreed to for insecticidal preparations.

The "excellence" of the service made available to doctors in this country by the poisons information services, whether national, regional, or any other,² does not even begin to cope with the problem of accidental poisoning of children by commercial products in the home. These services come into the picture only after the poisoning has occurred. Parents in this country should receive the same consideration given to parents in other countries such as Sweden and the U.S.A., where the labels of the containers of commercial preparations containing dangerous substances have this fact clearly stated on the label in one form or another so that care can be taken to keep these particular preparations locked away well out of the reach of young children. These countries also put the name of the dangerous substance present on the label, while in addition Sweden adds a "poisons symbol" in the form of a skull.

In answer to the third point that the Home Office put forward it should be clearly stated and understood now that no practical difficulties would be presented if legislation requiring all toxic substances to be labelled is contemplated. It has been done in several countries including Sweden and the U.S.A., and in fact Sweden implemented its own legislation relating to this matter as recently as July 1967.

If the Home Office and its advisers find difficulty in producing lists of the dangerous substances which require cautionary labelling, then I for one, and I am sure many other doctors, would be very willing to help compile such lists. Incidentally, when firms export commercial products containing dangerous substances to such countries as Sweden and U.S.A. the firms comply with the rigid cautionary labelling requirements of these countries without any trouble. I have also had an example of a commercial preparation which was put out on to the British and American markets at the same time by one firm and found that those on the American market had a cautionary label on them while those on sale here had no cautionary label at all.

I sincerely hope that the Public Health Committee will continue to pursue this matter further with the Home Office until a satisfactory form of labelling is agreed to. My experience and that of my colleagues indicates that there would be overwhelming medical and public support for such a course of action by the B.M.A.—I am, etc.,

NOLAN A. WYNNE.

Poisons Information Centre,
Royal Victoria Infirmary,
Newcastle upon Tyne.

REFERENCES

- Davidson, S., *The Principles and Practice of Medicine*, 9th ed., 1968, p. 91. Edinburgh, Livingstone.
- Which*?, July, 1969, p. 202.

Authorship of Medical Publications

SIR,—In recent years the standards of scientific ethics of authorship of medical publications has fallen. There are two common sins. One is for the name of one or more of the major participants in the research to be either omitted completely from the paper or to be acknowledged at the end with a mere "thanks for help." The other is for the names of one or more authors to be attached to a paper when they may neither have taken an active part in the research nor collaborated in the writing, nor even seen the typescript submitted for publication. Even the order of names (which often does have significance) may not truly reflect the relative contributions of the authors. Occasionally papers are seen which obviously lean heavily on the work of other departments, but where no acknowledgement is made.

Unjustifiable investigations on humans have become fewer since editors of medical journals made the specific decision to refuse to publish articles which did not specify the informed consent of the subject. I am writing to suggest that medical editors could equally diminish, if not eliminate, the problems of authorship attribution. They should require at the submission of a paper for publication a declaration signed by *all* authors that they have approved the sub-

mitted typescript, and the names and order of the authors, and that they have not omitted the names of any participants in the work.

The victims of these offences usually find it difficult to complain publicly (or even privately) without grave embarrassment.—I am, etc.,

D. N. BARON.

Department of Chemical Pathology,
Royal Free Hospital School
of Medicine,
London W.C.1.

Contraceptives and Cervical Carcinoma

SIR,—I am obliged to Dr. Myron R. Melamed (1 November, p. 302) for pointing out my misinterpretation (23 August, p. 471) of Table I of the paper by him and his colleagues (26 July, p. 195). His explanation does indeed clarify the matter, which, I submit, was not put as clearly as it might have been, and which I suspect may well have been misinterpreted by others than myself.

However, even allowing for the doubts which could exist about the accuracy of recalled information on previous contraceptive practice by women before they entered the study, data on prevalence rates of carcinoma in situ at the time of entry into the study would still be of considerable interest. If, for example, they showed a significant difference between the two groups (those choosing oral contraceptives and those choosing the diaphragm), they would raise the suspicion of heterogeneity, such as I had, apparently mistakenly, inferred. Perhaps Dr. Melamed could provide the figures for prevalence, at the time of entry to the study, for the two groups of women concerned.—I am, etc.,

G. I. M. SWYER.

London N.W.1.

Sugar and Ischaemic Heart Disease

SIR,—Some of the points raised in the letter of Professor J. Yudkin and Dr. R. H. J. Watson (11 October, p. 110) demand an answer, notwithstanding the fact that 12 years after the publication of Professor Yudkin's hypothesis the only support he cites from trials in man is restricted to two papers from his own team. On the other hand, five very recent papers^{1–5} have failed to substantiate this hypothesis.

Professor Yudkin and Dr. Watson appear to attach great importance to the question of changing sugar habits with age. We considered that age-standardization would have introduced complications for readers, unjustifiably, in view of its insignificant effect in our series, where intake tended to fall slightly with age in both groups. A study of Table V in our article (19 July, p. 145) supports this view, and the Fig. giving mean weekly sugar intake by age again shows no pattern of increased consumption in the ischaemic heart disease (I.H.D.) group. The experience of those with unchanged sugar habits is also not significantly affected by age-standardization. Similarly, there were minor differences in sugar intake by social class, but since class-standardization did not substantially affect the results we again refrained from complicating the Tables. The Table overleaf quite clearly supports this view.

It cannot be too strongly emphasized that this study was not designed to discover variations in intake levels of low statistical significance. Not only is Professor Yudkin's questionnaire too blunt an instrument for this, but interpretation of such circumstances is difficult because of statistical variation and the intrusion of minor variables, known and unknown. The object, with no preconceived ideas to advance or defend, was to confirm or refute, with larger numbers and using a fairly "normal" population, Professor Yudkin's earlier finding of a very significant difference in sucrose intake between I.H.D. and control groups.

Mean Sugar Intake (g./week) in 1,158 Controls and 170 I.H.D. Patients by Age Grouping and Social Class

Social Class	Controls Age			I.H.D. Age		
	< 44	44-48	49 and Over	< 44	44-48	49 and Over
1	543	546	523	502	523	319
2	622	524	527	571	353	412
3	506	574	540	359	428	580
4 and 5	589	600	504	605	474	438

We would agree that measurement of sugar consumption over a lifetime cannot be precise, particularly if a questionnaire is the instrument. It appears to us, however, that the real significance of changing sugar habits with time (and hence with age) is that few people retain unchanged habits over a lifetime. Clearly there is a danger that those who remain consistent in their lifetime habits will be a minor, biased, sample from which it would be inappropriate to extrapolate results to cover the whole population. Yet Professor Yudkin and Dr. Watson would apparently restrict investigations to those with unchanged sugar habits.

Since neither our paper, nor any other recent article which has come to our notice has offered any convincing evidence that "high sugar intake is the chief dietary factor causing ischaemic heart disease," we thought it eminently reasonable to suggest that considerably more confirmation in man is required before the hypothesis is accepted. We did not "reject sucrose"—as they suggest—nor did we say the hypothesis is wrong. Since our study did not extend to include fats or other dietary factors we have no comment to make, though we did find that no correlation

existed between sugar consumption and serum cholesterol, β -lipoprotein, uric acid levels, or other biological tests. This important finding should not be overlooked.

If Professor Yudkin and Dr. Watson would have us speculate on dietary sugar and I.H.D. then we would suggest that rather than the critical individual level of intake apparently suggested by them there may be a population threshold of intake generally exceeded in the Western world. This may not be incompatible with some observations of Cleave and Campbell⁶ in their study of sugar, carbohydrates, and other dietary factors. Nor would it be inconsistent with recent papers¹⁻³ which failed to find individual susceptibility to I.H.D. by intake level.—We are, etc.,

R. W. HOWELL.

United Kingdom Atomic Energy Authority,
Harwell.

D. G. WILSON.

United Kingdom Atomic Energy Authority,
Dounreay.

REFERENCES

- Howell, R. W., and Wilson, D. G., *British Medical Journal*, 1969, **3**, 145.
- Burns-Cox, C. J., Doll, R., and Ball, K. P., *British Heart Journal*, 1969, **31**, 485.
- Paul, O., MacMillan, A., McKean, H., and Park, H., *Lancet*, 1968, **2**, 1049.
- Finegan, A., Hickey, N., Maurer, B., and Mulcahy, R., *American Journal of Clinical Nutrition*, 1968, **21**, 143.
- Finegan, A., Hickey, N., Maurer, B., and Mulcahy, R., *American Journal of Clinical Nutrition*, 1969, **22**, 8.
- Cleave, T. L., and Campbell, G. D., *Diabetes, Coronary Thrombosis and the Saccharine Disease*, 1966. Bristol, Wright.

Management of Unconscious Poisoned Patients

SIR,—Drs. N. Wright and J. A. Raeburn (13 September, p. 656) compared the duration of unconsciousness after poisoning with different types of barbiturates in terms of equivalent severity of poisoning. My remarks (19 July, p. 177) referred to the effects of equal amounts of drug. My own data, accumulated several years ago, are admittedly very limited and no doubt importantly influenced by the fact that all patients were treated energetically with forced alkaline diuresis from the time of admission. However, four patients admitted comatose, having taken an average of approximately 3 g. of phenobarbitone as the sole drug (other than alcohol), were in coma for an average of 18 hours. Ten patients having taken an average of approximately the same amount of pentobarbitone, amylobarbitone, or Tuinal (quinalbarbitone sodium and amylobarbitone sodium) as sole drug were in coma for an average of 28 hours.

Setter *et al.*¹ in their large series in fact quote an equal duration of unconsciousness for phenobarbitone and the alkyl-substituted barbiturates, the average quantity ingested, where known, being the same in the two groups. However, the amount ingested was known in only about half the cases, and in a considerable number other drugs were involved as well. These authors particularly remark on the fact that intoxication with "short-acting" drugs caused deeper and more rapid onset of coma and more severe complications than phenobarbital poisoning (my italics). Maher and Schreiner² emphasized the slower removal of "short-acting" (alkyl-substituted) barbiturates by haemodialysis. Bloomer³ emphasized also the relatively lesser efficacy of alkaline diuresis and

peritoneal dialysis in the case of pentobarbital intoxication. Thus differences in duration of effect are enhanced by active eliminative treatment.

There would seem therefore to be grounds for at least dubiety as to the longer action of phenobarbitone in cases of poisoning, and my contention is that there is little point in retaining a classification so based when there is a more logical way of classifying barbiturates on speed of onset of effect. This is valid at any dosage level and holds irrespective of subsequent events.

However, I would not wish controversy over this rather academic point to obscure the much more important one on which we are in agreement—namely, the desirability of abandoning the use of barbiturates as hypnotics at least for outpatients.—I am, etc.,

E. G. MCQUEEN.

National Poisons Information
Centre,
Dunedin, New Zealand.

REFERENCES

- Setter, J. G., Maher, J. F., and Schreiner, G. E., *Archives of Internal Medicine*, 1966, **117**, 224.
- Maher, J. F., and Schreiner, G. E., *Transactions of the American Society for Artificial Internal Organs*, 1965, **11**, 349.
- Bloomer, H. A., *New England Journal of Medicine*, 1965, **272**, 1309.

Treating Erythema Nodosum

SIR,—Erythema nodosum is often a very painful condition. The skin lesions themselves can be painful and tender, and there may be local muscle aching and tenderness associated with arteritis, panniculitis, or (especially in sarcoidosis) myositis. Skin lesions and muscle pains were rapidly relieved by oxyphenbutazone in the following case.

A 33-year-old woman was seen at home in May 1969 with erythema nodosum of the legs and pain sufficiently severe to prevent her standing for more than a few minutes. The skin lesions were preceded by arthralgia in the shoulders, ankles, and feet, but at the time of examination there was no joint swelling or tenderness. Neither aspirin, 40–50 mg., nor indomethacin, 75 mg., had been effective in relieving the leg pains. There was a history of pulmonary tuberculosis in childhood. She was admitted to the Princess Alexandra Hospital, Harlow, where the results of investigations were: Hb 92%, E.S.R. 21 mm./hr. (Westergren), W.B.C. 5,700/cu. mm. (differential W.B.C. normal), serum calcium and phosphate normal, protein electrophoresis: slight elevation of gamma globulin, S.G.O.T. 56 units, serum aldolase and creatinine phosphokinase normal, urine analysis normal, throat swab negative, A.S.O. titre=300 Todd units, Rose-Waaler negative, Mantoux 1:1,000 and 1:100 negative, Kveim test positive. A chest radiograph showed bilateral hilar adenopathy, consistent with sarcoidosis.

A diagnosis of erythema nodosum associated with sarcoidosis was made. She was initially treated with bed-rest and salicylates (soluble aspirin, 60 gr. daily), the skin lesions remained static, and she was given oxyphenbutazone, 400 mg., daily. Within four days the lesions had completely subsided and she was pain-free; stopping the oxyphenbutazone was followed by recurrence of the rash, so she continued the drug in a dosage of 300 mg. daily after discharge on 25 June. A chest