Middle Articles

CLINICAL PROGRESS

Deep Vein Thrombosis

In the articles below three contributors discuss recent progress in the diagnosis and medical and surgical management of deep vein thrombosis. Mr. Norman Browse concludes that not only is phlebography the most valuable method for diagnosis but that it may help correct management. Dr. P. T. D'Alute emphasizes that anticoagulants may be ineffective in preventing pulmonary embolism from recent thrombi in the popliteal or more proximal deep veins, and that fibrinolytic drugs are a possible alternative to surgery. Considering surgical management, Mr. G. E. Mavor illustrates the usefulness of classifying deep vein thrombosis as peripheral or iliofemoral.

Diagnosis

NORMAN BROWSE,* M.D., F.R.C.S.

British Medical Journal, 1969, 4, 676-678

Deep vein thrombosis, often associated with pulmonary embolism, is a common condition occurring in every type of patient—surgical, medical, and obstetric. It is not uncommon in apparently normal, active subjects, and women taking oral contraceptives are particularly susceptible. A simple method of making an accurate diagnosis in the early stages would be of considerable clinical value and also help those engaged in the study of methods of prophylaxis.

Clinical Examination

Once it was appreciated that pulmonary emboli came from the leg veins, and that even large lethal emboli could occur without any premonitory symptoms or signs in the legs, it became recognized that deep vein thrombosis could occur without physical signs. Even so, it is extremely difficult to look at a pair of apparently normal legs and actively consider the possibility that they are full of thrombus.

Physical signs depend on two distinct features of venous thrombosis. Swelling, venous distention, and an increase in temperature are due to a functional obstruction to the venous return from the legs. Pain and tenderness are apparently related to phlebitis—that is, an inflammatory response in the vein wall. Why phlebitis occurs with some deep vein thrombi and not with others is not known. The presence or absence of the phlebitis is certainly not related to the extent of the thrombosis. Thus an extensive deep vein thrombosis can occur *without* physical signs if the major veins are not completely blocked, if there are adequate collateral vessels around any totally occluded vessels, or if there is no phlebitis in the vein wall. At least half, probably two-thirds to four-fifths, of the patients who get deep vein thrombosis have no physical signs. One very important point arises from this situation; all of the many clinical investigations of deep vein thrombosis that have been based on physical signs for assessment are suspect, as they will not have ascertained the true incidence of the disease. A case could be made for saying that they are all valueless.

The above statements do not mean that physical signs *when present* are valueless. J. McLachlin and his colleagues showed by comparing physical signs with an objective means of diag-

nosis—necropsy examination—that ankle oedema was a reliable sign of underlying deep vein thrombosis in 80% of cases.¹ Other physical signs such as tenderness and Homan's sign were correct in only 50% and 8% of cases, respectively. These findings mean that some physical signs are not only rare, but they may be misleading. When a patient complains of pain and tenderness in the calf a definite diagnosis of deep vein thrombosis should not be made, as there is at least an 11% chance that the deep veins are normal.¹ It would be much more realistic to ignore physical signs and work on the assumption that one-third of all patients in hospital have thrombi in their deep veins.

The inadequacy of clinical examination has led many workers to search for objective means of ascertainment of the incidence of deep vein thrombosis. The oldest and perhaps the most accurate is extensive necropsy examination.² This method of course has a limited value and can only be used for clinical studies where, for some reason or other, there is a high mortality rate.³

Phlebography

Phlebography is the second oldest method of investigation. First described by Dos Santos in 1938,⁴ it was developed by him and the Swedish school⁵ during the early '40s. These workers proved phlebography to be an accurate and acceptable technique. Unfortunately, its value was not appreciated by clinicians and radiologists in Britain; only in the last five years has it been used here extensively as a diagnostic aid. One of the early problems of the technique was difficulty in recognizing streamline flow and poor mixing, a cause of false-positive results. The most valuable advance in diagnostic technique has been the use of television monitoring during the injection of the dye. The radiologist can now watch the veins filling and take films at the best moments rather than relying on the older method of taking exposures at set times after the injection.

Three methods of phlebography are in common use:

(1) Inject a radiopaque dye into a vein in the dorsum of the foot after occluding the superficial veins at the ankle with a rubber tourniquet (*ascending phlebography*). With this method it is possible to visualize the calf, popliteal, superficial femoral, common femoral, and iliac veins. The mouth of the profunda femoris vein is displayed if the patient performs a Valsalva manoeuvre.

* Reader in Surgery, St. Thomas's Hospital Medical School, London S.E.1.

(2) If the major veins in the upper thigh are obstructed ascending phlebography will not show the veins above the obstruction. The femoral and iliac veins and vena cava can be visualized by injecting a contrast medium directly into the femoral vein (*percutaneous femoral phlebography*).

(3) An excellent demonstration of the iliac veins, particularly the internal iliac vein, may be obtained when the common femoral vein is blocked by injecting the dye into the medulla of the upper end of the femur (*intraosseous petrocanteric phlebography*).⁶

The phlebographic diagnosis of deep vein thrombosis is based on the presence of well-defined filling defects in opacified veins, and on finding these defects on at least two radiographs.⁷ Recent thrombus appears as a rounded or cylindrical filling defect in a contrast-filled vein giving it a "ground-glass" appearance. The thrombus is separated from the wall of the vein by a thin white line representing contrast medium around the thrombus. Old thrombus shows as a filling defect projecting directly from the wall of the vein. Non-filling of a vein is accepted as a thrombotic occlusion only if the veins above and below the block are displayed, together with the collateral veins around it.

Some groups of veins cannot be displayed by phlebography. The venous sinuses in the calf are not usually seen, a serious deficiency, as they are a common site for deep vein thrombosis. Fortunately thrombosis in the muscle sinuses is usually associated with thrombus in the venae cavae of the arteries of the leg, which are easy to display. The tributaries of the profunda femoris vein are not seen unless the superficial femoral vein is blocked, but in my experience these vessels are not a common site for deep vein thrombosis. The tributaries of the internal iliac vein can be displayed only if the petrochanteric intraosseous technique is used, but again in my experience pelvic vein thrombosis is uncommon.

Despite these disadvantages phlebography probably displays 90 to 95% of existing venous thrombi, though there have been no careful studies comparing phlebography during life and necropsy findings. I think it is the most valuable investigation available; not only does it confirm the diagnosis, but it also gives a considerable amount of information on the exact site, extent, nature, and age of the thrombus, which can be extremely valuable when deciding on treatment. Nevertheless, unless the investigation is done with care by an enthusiast the results are poor. Hence research has also been carried out into other methods for detecting deep vein thrombosis.

Labelled-fibrinogen Uptake

In 1960 J. T. Hobbs and J. W. L. Davies described the preferential uptake of ¹³¹I-fibrinogen into fresh thrombi in rabbit leg veins,⁸ suggesting that this might form the basis of a useful clinical technique; subsequently this suggestion was confirmed.⁹⁻¹⁰ Subsequently the technique was satisfactorily modified to use an isotope with a longer half life—¹²⁵I.¹¹⁻¹²

Given intravenously 100 μ c ¹²⁵I-fibrinogen is incorporated as fibrin into any new thrombus that forms, and also passes into existing thrombus. The radioactivity can be detected with a scintillation counter. The routine of the test is to give the labelled fibrinogen, after blocking the uptake of iodine by the thyroid with potassium iodide, and then count the radioactivity at fixed points down the legs. The count rates in the legs are expressed as a percentage of the heart count. If one area shows a count rate of more than 15% above the adjacent areas, or than the same area on the other leg, then deep vein thrombosis may be confidently diagnosed. False-positive results occur if there has been any trauma to the leg or if there is gross oedema. Fig. 1 shows the counts in a man's legs seven days after ureterolithotomy. The phlebogram showed a 4 in. (10 cm.) long thrombus in the posterior tibial vein of the right leg. All workers in the field have shown over a 90% positive correlation between the fibrinogen uptake and phlebo-

graphy. In practice this technique shows a higher incidence of deep vein thrombosis than phlebography because it detects small thrombi in the calf sinuses more efficiently.

Fig. 1.—¹²⁵I-fibrinogen uptake. The radioactivity in the legs of a 63-year-old man seven days after a utero-lithotomy. The labelled fibrinogen has been taken up in a thrombus in the lower right calf giving count rates 20% higher than adjacent points and the same point on the opposite leg. Differences of 15% or more are diagnostic of deep vein thrombosis.

The biological half-life of the labelled fibrinogen is only three to four days, so that counting can continue for only six to seven days. Hence if the effect of treatment is being followed a booster dose of fibrinogen must be given. A second and more important disadvantage is that the technique cannot detect thrombi at the top of the thigh and in the iliac veins and vena cava. At the root of the leg and in the abdomen the background count from other tissues—and above the pubis the bladder—swamp any changes in the count rates from the veins. The fact that the ¹²⁵I-fibrinogen uptake is of doubtful value in the upper thigh and of no value at all above the inguinal ligament is a serious disadvantage, as the large loose thrombi that cause lethal pulmonary emboli often arise in the common femoral and iliac veins.

I have found the method to be particularly valuable for research, as it provides an objective method for detecting deep vein thrombosis, thus enabling accurate studies of treatment and prophylaxis. It is also valuable for the diagnosis of existing thrombosis. For example, in many patients presenting with pain in the chest and dyspnoea the differential diagnosis lies between pulmonary embolism or pneumonia; the detection of peripheral venous thrombi may be of considerable help in deciding on the correct diagnosis.

Ultrasound Techniques

When a beam of ultrasound is reflected from a moving object its frequency is altered according to the rate at which the object is moving, owing to the Doppler effect. The particles in blood act as reflectors, so that the frequency of an ultrasound beam passed through a moving column of blood is altered according to the rate of flow. This change can be recorded on paper or amplified into an audible signal. Because ultrasound is reflected off any interface between substances of different densities, a beam can be passed across a vessel, reflected off the distal wall or particles in the blood and picked up with a receiver adjacent to the transmitter. By using the ultrasound flowmeter to detect flow in a major vein one can discover whether the vessel is patent or blocked.¹³⁻¹⁵ The commonest cause of venous obstruction is acute thrombosis.

One method uses a 2MHz beam (Sonicaid Ltd.)¹⁵; the major veins along the leg are examined by placing the ultrasonic probe over a vein and increasing the flow by squeezing the vein up-

stream to the transmitter. Fig. 2 shows the type of tracing obtained with a complete block of the deep veins of the left leg.

FIG. 2.—The ultrasound flowmeter record of a man with a total left deep vein block. Squeezing the calf or thigh on the right side increases the flow and produces an audible noise, recorded as an upswing (A wave) on the record. Squeezing the left leg produced no change in flow indicating a blocked vein. (Reproduced by kind permission of the authors, D. S. Evans and F. B. Cockett, *British Medical Journal*, 1969, 2, 802.)

This method is quick and extremely simple and studies the important source of *major* emboli—the femoral and iliac veins and the vena cava. On the other hand, a disadvantage of the method is that flow in large collaterals or in a very large superficial vein will produce a sound similar to that produced by flow in a patent femoral vein. These circumstances may produce a false-negative reading. Early thrombi in the femoral vein sometimes propagate along the vein without producing any significant obstruction, and this type of thrombus may also be missed. Small thrombi in the calf veins, the tributaries of the profunda vein, or the internal iliac veins cannot be detected because they do not cause major vein obstruction. Despite these drawbacks, however, ultrasound is a valuable and rapid method of examining the major veins of a leg for large occlusive thrombi.

Perhaps some caution should be exercised while squeezing the leg to produce the increase in flow, as young thrombi are known to be very loose and friable; theoretically a thrombus might be broken free if the leg is squeezed too vigorously.

When the absence of flow suggests an occlusion of a major vein a phlebogram must still be performed because possibly the occlusion is old and not due to fresh, loose thrombosis. Before proceeding to any form of treatment more detailed information is needed concerning the nature and extent of the thrombus.

Selection of Tests

Phlebography is undoubtedly the most accurate technique; it is essential before any surgical treatment and is important and helpful before deciding on medical treatment, such as thrombolysis.

^{125}I -fibrinogen is a very accurate technique for detecting thrombus below the upper thigh, but there is a delay of 6–24 hours after giving the fibrinogen before the count rates are of diagnostic value. However, the results from using this method may be available earlier if phlebography cannot be carried out immediately, and the whole technique can be carried out by a technician. I use the method extensively for *confirming* the diagnosis of calf vein thrombosis and no longer treat any patient solely on the basis of a clinical diagnosis, because some patients with pain and tenderness in the calf do not have venous thrombosis. Once the diagnosis is made, daily leg scans will show whether the thrombosis is progressing or regressing, thus providing an objective assessment of the effectiveness of treatment.

The ultrasound flowmeter is the simplest of the screening methods but is applicable only to the major veins.

My current policy is to divide patients with deep vein thrombosis into two groups: those with pulmonary embolism and those without. If the patient has had an embolus I order an emergency phlebogram to obtain the maximum information as soon as possible so that further emboli can be prevented.¹⁶ I approach the patient with a suspected deep vein thrombosis, but no pulmonary embolus, with both the ultrasound flowmeter and a syringe of labelled fibrinogen. If the flowmeter shows the large veins to be clear I give the fibrinogen, scan the legs the following day, and then begin treatment if the diagnosis is confirmed. The legs are observed daily with the scintillation counter, and if in spite of treatment there is any extension of the thrombus I order a phlebogram. If the flowmeter shows a major vein block phlebography is carried out immediately in case the obstructing thrombus is fresh and loose and amenable to thrombectomy.

I believe that these three techniques, which any hospital can undertake, enable us to treat deep vein thrombosis in a logical and objective manner. They should also eventually produce a tremendous increase in our knowledge and understanding of the aetiology, natural history, and treatment of this common but very serious disease.

Medical Management

P. T. FLUTE,* M.D., M.R.C.PATH.

British Medical Journal, 1969, 4, 678–680

To understand the action of anticoagulants and fibrinolytic drugs the whole process of thrombosis must be viewed as the end result of the opposing action of two forces, one leading to the deposition and the other to the removal of fibrin. Disorders of blood flow and of the vein walls, platelet adhesion and aggregation, and blood coagulation contribute to the deposition of fibrin. The fibrinolytic enzymes and cells of the reticulo-endothelial system contribute to its removal. Hence, paying attention to any one of these variables may be valuable in prophylaxis.

Once formed, fibrin may dissolve completely under the influence of the fibrinolytic enzymes or become organized with

permanent damage to the veins. The true incidence of spontaneous thrombolysis is not yet known, but it has been found in a surprisingly high proportion of patients early in the post-operative period.¹⁷

Anticoagulants restrict the further deposition of fibrin but have no direct action on an established thrombus. Nevertheless, they may so limit its extension that they improve the chance of spontaneous thrombolysis. In assessing this objective methods must be used to measure thrombi, for recovery from clinical signs and symptoms often follows the opening of new collaterals while the thrombus remains intact.

The fibrinolytic drugs have a direct thrombolytic effect, which is achieved either by activating the fibrinolytic enzyme system or by an independent proteolysis of fibrin.

*Reader in Haematology, King's College Hospital Medical School, London S.E.5.