Like most of us, the young doctor is faced with the painful necessity of having to earn a living, particularly as he is likely to be married and often has a family to support. After, say, two years' work, he can enter general practice, an interesting and rewarding life with much variety, or decide on radiology as a career. In the former case he will have a fairly wide choice of district and start at £3,000-£4,000 p.a., achieving a high salary in the not too distant future. If he should choose radiology he will start as registrar for two years or more, obtain his D.M.R.D. and then become senior registrar for three years, altogether about five years' training. He can then take a further qualification if he wishes.

He will then start as a consultant and after 10 years or more reach the maximum salary. This will still be less than in general practice, unless he is fortunate enough to obtain a merit award or a lucrative private practice, both of which are rare in radiology.

What does the future hold for radiology in the United Kingdom? If things go as at present, it is a dying specialty. It is true many are in training, but how many will remain in Britain under the present system?—I am, etc.,

GEOFFREY MEDHURST.

General Hospital,
Kettering, Northants

ABO Blood Groups and Myelomatosis

SIR,—Dr. D. Catovsky and his colleagues (22 August, p. 438) have reported a 3.2% current incidence of thromboembolic disease in the M.R.C. myelomatosis therapeutic trial series, and they believe that this percentage will increase. On the other hand, each of three series of patients with thromboembolic disease in general¹⁻³ has shown a significant deficit of blood group O and excess of A. It is therefore worthy of note that there is an appreciable, though far from significant, deficit of O and excess of A in a series of 120 North-east Scotland cases of myelomatosis blood-grouped prior to transfusion in 1955-70 (see Table).

As many more years would have to elapse before this result could be either refuted or confirmed in North-east Scotland, with its very small population, it is to be hoped that further series will be compiled by others with access to much larger numbers. In this connexion, however, concerted action^{2,5} would clearly be particularly desirable, because of the uncommonness of myelomatosis, plus the strength of Wiener's⁶ argument that the study of the blood groups in relation to disease is called in question, to an unknown extent, by the putative non-publication of negative findings. The same would be even more true in respect of amyloidosis, whose relationship to myelomatosis and thromboembolism is discussed by Dr. Catovsky and his colleagues.

The possibility of a blood group association between thromboembolism and cancer

was suggested by Dick *et al.*,¹ who, referring to the relationship of thrombosis and cancer in general, pointed out that the significant excess of A in their series of thromboembolism cases was confined to the subgroup A₁, and that the same had been true in respect of their series of stomach cancer cases. Accordingly it is interesting that there is not only "evidence of some strength" of an association between group A and cancer of the pancreas,⁵ but also some evidence of a particularly high incidence of thrombosis (single or multiple) in cancer of the body or tail of the pancreas.⁷ Thus it is possible that, from the standpoint of ABO blood group distribution, a division of cases by site and type, not to mention secretor status, would be as profitable in pancreatic cancer as it has been in peptic ulcer.—I am, etc.,

T. M. ALLAN.

Blood Transfusion Centre,
Royal Infirmary,
Aberdeen.

REFERENCES

1. Dick, W., Schneider, W., Brockmüller, K., and Mayer, W., *Thrombosis et Diathesis Haemorrhagica*, 1963, 9, 472.
2. Jick, H., *et al.*, *Lancet*, 1969, 1, 539.
3. Talbot, S., Wakley, E. J., Ryrie, D., and Langman, M. J. S., *Lancet*, 1970, 1, 1257.
4. Kopeć, A. C., *The Distribution of the Blood Groups in the United Kingdom*, p. 34. London, Oxford University Press, 1970.
5. Aird, I., Lee, D. R., and Roberts, J. A. F., *British Medical Journal*, 1960, 1, 1163.
6. Wiener, A. S., *Journal of Forensic Medicine*, 1960, 7, 166.
7. Sprout, E. E., *American Journal of Cancer*, 1938, 34, 566.

ABO Blood Group Distribution in Myelomatosis in N.E. Scotland, 1955-70

ABO Blood Group	Number of Myelomatosis Patients in N. E. Scotland, 1955-70				
	Observed	Expected ^a	Difference (O. minus E.)	Ratio O./E.	(O. E.) ^a E.
A	53	42.4	+10.6	1.25	2.64
AB + B	16	16.6	0.6	0.96	0.02
O	51	61.0	-10.0	0.84	1.64
Total	120	120.0	Nil	1.00	4.30

$$\chi^2 = 4.30 \text{ for } 2 \text{ d.f.; } p > 0.1$$

Internal Sphincterotomy

SIR,—For almost two years we have been using lateral subcutaneous internal sphincterotomy for fissure-in-ano and our results¹ are similar to those of Mr. D. C. Hoffmann and Professor J. C. Goligher (19 September, p. 673). We wish to make some

observations from our experience.

As an alternative to the von Graefe cataract knife a no. 11 Bard-Parker blade may be used and indeed is often more readily available. We find it entirely satisfactory. Like Notaras² we divide the sphincter

from within outwards. The point of the knife-blade is introduced in the perianal region and advanced under the skin towards an anal papilla at the base of a column of Morgagni. Thus perforation into an anal crypt is avoided. Then by cutting laterally that part of the internal sphincter below the dentate line is firmly divided. Conversely when cutting towards the anal lumen, as advocated by Mr. Hoffmann and Professor Goligher, we find mucosal perforation more common. However, a small perforation is not serious. It heals without fistula formation. It is neither necessary to suture the defect nor to convert it to an open sphincterotomy.

In our clinic local infiltration anaesthesia with 0.25% bupivacaine containing adrenaline and hyaluronidase is used for many anorectal operations. This gives several hours of complete postoperative freedom from pain. Furthermore, outpatients may go home immediately after surgery, thus reducing the work load on nursing staff.—We are, etc.,

ANTHONY P. CLERY.

MICHAEL FEELY.

St. Lawrence's Hospital,
Dublin 7.

REFERENCES

1. Clery, A. B., and O'Riordan, J. B., *Journal of the Royal College of Surgeons in Ireland*, 1970, 6, 10.
2. Notaras, M. J., *Proceedings of the Royal Society of Medicine*, 1969, 62, 713.

Varicella and Herpes Zoster

SIR,—Varicella was at one time regarded as a relatively benign childhood ailment, but it is becoming increasingly realized that the virus of varicella may lie dormant in the central nervous system for many years to erupt later as herpes zoster, a much more serious complaint that causes great pain and suffering. Apparently this is associated with a waning of immunity to the varicella virus.

Surely the time is ripe for a vaccine to be made available to adults which could boost their immunity and so reduce the risk of their ever developing shingles.—I am, etc.,

T. H. GILLISON.

Ilkeston, Derbyshire.

Teratogenic Drugs—R.C.G.P. Survey

SIR,—In a leading article (14 March, p. 642) a survey conducted by the Royal College of General Practitioners was mentioned as a source of further information about possible relationships between aspirin consumption by women in the early weeks of pregnancy and subsequent congenital malformation in the child.

The college survey, now completed, is based on a study of approximately 10,000 women. Both self-administered drugs and prescriptions issued to these women in the period from six weeks before the last menstrual period to 22 weeks after the last menstrual period were recorded. The numbers given below, unless otherwise stated, refer to individual prescriptions by the practitioner or to reports by the woman to her practitioner of any drugs she has taken herself. The numbers of expected prescriptions based on the numbers issued to all women who reached term in the corresponding drug categories are in brackets.