"safe and effective prophylactic," and because "50% of acute infarcts develop major ventricular dysrhythmias."

(4) Consider intravenous atropine for bradycardia.

(5) If the pulse is irregular with runs of ectopic beats 50 mg. of lignocaine slowly intravenously and the rest of the syringe pack intramuscularly.

(6) Finally, beseech the ambulance men to give a smooth, steady, clammer-free ride to Trellis's coronary care unit.—I am, etc.,

R. D. MARTIN.

Newquay,
Cornwall.

Bell's Palsy and Herpes

SIR,—In your leading article on Bell's palsy and surgery (17 October, p. 126), you twice say that surgical decompression for the relief of the condition is controversial. I agree that this is so, and would indeed echo Professor Ritchie Russell's teaching that if you want to succeed with a decompression operation you must operate before the onset of the lesion. There is no magic moment after the paralysis has occurred.

Where the controversy comes in, however, is in relation with imperfect diagnosis. Too often a herpetic lesion is wrongly categorized as a Bell's palsy, and creeps into the statistical surveys on one side or the other. Bell's palsy is characterized by sudden and only slightly progressive paralysis, without much pain. In facial nerve herpes there is severe faceache which often precedes the paralysis by as much as 24 hours. The paralysis at its onset is usually incomplete, but it progresses over the next 24-48 hours.

My experience has been that this latter is the kind of case which has a bad prognosis if left alone and which rewards decompression. The time to do this is while the paralysis is progressing. Although the lesion may well be ganglionic basically, the nerve sheath becomes very inflamed, with vascular engorgement and an almost shaggy appearance. This is different from the "swollen spaghetti" look of the Bell's palsy nerve.

While agreeing that many herpetic cases do recover on masterly inactivity, I would urge that surgery should be thought about early and undertaken if the paralysis is progressing. But the differential diagnosis is the main issue, and here the virologist may help.—I am, etc.,

RONALD MACBETH.

Oxford.

Children in Intensive Therapy Units

SIR,—Drs. J. R. Harper and S. Varakis (13 June, p. 670) have pointed out that any hospital dealing with acutely ill children must solve its problems according to the local conditions. I strongly support the view that in a peripheral hospital the nursing of children in general intensive care units is a forward and not a retrogressive step.

My own service looks after a population of over 300,000. During the past 14 years we have been able to engage only two nurses with paediatric nursing qualifications; we would have been glad of others, but they are just not available.

The intensive care unit at Wigan Infir-

mary has now been in operation for just over three years. A total of 700 patients have been admitted and of these 50 (7%) have been children aged 12 and under. A very wide range of ages and of clinical conditions have been handled. We are fortunate that the unit is situated a few yards from the main children's ward, so that a mother living-in with her child in the hospital can use one of the rooms available on the children's ward. The transfer of a child to or from the intensive care unit at the appropriate time is also very easy.

I have not inquired in detail from the families involved what they thought about their experiences. This would be interesting, but in my case largely irrelevant. I am certain that the intensive care unit has saved lives which would otherwise have been lost, and the doubts of any of the families would pale into insignificance compared with the certainty of the medical and nursing staff of the hospital that the intensive unit has been invaluable.

While I agree with Drs. W. J. Appleyard and M. C. Joseph (16 May, p. 423) that the trend should be towards nursing children in children's units, it must never be at the expense of safety and technical efficiency. In the average peripheral hospital the staffing situation would seldom reach the levels to allow these standards of intensive care on the children's ward itself, and the paediatrician would find it very hard to justify such a concentration of nurses and equipment in competition with his colleagues in other disciplines, with whom these rare and expensive facilities can easily and effectively be shared.—I am, etc.,

R. M. FORRESTER.

Royal Albert Edward Infirmary,
Wigan, Lancs.

Prophylaxis of Postoperative Pulmonary Atelectasis

SIR,—Space permits us to comment on only a few of the points raised by Mr. M. O'Driscoll (3 October, p. 26). In brief, we find the design of his experiment inadequate, confused, and based on a false premise, the methods unsound, the results and their interpretation contradictory, the discussion untenable, and consequently the summary misleading.

Of the eight references that Mr. O'Driscoll cites in his first paragraph five have been misrepresented. Three of these authorities did not make the statements ascribed to them; two are quoted out of context and without those authors' reservations.

We are told that the study group consists of 181 patients divided into two unmatched groups. In Table I a further 116 patients mysteriously appear. 235 of Mr. O'Driscoll's patients received halothane as an anaesthetic while a further 52 were given Fluothane (halothane).

The latter part of this paper concentrates on those patients who did not receive bicarbonate and in whom chest complications were said to have occurred. It is claimed that a "particularly pronounced" metabolic acidosis was seen in these patients. Table III shows this group to have been alkalotic with a mean pH of 7.43. To describe a mean base deficit of less than 4 mEq/l. as diagnostic of marked metabolic acidosis shows a lack of appreciation of the normal

variations in acid-base state. Deductions made on the basis of isolated serum venous bicarbonate levels (presumably performed by an autoanalyser technique) are totally meaningless. These patients had a mean PCO_2 of 27.6 mm. Hg. Mr. O'Driscoll is most reluctant to ascribe this respiratory alkalosis to increased alveolar ventilation. He believes that this hypocarbia demonstrates a reduction in physiological dead space and a probable reduction in carbon dioxide production. This is not so. Alveolar dead space is normally small and even its reduction to zero would have an insignificant effect upon arterial PCO_2 . Anatomical dead space could hardly decrease by the necessary order implied in the data. A reduction in CO_2 production of the order of 73% would be required in order to produce the changes in PCO_2 seen. In any event falls in both CO_2 production and in dead space are usually accompanied by hypoventilation in order to maintain normal alveolar PCO_2 values.

Because of the known difficulties in measuring ventilation with a respirometer and face mask, the known errors of blood-gas analyses performed on arterialized venous blood, the lack of details of any correction of acid-base data for changes in temperature or oxygen saturation, and the apparent confusion between Astrup and venous bicarbonates, we feel that the most likely explanation of Mr. O'Driscoll's findings is a methodological error.—We are, etc.,

C. M. CONWAY.

J. M. LEIGH.

Department of Anaesthetics,
Westminster Hospital,
London S.W.1.

Allergy to Iprindole (Prondol) with Hepatotoxicity

SIR,—We wish to add four further cases of iprindole-induced hepatitis to those previously reported in your correspondence columns (7 February, p. 367, and 25 April, p. 238).

Case 1.—Female aged 48 years. Iprindole prescribed for mild depressive symptoms. Hb and E.S.R. normal. Many years ago she had taken amitriptyline for similar symptoms without ill effects.

Seven days after starting iprindole noticed definite improvement in mood. On the eighth day sudden onset of "influenzal" symptoms—shivering, back pain, malaise, anorexia, and headache. The patient took her own temperature and found it to be 104°F. (40°C.). At this point she stopped the drug. The following day she developed pain in the right hypochondrium and passed dark urine.

When examined on the tenth day after starting iprindole the patient was found to be mildly jaundiced and tender beneath the right costal margin. The temperature was not raised. Urine contained bile on inspection, and this confirmed by Ictotest.

Liver function tests taken the following day were as follows: Serum bilirubin 1.1 mμ/100 ml.; S.G.O.T. 36; S.G.P.T. 93 units/ml.; cephalin cholesterol flocculation positive; colloidal gold flocculation negative; alkaline phosphatase 20 K.A. units. Albumin 3.4; globulin 3.1; total serum protein 6.5 g./100 ml.; zinc sulphate turbidity 5 units.

Jaundice rapidly disappeared and bilirubin was no longer present in the urine four days after its first detection.

The patient made a rapid and complete recovery. Cholecystography showed a normal gall bladder.

Case 2.—Female aged 23 years. Iprindole prescribed for postinfluenzal depression. Hb and E.S.R. normal. Eight days after first taking the drug suddenly developed pain in the right hypochondrium and right posterior chest with transient fever (temperature taken by the patient and found to be 101°F. (38.3°C.)).

When seen on the tenth day, apyrexial, marked tenderness present in the right hypochondrium, definite clinical jaundice. Urine strongly positive for bile.

Liver function tests taken the same day were as follows: Serum bilirubin 1.0 mg./100 ml.; S.G.P.T. 45 units/ml.; cephalin cholesterol flocculation positive; colloidal gold flocculation negative; alkaline phosphatase 23.5 K.A. units; zinc sulphate turbidity 8 units; total serum proteins 7.8; albumin 4.0, globulin 3.8 g./100ml.

The patient made a rapid recovery, jaundice disappearing by the twelfth day and bilirubinuria clearing by the fourteenth day. Reticulocyte count and blood film were normal and liver function tests substantially unchanged on the twelfth day after beginning the drug.

Case 3.—Female aged 30 years. Iprindole prescribed for anxiety-depressive symptoms, as amitriptyline caused drying of the mouth. Eight days after starting the drug she developed upper abdominal pain, fever ("felt as if I had 'flu'"), and flatulence.

When seen the following day apyrexial, not icteric, slight hepatic tenderness, urine negative for bile. Iprindole was discontinued.

Liver function tests the same day: Bilirubin 0.4 mg./100 ml.; S.G.P.T. 52; cephalin cholesterol flocculation positive; colloidal gold flocculation negative; alkaline phosphatase 11.5 K.A. units; zinc sulphate turbidity 6 units; total serum protein 7.4; albumin 4.2, globulin 3.2 g./100 ml.

By the thirteenth day the patient felt well. There were no abnormal signs. All the liver function tests were essentially unchanged.

Case 4.—Female aged 38 years. Developed upper abdominal pain 11 days after starting iprindole for depression.

Examination at this time revealed tenderness in the right hypochondrium but no jaundice or fever. Urine negative for bilirubin. Liver function tests not done. Iprindole was discontinued and all symptoms subsided completely over the next four days. This was our first case, and the significance of her symptoms was appreciated only in retrospect.

We suggest that these patients exhibit similar features to those already reported. They show evidence of hepatic dysfunction about one week after starting iprindole. In view of the short time that this drug has been in general use the evidence of its hepatotoxicity which has already appeared is disturbing. As alternative tricyclic drugs are available with little or no evidence of adverse reactions, we would suggest that iprindole be withdrawn by the manufacturers.—We are, etc.,

D. F. HARRISON.

I. M. STANLEY.

Sherburn-in-Elmet,
Leeds.

Pulmonary Failure after Peripheral Injury

SIR,—I have experience of two cases of the condition described in your leading article (3 October, p. 5). The patients were previously healthy young people suffering from multiple fractures sustained in road accidents. In both cases the initial shock was fully corrected by transfusions of blood

and low molecular weight dextran but they had not been overtransfused.

In one case the onset of haemorrhagic pulmonary oedema was very dramatic and the patient literally drowned in a matter of minutes in the haemorrhagic fluid which welled up from the trachea. In the other case the condition was corrected by an immediate venesection and administration of diuretics and oxygen. In neither case had gas mixtures with a high concentration of oxygen been given in the early stages.

The administration of low molecular weight dextran would seem to be a dangerous procedure once pulmonary oedema is established. Low molecular weight dextran is hyperoncotic and leads to a further increase in the circulating blood volume.

The part played by fat embolism in the aetiology of this condition may be considerable, but the syndrome differs so markedly from the usual picture of fat embolism that it would appear to be fundamentally different in some ways.—I am, etc.,

J. A. O'GARRA.

Guest Hospital,
Dudley.

Vasculitis in Coeliac Disease

SIR,—We were interested in Professor C. C. Booth's description of arteritis complicating coeliac disease (3 October, p. 14), as we have under observation a patient with coeliac disease with many unusual manifestations, including evidence of vasculitis in the skin.

A housewife aged 50 developed symptoms of anaemia in 1967. This was investigated elsewhere and evidence was found of folate and iron deficiency. Later she developed oedema due to hypoalbuminaemia and coeliac disease was suspected on the basis of low D-xylose excretion. She was placed on a gluten-free diet but did not adhere to this strictly.

She was referred to one of us (S.O.) in October 1968 very ill with weight loss, diarrhoea, and gross ascites and oedema. Investigations showed evidence of malabsorption (faecal fat 15 g./day, D-xylose excretion 2.7 g. in 5 hours), and jejunal biopsy showed subtotal villous atrophy. As in Professor Booth's patient, the enterocytes were flat and in some places absent, and there was prominent collagen beneath them. Ileal involvement was suggested by a serum B₁₂ level of 50 µg./ml. There was hypo-proteinaemia (serum albumin 2.1 g./100 ml., globulin 1.4 g./100 ml.) and labelling of the serum proteins with ⁵¹CrCl₃ showed this to be due to protein-losing enteropathy. After intravenous injection of 100 µCi ⁵¹CrCl₃, 25% of the dose was lost in the stools in 5 days (normal <1%) and plasma half-life of the isotope was 1.5 days (normal 9.5 days¹).

She was treated with a gluten-free diet, replacement therapy, and diuretics, but remained critically ill. In view of the protein-losing state and ileal involvement prednisolone 30 mg./day was added to her regimen, and this caused a remarkable improvement, with clearing of the oedema and ascites and a rise in serum proteins (albumin 3.7 g./100 ml., globulin 2.2 g./100 ml.).

Repeat jejunal biopsy in March 1969 showed improvement, with only partial villous atrophy. Prednisolone dosage was therefore reduced to 15 mg./day with no symptomatic change, but there was a fall in serum proteins and evidence of protein-losing enteropathy persisted (5-day ⁵¹CrCl₃ loss in stools 9% of dose). In June 1969 she developed a dermatosis which rapidly became generalized; this was thought to be pemphigus erythematosus (Dr. G. Auckland). Skin biopsy

showed a chronic inflammatory infiltrate closely applied to the basal layer of epidermis and also around capillaries in the dermis. Treatment with local hydrocortisone and increase in oral prednisolone to 40 mg./day produced a slow improvement in her condition, but during this time she had several grand mal convulsions. These were controlled with phenobarbitone.

The rash eventually cleared completely and her condition has remained very satisfactory apart from occasional fits. Serum proteins returned completely to normal (albumin 4.5 g./100 ml., globulin 2.2 g./100 ml.). She remains on a strict gluten-free diet, folic acid 5 mg. t.d.s., prednisolone 30 mg./day, and phenobarbitone 30 mg. t.d.s.

The diagnosis of coeliac disease is not fully established in this patient in that challenge with gluten has not been tried. In view of the severity of the initial illness and deterioration on reduced steroid dosage we have not felt justified in attempting this test. However, her clinical picture at the onset seems characteristic and her well-being at the present stage makes it unlikely that she has a lymphoma of the small intestine. There is undoubted evidence of vasculitis in the skin biopsy, similar to that of lupus erythematosus, but L. E. cells were not found and antinuclear factor was absent. Numerous dermatoses have been described in association with coeliac disease;² to our knowledge, however, pemphigus erythematosus has not been previously reported.

Neurological manifestations of coeliac disease, including convulsions, are uncommon and their pathogenesis unclear. Pyridoxine deficiency has been suggested,³ but a course of this in our patient had no effect on the E.E.G. or on the frequency of the fits. It is conceivable that they could be a manifestation of vasculitis.—We are, etc.,

D. J. HOLDSTOCK.

S. OLEESKY.

M3 Unit,
Royal Infirmary,
Manchester.

REFERENCES

- 1 Walker-Smith, J. A., Skyring, A. P., and Mistilis, S. P., *Gut*, 1967, 8, 166.
- 2 Wormsley, K. G., *The Skin and Gut in Disease*. London, Pitman, 1964.
- 3 Morris, J. S., Ajdukiewicz, A. B., and Read, A. E., *Gut*, 1970, 11, 549.

Other Side of Table Mountain

SIR,—Leaving aside Dr. Joyce E. Leeson's rather unfair comment (12 September, p. 648) that "Professor Jackson would not know about that"—"that" being tuberculosis and malnutrition among the non-White population—I should like to ask her just one question. Does she think that doctors who leave South Africa "for the sake of their principles" do more good than those who stay and work among and for the non-White people to the best of their medical ability?

Perhaps some doctors who hold Dr. Leeson's views should join the splendid man she mentions in her letter (Dr. Guy Daynes in the Transkei).—I am, etc.,

W. P. U. JACKSON.

Department of Medicine,
University of Cape Town,
South Africa.