- Braude, A. I., Carey, F. J., and Zalesky, M. (1955). *Journal of Clinical Investigation*, 34, 858.
- Campbell, R. M., Cuthbertson, D. P., Matthews, C. M., and McFarlane, A. S. (1956). *International Journal of Applied Radiation and Isotopes*, 1, 66.
- Caroli, J. (1958). *Revue Médico-Chirurgicale des Maladies du Foie, de la Rate et du Pancréas*, 33, 187.
- Chedid, L., Parant, M., and Parant, F. (1970). *Journal of the Reticuloendothelial Society*, 7, 238.
- Crosnier, J. (1958). *Revue Médico-Chirurgicale des Maladies du Foie, de la Rate et du Pancréas*, 33, 186.
- Dawson, J. L. (1968). *Annals of the Royal College of Surgeons of England*, 42, 163.
- Farrar, W. E., Eidson, M., and Kent, T. H. (1968). *Proceedings of the Society for Experimental Biology and Medicine*, 128, 711.
- Funck-Brentano, J. L., Méry, J. Ph., Vantelon, J., and Watchi, J. (1963). *Presse Médicale*, 71, 1039.
- Good, R. A., and Thomas, A. L. (1952). *Journal of Experimental Medicine*, 96, 625.
- Gronvall, J. A., and Brunson, J. G. (1956). *Archives of Pathology*, 62, 324.
- Hamburger, J. (1958). *Revue Médico-Chirurgicale des Maladies du Foie, de la Rate et du Pancréas*, 33, 181.
- Hepp, J. (1958). *Revue Médico-Chirurgicale des Maladies du Foie, de la Rate et du Pancréas*, 33, 184.
- Herring, W. B., Herion, J. C., Walker, R. I., and Palmer, J. G. (1963). *Journal of Clinical Investigation*, 42, 78.
- Hjort, P. F., and Rapaport, S. I. (1965). *Annual Review of Medicine*, 16, 135.
- Huth, K., Schoenborn, W., and Knorpp, K. (1967). *Thrombosis et Diathesis Haemorrhagica*, 17, 129.
- Kliman, A., and McKay, D. G. (1958). *Archives of Pathology*, 66, 715.
- Lázár, G., and Karády, S. (1965). *Journal of Pharmacy and Pharmacology*, 17, 517.
- Lázár, G., Karády, S., and Husztik, E. (1969). *Thrombosis et Diathesis Haemorrhagica*, 21, 159.
- Lee, L. (1962). *Journal of Experimental Medicine*, 115, 1065.
- Lendrum, A. C., Fraser, D. S., Slidders, W., and Henderson, R. (1962). *Journal of Clinical Pathology*, 15, 401.
- Lerner, R. G., Rapaport, S. I., and Spitzer, J. M. (1968). *Proceedings of the Society for Experimental Biology and Medicine*, 127, 904.
- Lipiński, B., and Warowski, K. (1968). *Thrombosis et Diathesis Haemorrhagica*, 20, 44.
- McKay, D. G., Margaretten, W., and Rothenberg, J. (1964). *Laboratory Investigation*, 13, 54.
- Nolan, J. P., and Ali, M. V. (1968). *Proceedings of the Society for Experimental Biology and Medicine*, 129, 29.
- Shaldon, S., Chianussi, L., Guevara, L., Caesar, J., and Sherlock, S. (1961). *Journal of Clinical Investigation*, 40, 1346.
- Thomas, L., and Good, R. A. (1951). *Proceedings of the Society for Experimental Biology and Medicine*, 76, 604.
- Vassalli, P., and Richet, G. (1961). *Proceedings of the First International Congress of Nephrology*, ed. G. Richet, p. 236. Basel, Karger.
- Weil, M. H., MacLean, L. D., Visscher, M. B., and Spink, W. W. (1956). *Journal of Clinical Investigation*, 35, 1191.

Medical Memoranda

Neuropathy, IgM Paraproteinaemia, and Autoantibodies in Hypernephroma

British Medical Journal, 1970, 4, 474

In a recent review of 72 patients (Croft and Wilkinson, 1969) neurological symptoms preceded the finding of malignancy in 49%, the mean interval before discovery of the neoplasm being two years. In the case reported here neurological symptoms preceded diagnosis of hypernephroma by four and a half years.

CASE REPORT

A 58-year-old timber salesman had been in good health until 1964, when he developed weakness of his left hand associated with numbness of the ring and little fingers. A few weeks later he noticed weakness of his right foot and intermittent paraesthesia over the dorsum of his left foot.

On examination he was normotensive and had a left ulnar nerve palsy, a right lateral popliteal palsy, absent reflexes in the legs, and flexor plantar responses; vibration sense was diminished at both ankles. The clinical signs were those of mononeuritis multiplex in the context of a more generalized peripheral neuropathy. All investigations performed at that time were normal.

In July 1967 he was admitted to hospital for further assessment. Apart from the development of wasting, his physical signs remained unchanged, but investigations showed several new features. His E.S.R. had risen to between 42 and 60 mm. in the first hour and the C.S.F. protein to 80 mg./100 ml. All other investigations, including biopsy of the right sural nerve and underlying muscle, were normal.

During the next 18 months there was progressive deterioration with increasing weakness and general malaise. Examination in October 1968 disclosed bilateral wasting of the interossei muscles and hypothenar eminences associated with generalized weakness of the arms, most pronounced distally. There was global wasting of the right leg with gross distal weakness and associated foot drop. Power in the left leg was normal. All the reflexes were absent except the left triceps reflex, which was brisk; the left plantar response was extensor and the right plantar response produced fanning of the toes. A sensory loss in the territory of the ulnar nerve was present in both hands, and there was impairment of vibration and joint position sense at both ankles. Romberg's sign was positive and his gait ataxic.

Investigations.—Haemoglobin 112%; E.S.R. 68–71 mm. in the first hour; C.S.F. protein 70 mg./100 ml.—immunoelectrophoresis normal. Serum electrophoresis showed a non-specific increase in gammaglobulins—total globulin 4.1 g./100 ml.—and immunoelectrophoresis an IgM paraproteinaemia. IgG 1,280 mg./100 ml., IgM 74% of standard, IgA 136% of standard. Sternal marrow aspirate showed a hypercellular picture with evidence of iron deficiency; staining with fluorescent anti-human IgM showed no significant uptake of the antibody. The titre of organ-specific anti-brain

antibodies was 1:1,024. A barium-meal examination was performed. The films showed an enlarged left kidney, and a large vascular tumour in the lower pole was demonstrated by aortography.

Exploration was performed by Mr. Dawson Edwards and a large hypernephroma removed. At operation blood samples were taken from both renal artery and renal vein for immunoglobulin estimations; no significant difference was found. The tumour was also stained with fluorescent anti-human IgM, but no convincing uptake of the fluorescent stain occurred.

COMMENT

Though neurological syndromes associated with renal tumours have been previously described in the literature (Swan and Wharton, 1963; Madanagopalan and Saratchandra, 1966) this association is uncommon.

IgM paraproteinaemia has been reported in association with malignancy, and in this patient it was initially considered whether the paraprotein represented an antibody response to antigens in the hypernephroma. Immunofluorescent studies with the use of labelled anti-human IgM, however, failed to show any uptake of the patient's IgM by the tumour, and the IgM paraproteinaemia was thought to be of the essential monoclonal benign type (Waldenström, 1952).

Complement-fixing organ-specific anti-brain antibodies have previously been reported in four cases of sensory neuropathy and one with a neuromyopathic syndrome, all associated with carcinoma of the bronchus (Wilkinson, 1964). It is of interest that in this patient the autoantibodies were in the IgM fraction of the immunoglobulins and that they occurred in a high titre of 1:1,024. The IgM anti-brain antibodies could also be detected by immunofluorescence to a titre of 1:160. These titres, though significant, are not high enough to enable one to conclude that the IgM paraprotein was specifically directed against brain tissue. The relationship between the patient's neurological symptoms, autoantibodies, and hypernephroma remains obscure.

I should like to thank Dr. Michael Small for his encouragement in the preparation of this paper, and Dr. R. W. Thompson and Dr. N. Williamson for their interest and helpful advice in the immunological and immunofluorescent studies.

D. C. THRUSH, M.B., M.R.C.P.
Neurological Registrar, Queen Elizabeth Hospital
Birmingham 15.

REFERENCES

- Croft, P. B., and Wilkinson, M. (1969). *Brain*, 92, 1.
- Madanagopalan, N., and Saratchandra, R. (1966). *Lancet*, 1, 1351.
- Swan, C. H. J., and Wharton, B. A. (1963). *Lancet*, 2, 383.
- Waldenström, J. (1952). *Advances in Internal Medicine*, 5, 417.
- Wilkinson, P. C. (1964). *Lancet*, 1, 1301.