37th week of pregnancy by demonstrating raised levels of 17-ketosteroids and pregnanetriol in amniotic fluid. In this condition, which may be difficult to diagnose at birth, there is a risk of sudden death in the early neonatal period and prenatal diagnosis may therefore be helpful.

Determination of Fetal Sex

Genetic diagnosis has extended into the prenatal period¹⁷ and it is now possible to determine fetal sex by finding Barr bodies or by showing fluorescence of the Y chromosome in amniotic fluid cells. This is most useful if the mother is a carrier of an X-linked gene for serious disorders such as haemophilia or Duchenne muscular dystrophy when males carry a 50% risk of having the disease.

Amniotic cell culture may be used to demonstrate abnormal chromosomes. For example, there is a high risk of Down's syndrome in the offspring of G/D translocation carriers. If the disorder is diagnosed in the fetus in early pregnancy, termination may be considered. Unfortunately, amniocentesis is not feasible before the 12th week of pregnancy and cell culture takes another four weeks, by which time pregnancy is well advanced. Nevertheless with further progress it may be possible to reduce the culture time.

Over 30 rare metabolic disorders can be recognized by biochemical study of cultured cells. The field is a rapidly expanding one and is increasingly being applied to prenatal diagnosis. Many of these disorders are inherited as recessives,

when the recurrence risk is 25%, but with these new techniques it should be possible in future to give a positive diagnosis with an accuracy approaching 100%.

References

- ¹ Royal College of Physicians of London, *Smoking and Health Now*. London, Pitman, 1971.
- ² Russell, J. G. B., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1969, **76**, 208.
- ³ Campbell, S., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1969, **76**, 603.
- ⁴ Willocks, J., Donald, I., Campbell, S., and Dunsmore, I. R., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1967, **74**, 639.
- ⁵ Brosens, I., and Gordon, H., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1966, **73**, 88.
- ⁶ Lind, T., and Billewicz, W., *British Journal of Hospital Medicine*, 1971, **5**, 681.
- ⁷ Liley, A. W., *American Journal of Obstetrics and Gynaecology*, 1961, **82**, 1359.
- ⁸ Hibbard, B. M., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1964, **71**, 529.
- ⁹ Saling, E., *Foetal and Neonatal Hypoxia in Relation to Clinical Obstetric Practice*. London, Arnold, 1968.
- ¹⁰ Fedrick, J., *Annals of Human Genetics*, 1970, **34**, 31.
- ¹¹ Torpin, R., *Fetal Malformations Caused by Amnion Rupture during Gestation*. Springfield, Illinois, Thomas, 1968.
- ¹² Pedersen, L. M., Tygstrup, I., and Pedersen, J., *Lancet*, 1964, **1**, 1124.
- ¹³ Farquhar, J. W., *Archives of Disease in Childhood*, 1969, **44**, 36.
- ¹⁴ Nelson, M. M., and Forfar, J. O., *British Medical Journal*, 1971, **1**, 523.
- ¹⁵ Stevenson, A. C., in *Ciba Foundation Symposium on Congenital Malformations*, ed. G. E. W. Wolstenholme and C. M. O'Connor, p. 241. London, Churchill, 1960.
- ¹⁶ Queenan, J. T., and Gadow, E. C., *Obstetrics and Gynecology*, 1970, **35**, 648.
- ¹⁷ Nadler, H. L., *Journal of Pediatrics*, 1969, **74**, 132.

Medical Education

Medical Television in the Inner London Area

MICHAEL CLARKE, IAN GILLILAND

British Medical Journal, 1971, **4**, 108-109

The Inner London Education Authority has established a closed circuit television network in Inner London by means of a cable laid by and rented from the Post Office. Channel 7 of this network is for higher education. The University of London has developed an Audio-Visual Centre, which, as part of the work which the centre does for the many disciplines of London University, has prepared television recordings for postgraduate medical education. These programmes are produced in co-operation with the British Postgraduate Medical Federation and with financial support from the four metropolitan regional hospital boards. The medical programmes will be transmitted on Channel 7.

The University Audio-Visual Centre's headquarters are at 11 Bedford Square in London, and it is here that programmes are planned and produced. In this building there is a small studio for television recording and equipment for film making, including animated film, photography, sound recording, and

graphic design. Most of the medical programmes are made in the centre but, where necessary, a mobile unit can go outside to record material. This mobile unit is in a large van which contains a control room with three cameras and the necessary staff. Recordings can thus be made anywhere in the metropolitan area—in clinics, laboratories, theatres, or wards. Film can also be made where extra precision and control are needed and the film subsequently included in a television recording.

The British Postgraduate Medical Federation, as part of the University of London, is responsible for medical postgraduate education in a large area of southern England which includes some 72 academic centres in district hospitals of the four metropolitan regional hospital boards. Soon the 12 undergraduate teaching hospitals, the 15 postgraduate institutes, and 11 of the major district hospitals in the Inner London area will be connected to the network, so that there is a large potential audience. In all these attached centres medical programmes can be seen by switching the receiver at the appropriate time to Channel 7.

Object of Programmes

The object of these programmes is to enable as many medical personnel as possible to see and hear expert teachers who cannot visit all these centres in person, while at the same time providing the experts with all the aids which they might use in their usual teaching—graphs, diagrams, charts, photo-

University of London

MICHAEL CLARKE, M.A., Director, University of London Audio-Visual Centre

IAN GILLILAND, M.D., F.R.C.P., Assistant Director, British Postgraduate Medical Federation

graphs, film sequences, still or moving micrography, or radiography, all of which can be included in television recordings. From the standpoint of the viewers this obviously allows them to "attend" formal lectures and engage in subsequent discussion without moving from their place of work. Most of the recordings have been made to last 20-25 minutes, as they are intended to initiate discussion.

Immediate production is confined to four series which have been designed to fit the academic year, the first taking place on 11 October. Times of transmission of programmes, which will be repeated at different times on various days, will be posted where television receivers are found. Each institute or hospital will receive advance programmes and will make its own arrangements for notification and for viewing. Viewers will be given the opportunity to comment on the programmes; indeed, feed-back will be encouraged for assessment of the value of the transmissions.

The four series in current production are:

The Scientific Basis of Medicine.—This series of public lectures by leading research workers in sciences related to medicine has an established place in British medicine. They will now take the form of weekly television discourses, each broadcast on four occasions during a week.

Clinical Medicine.—This series is particularly aimed at junior doctors. For the first two terms the topic is "Emergency Procedures" and such situations as acute renal failure, cardiac arrest, status asthmaticus, and others will be covered, so it will also be appropriate for all those dealing personally with emergency situations.

Up to Date.—This series will endeavour to keep the audience abreast of new developments, as well as reviewing fundamental concepts. A single field of medicine will be covered each term. The autumn term topic is cardiology, and endocrinology will be the subject for the spring term. In each case a wide range of speakers will be drawn from available experts in the London area and sometimes from outside as well.

Medicine This Week.—This magazine programme will deal with current topics, containing medical news and reminders of meetings. There will also be interviews with distinguished visitors and medical educationists.

In the immediate future centres connected with the I.L.E.A. cable will be at an advantage. Those with a videotape recorder can make a record of the programmes for themselves and use them again whenever they wish, particularly at times other than those advertised. Outside the Inner London area recordings will be made available through the regional hospital boards. Television receivers will be needed to view these recordings and some type of videotape recorder to replay them. The Audio-Visual Centre can make recordings for replay on Sony, Ampex, or I.V.C. recorders. A number of academic centres have already acquired small Sony $\frac{1}{2}$ in videotape recorders, and these and other makes can be used with any television receiver or monitor which has a video input device. Advice on this can be obtained from the centre.

In the fairly near future video-cassette recorders will be available which can be used in any postgraduate centre or even in private homes. This is a device rather like an ordinary tape recorder in size, into which a cassette of videotape fits; by connexion to an ordinary television set, the device will play back the material recorded. It can also be used to make a record of any particular programme on broadcast television, such as *Medicine Today* or even a Cup Final. When these machines have been evaluated and are freely available medical recordings will be transferred on to the appropriate type of cassette. The cost of such devices at the moment is just under £300, so possibly even isolated general practitioners in the countryside will be able to subscribe to a medical cassette library and find it a welcome supplement to the present journals.

Finally, the importance of audience participation cannot be overstressed. The policy committee, under the chairmanship of Professor Norman Ashton, F.R.S., whose members represent the different disciplines of medicine and career grades, can function effectively only if it is made aware of the opinion of its medical colleagues on the programmes. This committee's assessment of current medical educational needs will be influenced by demand from and comment by the viewers.

Any Questions?

We publish below a selection of questions and answers of general interest

Risk of Cataract with Prednisolone

A middle-aged woman with disabling asthma is controlled with prednisolone. She has only one eye. What is the risk of her developing cataract in the remaining eye if prednisolone is continued over long-term?

The incidence of posterior subcapsular cataract during systemic steroid therapy increases with the dose and duration of treatment. More cases of cataract have been reported in patients with rheumatoid arthritis than in patients with asthma. There is probably a one-in-ten chance of cataract formation with a dosage of 10 mg prednisolone daily over some years, but it must be emphasized that the typical subcapsular opacities induced have little effect on vision. In the patient referred to, relief of the disabling asthma clearly outweighs the risk of cataract formation.

Micropenis

Two brothers aged 10 and 2 have micropenis. They are mentally normal, have no physical stigmata, and are not obese.

There is no hypospadias. There are no associated symptoms other than the onset of problems from the older boy who realizes he is different from other boys. What investigations should be carried out, and is there any treatment?

The questioner does not mention the condition of the testicles in the boys and this is of some importance. There are some examples, apparently with a familial basis, in which the testicles appear to be normal but the penis is particularly small, although normal in its anatomy. In these cases treatment by hormones does not appear to be in any way effective and it must be presumed that there is a failure of end organ response to the normal stimulus. The only possible treatment in these cases is the construction by plastic surgery of some appendage resembling a penis into which the more adventurous surgeons may insert some silicone stiffening with the object of making sexual intercourse possible if not enjoyable.

If, however, the micropenis is accompanied by absence of palpable testicles then a much more thorough investigation is required, including chromosome analysis and possible laparotomy to determine the precise anatomy. Some of these patients will undoubtedly have much more complicated intersex type of anomalies.