mimicked by multiple abscesses communicating with the intrahepatic ducts.—I am, etc.,

JAMES G. McNULTY

Jervis Street Hospital,
Dublin 1

¹ Caroli, J., Soupault, R., Kossanowsky, J., Plocker, L., and Para Dowska, E., *Semaine des Hôpitaux de Paris*, 1958, 34, 128.

Operations for Obesity

SIR,—Your leading article entitled "Operations for Obesity" (30 October, p. 247) gave a comprehensive review of the value and hazards of small-bowel shunt surgery in the management of gross refractory obesity. May I comment on this form of treatment from my own limited experience of four patients upon whom I have performed jejuno-ileostomy within the past two years?

Three women and one man have undergone end-to-side jejuno-ileostomy of the "14+4" dimension for refractory obesity amounting to 95-135% above ideal body weight. The follow-up period has been 4-18 months. The figure shows that satisfactory reduction in weight

the laughing stock of the community. The obvious reversion to more normal clothed bodily contours has overcome this social disability, or is in the process of doing so.

With regard to the technique of the operation, I prefer to use a 6-8 in (15-20 cm) midline incision, closed with stainless steel wire, rather than the transverse "melon-slice" approach, deferring apronectomy to a later date. In this way there have been no problems of seroma, haematoma, dehiscence, incisional hernia, or postoperative respiratory difficulties. The operation is hardly formidable, though good strong retraction is necessary.

Though jejuno-ileostomy is still regarded as an experimental procedure, I believe that the "14+4 operation"¹ has now been performed with satisfaction by a sufficient number of surgeons as to justify a place in the management of gross intractable obesity. Not only is it effective and relatively uncomplicated, but it also has the advantage of being reversible. The main proviso to its use in selected patients is that they should be followed closely for several years to de-

and his colleagues agree that 20% is a more accurate measurement than 15%.⁹

There is no doubt that the ¹²⁵I-fibrinogen test is the most useful method at present available for screening large numbers of patients and detecting early, forming thrombi. It is particularly useful in post-operative patients. Nevertheless we have made it clear¹⁰ that there are defects. In the more proximal veins, in the groin and pelvis, there is not sufficient difference in the radioactivity over the veins and adjoining tissues to be confident about diagnosis of thrombosis. In addition, there are difficulties with established thrombosis where the process is static or resolving.

We have now completed an investigation into 82 patients with clinical signs suggesting established deep vein thrombosis. The results of this study are being published in detail elsewhere.¹¹ In each of the patients both the ¹²⁵I-fibrinogen test and phlebography were used. In 15% there was a false-positive finding with the radioactive test. The phlebograms established that there were, in fact, no thrombi there. Subsequently it was found that these patients suffered from such conditions as rupture of muscle fibres in the calf, haematoma within the calf, infection in the region, or other conditions. On the other hand, approximately 25% of old thrombi, proved phlebographically, were negative with the radioactive test. It is curious, and here we agree with Mr. Browse and his colleagues, that apparently static or dissolving thrombi can in a considerable proportion of cases become radioactive. Our practice, therefore, with so many false positives and negatives in suspected cases of established deep vein thrombosis is to use venography. Our experience indicates that this is the more accurate and practical way of assessing the condition at this stage and deciding on the management of established deep vein thrombosis.—I am, etc.,

J. G. MURRAY

Department of Surgery,
Kino's College Hospital Medical School,
London S.E.5

occurred in each, some stabilization developing in the one patient followed for 18 months. One patient has had one bulky stool each day, but the remainder have three to five loose unformed stools daily. This frequency has not been too much of a disability and has varied with dietary intake. No abnormalities of blood picture, haematological indices, or serum vitamin B₁₂ levels have occurred. Only in the patient followed for 20 months has the serum folate level fallen to a level demanding correction. Before operation serum iron levels were either low-normal or subnormal in all the patients, and this abnormality became more pronounced during the follow-up periods. Serum electrolytes, calcium, magnesium, and uric acid values have remained unchanged from preoperative levels. Serum albumin has fallen mildly to low-normal values in two patients. Dramatic falls in serum cholesterol of 32-50% have been seen in all patients.

So far each patient has been highly gratified with the operation and its results. All have lost their presenting symptoms such as breathlessness, bronchitis, and backache. All look healthy, and with their restored mobility two have returned to employment previously abandoned owing to their obesity. They all eat as much as they wish; a habit in which they have not indulged without guilt for some years. A further notable feature has been their social reacceptance; grossly obese patients tend to be cast out by their family or friends as well as being

test early any of the potential nutritional and metabolic abnormalities which may arise.—I am, etc.,

MICHAEL BADDELEY

General Hospital, Birmingham

¹ Payne, J. H., and DeWind, L. T., *American Journal of Surgery*, 1969, 118, 141.

Diagnosis of Established Deep Vein Thrombosis

SIR,—I was most interested to read the account of Mr. N. L. Browse and his colleagues (6 November, p. 325) on the use of the ¹²⁵I-fibrinogen test in the diagnosis of established deep vein thrombosis. This method has superseded the ¹³¹I-fibrinogen test and was developed in this department.^{1,2} Subsequently the accuracy of the method was shown by ascending functional phlebography³ and later confirmed by numerous workers.⁴⁻⁷ The original method used a scaler, but was superseded by a much simpler technique using a ratemeter.⁸ The results using a scaler and a ratemeter were compared.⁸ Well over 1,000 patients have now been investigated by us using the ¹²⁵I-fibrinogen test. It was pointed out when the ratemeter was introduced that the most accurate method in diagnosing a thrombus was a 20% difference in radioactivity. Subsequent experience has confirmed this. I am happy to see that Negus

- Atkins, P., and Hawkins, L. A., *Lancet*, 1965, 2, 1217.
- Atkins, P., and Hawkins, L. A., *British Journal of Surgery*, 1968, 55, 825.
- Kakkar, V. V., and Flanc, C., *British Journal of Surgery*, 1968, 55, 384.
- Flanc, C., Kakkar, V. V., and Clarke, M. B., *British Journal of Surgery*, 1968, 55, 742.
- Negus, D., Pinto, D. J., Le Quesne, L. P., Brown, N., and Chapman, M., *British Journal of Surgery*, 1968, 55, 835.
- Lambie, J. M., et al., *British Medical Journal*, 1970, 2, 142.
- Milne, R. M., Griffiths, J. M. T., Gunn, A. A., and Ruckley, C. V., *Lancet*, 1971, 2, 445.
- Kakkar, V. V., Nicolaides, A. N., Renney, J. T. G., Friend, J. R., and Clarke, M. B., *Lancet*, 1970, 1, 540.
- Pai, B. Y., and Negus, D., *Lancet*, 1971, 2, 1098.
- Murray, J. G., *Modern Trends in Surgery*, ed. W. T. Irvine, 3, p. 192. London, Butterworths, 1971.
- Kakkar, V. V., *Archives of Surgery*, 1971, in press.

SIR,—We were interested to read the paper of Mr. N. L. Browse and his colleagues (6 November, p. 325), but found their one-day measurement reflected the blood radioactivity in the thrombosed limb rather than the uptake of ¹²⁵I fibrinogen in a thrombus. Previous investigations have shown that serial measurements over several days will show a developing thrombosis, but that early rises lasting one day are of no significance.¹ Rosenthal and Greyson² injected 99mTC-labelled albumin macroaggregates (into the