lamines, might be fluctuating hypertensive filling of the atherosclerotic coronary vessels in a usually normotensive or hypertensive person. This may cause the trauma to the arterial wall which sets off changes leading to thrombus formation and cardiac infarction.

Treatment with β -blocking or hypotensive agents might be used to prevent unnatural rises of blood pressure, level out fluctuations, and thus avoid repeated trauma to the vulnerable vessel wall. Many potential patients are not hypertensive, and a raised blood pressure is therefore not the only reason why this approach should be tried.—I am, etc.,

ERIC FRANKEL

Wanstead Hospital,
London E.11

Incidence of Gonorrhoea

SIR,—Dr. W. H. Hughes and Mr. J. M. Davies (13 November, p. 424) suggest that in Britain and at this time there may not be a reservoir of silent female carriers of gonorrhoea of the size that has been supposed. The annual incidence of gonorrhoea is published by the Chief Medical Officer of the Department of Health using data supplied to him by the special clinics.¹ It is widely believed that these figures represent a fair sample of the cases occurring and are a high proportion of the total. In 1969 the incidence of new infections among girls of 18-19 years was almost exactly the convenient figure of 400 per 100,000 of the female population of that age and this was by far the highest incidence of any age group.

The investigation of Dr. Hughes and Mr. Davies took place between 11 March and 26 August 1971. It is the study of the prevalence of gonorrhoea among a sample of one third of the new patients at the Samaritan Hospital for Women near the time of their attendance or admission. The prevalence of previously undiscovered infection was 0.30%.

An estimate of the point prevalence in a representative sample of girls of 18-19 years would require at least a knowledge of the mean duration of an infection at this age. On this point there is no statistical evidence available. If to cover this deficiency the assumption is made that all infections discovered and recorded had been present for twelve months, then the estimated point prevalence equals the annual incidence and is 0.40%. Correction factors are required for this rough estimate. An allowance should be made for the probable short-fall in the recording of cases and the failure for a number of possible reasons to make a diagnosis. The result must be multiplied by the appropriate factor. If the assumption that each infection lasted for one year were not in accord with clinical experience, the correction factor required would almost certainly be less than one. The two corrections might therefore be self-balancing.

If the yield of screening programmes should prove to be of the order of, say, 0.5% then it would require 26,450 examinations using the 1969 figures to increase the discovery of female gonorrhoea in the special clinics by 132 cases or 1%. Many such examinations would be carried out primarily for reasons other than screening. The discovery of gonorrhoea would be a bonus to the patients

concerned but a relatively small advantage to the public health.—I am, etc.,

W. F. FELTON

St. Thomas's Hospital,
London S.E.1

¹ Department of Health and Social Security. *On the State of the Public Health. Annual Report of the Chief Medical Officer.* London, H.M.S.O. 1970.

Need for Asylums

SIR,—Dr. O. W. S. Fitzgerald (27 November, p. 556) writes of the considerable number of so-called criminals occupying our prisons who are in fact chronic schizophrenics or subnormals, and the difficulty he has in obtaining appropriate care for them when they are discharged from prison.

Having, like Dr. Fitzgerald, moved from a psychiatric hospital into the prison medical service (from which I recently retired), I can endorse his remarks. It was my experience that it was just as difficult to obtain psychiatric treatment or suitable institutional care for these people before committal to prison so that the courts had no choice about disposal.

The mentally ill and the subnormal are a burden to the penal establishments. Efforts are being made to provide suitable care, but there are many difficulties, overcrowding and shortage of suitably trained staff being the biggest. It seems surprising that in our welfare society provision is no longer made for the chronic untreatable mental case whose antisocial behaviour makes him too hot to hold in our "open door" psychiatric hospitals. Such people need long-term care and a prison sentence does not necessarily provide this. Frequently trivial minor offences are committed for which the offenders receive short sentences, and on coming out the whole cycle starts again. One wonders what the cost is in time spent by the police, courts, probation officers, and welfare workers.

It is to be hoped that with the reconsideration of the mental health services an answer will be found, and there will be better liaison between the various agencies involved in the care of these people.—I am, etc.,

FREDA S. REED

Chilham, Kent

Unfair Pressure for Abortion

SIR,—What is the position of a general practitioner when he disagrees with his patient's request for an abortion? She may decline his offer of social and antenatal support and press for abortion, usually on social grounds. Nor are gynaecologists immune from unfair pressure, and there may be more to come. The Medical Practitioners' Union is reported¹ as having recommended to the Lane committee that the 1967 Abortion Act be amended so that "doctors who refuse to perform abortions should be made to give their reasons in writing." It would seem unlikely that unfair pressure of this type would commend itself to a future Parliament, though such pressure does exist on the other side of the Iron Curtain, where the rule for gynaecologists and their trainees is "abort or get out."

The legal position under the 1967 Abortion Act is quite clear. A practitioner should examine his patient and offer his con-

sidered opinion on what he thinks is in her best interest. He may or may not advise a termination, but even in the face of rudeness from the patient, her husband, or her mother there is no legal or moral obligation to offer more than a medical opinion given in good faith. Naturally a Roman Catholic or other doctor with personal reasons making it difficult for him to take part in an abortion consultation should explain the position to his patient, arranging for her to consult a suitable medical colleague, perhaps his partner. Naturally he could not introduce his patient to an advisory agency known to arrange abortion on demand in private. The press and radio must take much of the blame for giving the public the false impression that Parliament in 1967 made it legal for any woman to request and to get an abortion either in the N.H.S. or privately.

Each of us must take up an ethical position, which need not lack compassion for the mother or her baby. I believe that good medical practice is based on kindness, a careful examination, and careful notes. This need not lead to bullying a patient. But, vice versa, a patient should not try to bully or threaten her medical adviser nor should she cross his palm with silver. There are nevertheless occasions when inevitably a patient and her doctor must disagree, but this should always be possible without rancour, and especially since the patient seeking the termination is often distressed, even distraught.—I am, etc.,

HUGH CAMERON McLAREN

Birmingham Maternity Hospital,
Queen Elizabeth Medical Centre,
Edgbaston,
Birmingham 15

¹ *Daily Telegraph*, 24 November 1971.

Australia Antigen in Nigerian Blood Donors and School Children

SIR,—The high prevalence of Australia antigen (Au(1), H.A.A.) in the "normal" population has been reported to be high in tropical areas.^{1,2} There have also been many reports of the prevalence of Au(1) in blood donors from several countries. There have, however, been no reports of the prevalence of Au(1) in blood donors in Nigeria. This information is supplied here. Eight thousand sera obtained from volunteer blood donors, 95% of whom are men aged between 18-60 years at the University College Hospital, Ibadan, Nigeria, between 1 February and 30 September 1971 were screened by the electroimmuno-osmotic diffusion method of Bedarida *et al.*³ The specificity of the detected Au(1) and its antibody was confirmed by double diffusion using standard reagents

Monthly Prevalence of Au(1) in Nigerian Blood Donors (February-September 1971)

Month	Total No. Donors	Au(1) Positive		Au(1) Antibody Positive	
		No.	%	No.	%
February	1,001	50	4.9	1	0.1
March	1,044	56	5.6	2	0.19
April	1,121	120	10.1	5	0.45
May	773	20	3.8	1	0.13
June	1,149	47	4.1	—	—
July	771	29	3.9	1	0.13
August	998	87	8.7	2	0.2
September	1,143	65	5.7	1	0.09