respiratory exchange ratio may be used to determine whether the ratio is outside the normal range (usually owing to an unsteady respiratory state), and in such cases it would be preferable to use equation (i) rather than to rely on the nomogram for measurement of oxygen consumption. When the respiratory exchange ratio was within the limits 0.70-0.93 the maximum error was 10 ml/min (S.D. 3.92).

Discussion

The respiratory exchange ratio can be derived from the nomogram with an error which is limited by visual discrimination on the nomogram. With normal eyesight it should be possible to be accurate within 0.01, which is sufficient for many purposes. It is certainly adequate for determining whether the value is

sufficiently far from the normal range to invalidate the derivation of oxygen consumption on the other half of the nomogram.

Oxygen consumption may be derived from the volume and oxygen concentration of expired air with an error of standard deviation 7.73 ml/min, the error compounding variations in respiratory exchange ratio, barometric pressure, and laboratory temperature. When patients with abnormal values for respiratory exchange ratio are excluded the error is substantially less, and acceptable for a wide range of clinical purposes. Both nomograms apply only to patients breathing room air.

Reference

Pappenheimer, J. R., *et al.* *Federation Proceedings*, 9, 602.

Nomograms for Calculation of Heat Loss

FRANK HOUSE, RAYMOND VALE

British Medical Journal, 1972, 4, 20-21

Summary

Two nomograms are presented. The first enables the mean surface and body temperatures and the body heat content of a patient of given weight to be determined from measurements of skin temperature at three sites and of the core (rectal) temperature. The second enables the change in heat content of such a patient to be determined from the change in mean body temperature.

Introduction

Heat production and loss from the body can be measured directly by calorimetry and indirectly by measuring the temperature at various sites on the body surface. The first method is too difficult for clinical use, but the latter is possible and useful in many circumstances. Changes in rectal temperature alone do not adequately explain discrepancies between heat production and heat loss. Heat debt calculated from the mean surface temperature "mixed" with rectal temperature was found by Burton and Edholm (1955) to give the best agreement with observed differences between heat production and heat loss. They found that the best value for the mixing coefficient was:

$$\theta = (0.7 \times T_{\text{core}}) + (0.3 \times T_s)$$

where θ represents the mean body temperature, T_{core} the core (rectal) temperature, and T_s the mean surface temperature.

The mean surface temperature is derived as follows:

$$T_s = (0.14 \times T_{\text{forearm}}) + (0.36 \times T_{\text{calf}}) + (0.5 \times T_{\text{abdomen}})$$

these temperatures being measured on the skin surface at these sites.

If heat balance is the point where heat production equals heat loss then:

$$\Delta \text{ heat balance} = \Delta \theta \times 0.83 \times \text{wt.}$$

where Δ heat balance = change in heat balance, $\Delta \theta$ = change in mean body temperature, 0.83 = specific heat of tissues, and wt. = body weight (kg).

Calculating these values arithmetically is laborious, and for use in clinical practice two nomograms have been devised.

Nomogram 1

A straight edge is laid across scale A and scale C (Fig. 1) at the measured temperatures of the skin of the calf and the skin of the forearm. The point of intersection on scale B is joined to the abdominal skin temperature on scale D and the mean surface temperature is read on scale C. The mean surface temperature on scale C is then joined to the core temperature on scale E and the point of intersection of this line on scale D gives the mean body temperature. If this point is joined to the body weight on scales F1, F2, or F3 the heat content can be read on the corresponding scale (G1, G2, or G3).

Nomogram 2 (Fig. 2)

Often it is only changes in heat content that are of interest. In such cases it may prove more convenient to use nomogram 1 to obtain mean body temperature at various times and then nomogram 2 to convert changes in temperature to changes in heat content. If the body weight on the first scale of nomogram 2 is joined to the change in mean body temperature on the third scale, the change in heat content can be read from the point of intersection on the second scale. For small body weights the two scales on the right of the nomogram are used.

Example of Use

The three sets of temperatures in the Table can be used to illustrate the use of the nomograms.

The mean surface temperature derived from the preinduction temperatures by means of nomogram 1 is 31.8°C and the mean body temperature is 35.0°C. Repeating the procedure with the postinduction temperatures (measured 15 minutes after the induction of anaesthesia) gives a mean surface temperature of

Temperatures (°C) at Various Sites Before and During Operation

	T _{core}	T _{forearm}	T _{abdomen}	T _{calf}
Preinduction	36.4	32.4	31.8	31.6
Postinduction	35.7	31.9	34.6	32.8
End of Operation	35.8	31.2	32.3	31.9

Discussion

Many workers have recorded changes in core temperature during anaesthesia and surgery, but few have measured changes in both core and surface temperatures. As will be seen from the example given, if meaningful conclusions are to be drawn, both must be measured. However, the significance of changes in

FIG. 1—Nomogram for calculation of heat content.

FIG. 2—Nomogram for calculation of change in heat content.

33.6°C and a mean body temperature of 35°C. Thus as a result of the vasodilatation produced by anaesthesia the core temperature has fallen by 0.7°C and there has been a redistribution of heat within the body without loss of heat.

The figures for the end of operation give a mean surface temperature of 32°C and a mean body temperature of 34.7°C. With the fall in mean body temperature of 0.3°C nomogram 2 shows a reduction in heat content of 17 kcal for a 70-kg man.

surface temperature at several sites is difficult to appreciate, and it is hoped that the present nomograms will enable these changes to be related to changes in heat content and thus increase the practical usefulness of temperature measurement.

Reference.

Burton, A. C., and Edholm, O. G. (1955). *Man in a Cold Environment*. London, Arnold.