CORRESPONDENCE

Correspondents are asked to be brief

Immunological Control of Schistosomiasis G. S. Nelson, M.D. 172	Survey of Alimentary Radiological Findings J. C. Petrie, M.R.C.P., and others 176	Acute Dystonia due to Metoclopramide P. S. Venkateswaran, M.R.C.P.E.D., and A. G. Otto, M.B. 178
Vocational Training in General Practice E. O. Evans, F.R.C.G.P. 172	Dialysis and Transplantation in Britain C. Shaldon, F.R.C.S., and others 176	Skin Exfoliation after Renal Failure C. Giordano, M.D., and others 179
Toxoplasmosis D. G. Fleck, M.D. 173	Tetracyclines for Malaria T. Wilson, M.D. 176	Abdominal Distension R. Heller, M.D. 179
How Infectious is Gonorrhoea? W. K. E. Bernfeld, M.D. 173	Abortion Deaths Josephine A. C. Weatherall, M.B.; Madeleine Simms, M.A. 176	Was it a Drug? T. K. Meikleham, M.B. 179
Puzzling Cases of Secondary Syphilis P. W. M. Copeman, M.D. 173	Rh Isoimmunization from Ruptured Tubal Pregnancy G. de B. Mitford-Barberton, M.R.C.O.G. 177	New Consultant Contract C. E. Astley, F.R.C.P.; J. S. McCann, M.D. 179
Treatment of Early Breast Cancer Sir Hedley Atkins, F.R.C.S., and others 173	Payment by Colour "Medical Graduate"; N. Pover, D.P.H. 177	Regional Hospital Staffing S. C. Simmons, F.R.C.O.G.; G. I. B. da Costa, F.R.C.S.E.D.; F. J. Bramble, F.R.C.S.; R. D. G. Creery, F.R.C.P. 180
Gas Gangrene and Hyperbaric Oxygen J. B. Bourke, F.R.C.S., and H. D. Ritchie, F.R.C.S.; D. J. D. Perrins, F.R.C.S.; A. J. Keogh, F.R.C.S. 174	Uganda Asians F. C. Harris, F.R.C.PATH. 178	Consultants and Registrars R. Strang, F.R.C.S. 181
Induction of Labour D. R. Firth, B.M. 175	Problems with Ketamine Anaesthesia Lieutenant-Colonel C. K. Davies, F.F.A.R.C.S. 178	Medical Teachers' Salary Awards M. D. Buckley-Sharp, M.B. 181
Diagnosis of Chronic Beryllium Disease W. J. Williams, F.R.C.PATH., and others 175	Leucocytosis after Tetracosactrin Lieutenant-Colonel A. R. Robinson, M.R.C.P.E.D. 178	Future of the B.M.A. R. D. Abernethy, M.B. 181
Fungal Contamination of Cervical Sponges A. D. Bremner, D.C.H., and Milicent H. S. Bell, M.B. 175	Reversal of Adult Mental Subnormality M. W. P. Carney, M.D., and others 178	Christmas Gifts Appeal Sir Thomas Holmes Sellors, F.R.C.S. 181

Immunological Control of Schistosomiasis

Sir,—Drs. R. B. Stott and H. J. N. Bethell (7 October, p. 49) refer to your leading article on immunological control of schistosomiasis (12 August, p. 366) and suggest that the elimination of this disease "is possible if doctors all over the world have the breadth of vision, courage, and determination to take medicine to the people and allow them to effect their own cures rather than awaiting miracles from the boffins."

As one who tried this approach for 13 years in East Africa and failed I would like to defend the boffins who are scorned by your correspondents for their "restrictive thinking" and "unjustified pessimism." Unjustified, or so they claim, because the doctors of the People's Republic of China have already demonstrated that all you have to do is educate everyone on the "life style of the snail and nature of the disease" and with the necessary enthusiasm of the people you will get rid of the problem. I suspect that Drs. Stott and Bethell have been reading reports that came out of China before the Cultural Revolution in 1966.

In a careful review of the Chinese literature from 1949 to 1970 Tien-Hsi Cheng,¹ who is enthusiastic in his praise for the achievements of China, notes that a great deal of the Chinese literature before 1966 is "premature, politically orientated and symbolic of propaganda for domestic and foreign consumption." He also reports that in the three years after 1966 the control programme was in disarray and "the disease was raging again in some localities where previously it had been brought under control." Even before 1966 there are very few

convincing accounts of anything more than limited local control of schistosomiasis in China. One of these accounts which is frequently quoted is by Mao and Huang,² who claimed that the snail hosts had been eradicated from 300,000 acres of farmland. This was a considerable achievement but it involved a minute part of the endemic area of China, and it was very much smaller than the Gezira scheme in the Sudan, where "boffins" from the London School of Hygiene and Tropical Medicine are collaborating with the Sudanese authorities in an attempt to eliminate the disease from two million acres of irrigated land.

Drs. Stott and Bethell are apparently unaware that the problem of controlling the *Biomphalaria* and *Bulinus* aquatic hosts of *Schistosoma mansoni* and *S. haematobium* is incomparably more difficult than getting rid of the amphibian *Oncomelania* snail hosts of *S. japonicum*. Indeed many of the techniques used in China would make the problem in Africa much worse. I have always been an optimist and I can see prospects for the successful control of snails and the disease in irrigated areas, but the most highly educated and highly motivated community is unlikely to have any impact on the snails in the vast swamps and rivers and lakes of Africa (even with the help of your correspondents' "snail spotters").

There are no prospects of totally eliminating the snail hosts of schistosomiasis from any of the main endemic areas and mass treatment without snail control is usually ineffective, especially in countries like China where more than 31 species of wild and

domestic animals are known to be reservoir hosts. That is why the boffins should be encouraged to pursue their studies in the hope of finding a more effective means of immunological control.—I am, etc.,

G. S. NELSON

London School of Hygiene and Tropical Medicine,
London W.C.1

- 1 Tien-Hsi Cheng, *American Journal of Tropical Medicine and Hygiene*, 1971, 20, 26.
- 2 Mao, S. P., and Huang, M. S., *Kexue Tonbao (Scientia)*, 1964, 11, 963.

Vocational Training in General Practice

Sir,—It was stated at the Annual Representative Meeting (*Supplement*, 29 July, p. 80) and in your leading article (30 September, p. 781) that there is a wide gap between the places available for vocational training and the number of doctors entering general practice each year. No doubt there is never enough training of the highest quality. However, the existing supply of training facilities still exceeds current demand, and there is evidence that even the potential gap is not as great as has been suggested.

Thus the average age of becoming a principal in general practice is over 29 years. Clearly, therefore, the main fault is the relevance of the training rather than the time spent over it. Some of the courses especially arranged for vocational training in general practice are still undersubscribed. The number of teaching practices likewise exceeds the supply of willing trainees. In Warwickshire, for example, there are several