CORRESPONDENCE

Correspondents are asked to be brief

Radiography of Potentially Pregnant Females G. M. Ardran, F.R.C.P., and F. H. Kemp, F.R.C.P.	422	I.U.D. and Uterine Bleeding N. K. Bhide, and S. T. Gogte.	426	Self-poisoning with Drugs C. L. Tolhurst-Cleaver, M.B.	429
Myeloid Leukaemia and Cot Deaths Alice M. Stewart, F.R.C.P.	423	Canavan's Disease and Spongiform Encephalopathy H. B. Pratt, M.B.	427	Self-poisoning with Paracetamol P. Barnes, B.M., and J. S. Prichard, D.Phil., B.M.	429
Erasures from the "Register" P. Noone, M.R.C.PATH., and others.	423	Inappropriate Secretion of ADH J. Sack, M.D., and M. A. Sperling, M.D.	427	Poisoning Treatments Centre I. Oswald, M.D.	430
Abortion Deaths Sir George Godber, F.R.C.P.	424	Australia Antigen and Pulmonary Tuberculosis J. M. Martinez-V., and others.	427	Accident Flying Squad D. Lamont, F.R.C.S.	430
Geriatric Accommodation in Acute Illness J. A. Fraiss, M.R.C.G.P.	424	Gentamycin in Urinary Infections J. H. Darrell, M.R.C.PATH.	427	Conscientious Objection to Abortion Margaret B. Wilkinson, D.C.H.	430
Drugs in Infertility G. I. M. Swyer, F.R.C.P.	425	Call the Doctor W. G. Davidson, M.D.	427	Influenza and Infant Mortality A. Czeizel, M.D.	430
Confinement of Subnormal Offenders D. A. A. Primrose, M.D.	425	Serum and Urine F.R.-Antigen in Renal Disease E. N. Wardle, M.R.C.P.	427	Acute Bed Requirements A. I. Ross, M.D.	430
Consultants in Mental Handicap D. A. Spencer, D.P.M.	425	Effect of Ultrasound P. Flamme, M.D., and C. Lardinois.	428	Raynaud's Disease W. Silber, F.R.C.S.ED.	430
Active Management of Labour M. K. O'Driscoll, F.R.C.O.G.	425	Phoenix from Physical Medicine J. H. Cyriax, M.D.	428	Insertion of Celestin Tubes B. S. Mantell, F.F.R.	431
Graphic Records in Labour J. W. W. Studd, M.D., and N. M. Duignan, M.R.C.O.G.	426	Electrotherapy C. B. Heald, F.R.C.P.	428	How Infectious is Gonorrhoea? W. F. Felton, D.P.H.	431
Vasectomy in the Surgery J. J. Hobbs, M.B.	426	Lumbar Disc Problems R. K. Jackson, F.R.C.S.	429	M.A.O.I. Treatment Cards A. J. Boakes, F.F.A.R.C.S.	431
Criteria for Free School Milk J. C. Waterlow, F.R.C.P.	426	Nurses for Nursing I. R. Gray, F.R.C.P.; T. C. Dann, M.D.	429	Careers in Community Medicine J. L. Gilloran, M.R.C.P.ED., and others.	431
				New Consultant Contract J. W. Paulley, F.R.C.P.	432

Radiography of Potentially Pregnant Females

SIR,—In June 1972 the Department of Health and Social Security published a revised *Code of Practice for the Protection of Persons against Ionizing Radiations arising from Medical and Dental Use* (London, H.M.S.O.). In the preface (para. 3) it is stated that "although the arrangements recommended relate primarily to institutions they should be applied, as far as practicable, by all medical and dental practitioners." The accompanying circular HM(72)39 from the Department imposes upon boards of governors and regional hospital boards the duty of implementing the code. The bulk of the code applies to those who administer the radiation, whether for diagnosis or treatment, but Section 7, dealing with the protection of the patient, applies also to those who request that such procedures be carried out. In particular we are concerned with paragraph 7.3.1. which reads as follows: "In all women of reproductive capacity the clinician requesting the examination should consider the possibility of an early stage of pregnancy. The date of the last menstrual period should be entered on the request form and it is the responsibility of the clinician requesting the examination to ascertain this. To reduce the likelihood of irradiation of a pregnancy, examinations involving the lower abdomen should, if practicable, be carried out within 10 days following the first day of the menstrual period."

Radiological Protection Committees are considering how these problems can best be dealt with to ensure that no embryo or fetus receives more radiation than is absolutely essential. The code and the circular do not

give specific details as to how the problem might be adequately tackled. As far as we are aware the bulk of medical practitioners have not been contacted to inform them of these requirements of the code and to suggest how the problem may best be dealt with.

Following the work of Dr. Alice Stewart and others it is now universally accepted that radiation to the fetus can cause leukaemia or other neoplastic disease. The hazards may be estimated in simple terms with sufficient accuracy for the present purpose in the following way:

The natural incidence of childhood neoplasia and leukaemia is about 1 in every 1,200 live births. The best estimates indicate that a dose of 1.4 r will double the natural incidence. Diagnostic examination of the pelvic regions can result in a dose to the fetus ranging from about 0.25 r up to perhaps 5 r or more in an extensive examination. This means that an average examination might double the natural incidence of leukaemia etc. This could also be expressed by saying that should leukaemia or neoplasia arise in a child who was radiographed as a fetus there is a fifty-fifty chance that the radiation caused it. All the evidence indicates that an early pregnancy is just as susceptible as a later one.

The code of practice clearly places the responsibility for finding out whether the patient might be pregnant or not on the referring clinician and requests that the date of commencement of the last menstrual period should be entered on the request form. Experience has shown that it is almost impossible for the x-ray department to cope with this problem without further assistance.

Many individuals do not know when their next period will occur and booking clerks cannot be expected to make these inquiries and arrange to book the patient within the first 10 days of the beginning of the next period (when the patient is unlikely to be pregnant) without considerable delays and difficulties. It is therefore suggested that the referring clinician, when requesting an x-ray or isotope investigation involving the area from the diaphragm to the knees, should ascertain if there is any possibility of pregnancy. He should also state whether the examination is of such urgency that it should be carried out regardless of whether the patient is pregnant or not. If the examination is not of immediate urgency or of a type for which an appointment must be booked, it is suggested that the referring physician should explain to the patient that because of a small hazard to the fetus it is best if the examination is carried out when the patient is not pregnant and the patient therefore should take the necessary steps to see that she does not become pregnant until after the examination.

To avoid difficulties when the patient arrives at hospital either for an outpatient appointment or with a view to admission it is further suggested that in any case in which there is a possibility of a diagnostic radiological investigation being required the patient likewise should be advised not to become pregnant until after the possibility of a radiological examination has passed, and it should be explained to her that if she fails to do this it will be her responsibility if any unfortunate consequences should occur.

We suggest that doctors should be supplied with a card (see Proforma) on which

they should certify that they have done this. They should obtain a signature from the patient acknowledging that this has been explained to her and has been understood. This card should be filed with the x-ray or the diagnostic isotope request card. It should be pointed out that when there is a possibility of a hysteroscopy examination this should not be carried out within the first 7 days after the onset of a period and that it is essential that the patient should not become pregnant before the examination has been performed.

Proforma

STRICTLY CONFIDENTIAL

X-RAY DIAGNOSIS

RECORD FORM FOR FEMALE PATIENTS BETWEEN PUBERTY AND THE MENOPAUSE To be completed by patient

Full Name
Date of birth
Date

The Code of Practice for the Protection of Persons against Ionizing Radiation (1972) requires that the clinician requesting the examination should consider the possibility of an early pregnancy.

Will you please answer the following questions:

- (1) Is there any possibility that you might be pregnant?
(Delete the word which is not applicable) YES OR NO
- (2) Please enter date of the 1st day of the last menstrual period
- (3) I understand that it is important that I should avoid possible pregnancy until the x-ray examination booked for me has been carried out. I undertake to inform the staff of the x-ray department if there is any chance that I might have become pregnant.

Signature of patient

To be completed by the doctor:

Please indicate which of the alternative procedures you wish to be adopted.

- (1) My patient M/s has been asked to undergo an immediate x-ray examination. The questions relating to pregnancy have been discussed.
- OR
- (2) My patient M/s is referred for a booked appointment. I have explained that she should not run a risk of pregnancy until the examination has been performed.

Signature of doctor

This form should be completed when the clinician requests an x-ray examination of any part between the diaphragm and the knees or a radioisotope investigation.

It is our view that the only way to cope with this problem is to place the onus on the patient to inform the doctor if there is any possibility of pregnancy and to take the necessary steps to see that she is not pregnant before being radiologically examined. It is clearly laid down in the code that it is the responsibility of the referring doctor to deal with this aspect. The x-ray department can then only do its best when this has been carried out. It is appreciated that this scheme might solve only about 95% of the problem, but this is probably the best that can be done. It is also appreciated that there may be difficulty with girls under the age of 16, but this should not deter us from attempting to deal with the main problem. It is suggested that a copy of Section 7 of the code should be sent to all medical practitioners with a covering letter, which might be based on the above.

Without the collaboration of referring physicians the problem, in our opinion, cannot be adequately solved and will inevitably result in delays and inefficiencies in any radiological service. We appreciate that whatever method is adopted would involve already

overworked people in more work and we would invite suggestions as to how the problem may be adequately dealt with more conveniently or expeditiously. Of course it must be stressed that any pregnant women who, for good medical reasons, need to be radiographed should be radiographed carefully and the small hazard accepted as inevitable, as other hazards are accepted daily when necessary.—We are, etc.,

G. M. ARDRAN
F. H. KEMP

Radcliffe Infirmary, Oxford

Myeloid Leukaemia and Cot Deaths

SIR,—In view of the following observations it has occurred to me that myeloid leukaemia might be a numerically important cause of cot deaths.

(1) Acute leukaemias are notoriously difficult to recognize postmortem, and the earlier the death the greater the difficulty.

(2) Between the initiation of leukaemia and the onset of symptoms (incubation period) there is a prodigious increase in the risk of dying from respiratory infections.¹

(3) Before the second world war the number of acute lymphatic leukaemias which were mistaken for fatal pneumonias was far greater than the number of recognized cases.²

(4) The postwar increase in leukaemia mortality has not affected infants or older children with myeloid leukaemia, and it was much slower to affect chronic lymphatic leukaemia—which is rarely diagnosed before 60 years of age—than other variants.

(5) During the first decade after the war there was a dramatic decrease in the infection death rate, and during the second decade there was an equally dramatic decrease in the number of deaths ascribed to senility,³ but there has been only a modest decrease in the number of stillbirths and neonatal deaths since 1945.

(6) The vast majority of cancers which prove fatal before 10 years of age are initiated before the third trimester of fetal life.⁴

(7) The number of identical twins who die from cancers within 10 years of birth and have not been exposed to obstetric radiography is much smaller than the expected number. On the other hand, the number of twin-concordant cancers in this age range is far larger than the expected number.⁵

(8) Myelocytes have short cell generation times and are produced by a tissue which is relatively mature at birth. By contrast, circulating lymphocytes have long cell generation times and are produced by a tissue which is relatively immature at birth.⁶

(9) Intervals between diagnosis and death are shorter for acute myeloid than acute lymphatic leukaemias, but the difference is much smaller after the age of 10 years than before the age of 3 years.⁵

(10) Most of the acute lymphatic leukaemias which are caused by obstetric radiography are diagnosed between 3 and 6 years of age, but the corresponding set of myeloid leukaemias are concentrated between 7 and 10 years of age.⁷

To an epidemiologist these miscellaneous observations suggest the following possibilities.

Early cancers have either germ cell or embryonic origins and when they cause blood dyscrasias instead of tumours they increase the risk of dying in utero, during parturition, and during childhood illnesses.

Cancers have distinctive intervals between initiation and diagnosis which are related both to the nature and to the maturity status of the original "malignant" cell. Consequently myeloid leukaemia is a more acute disease than lymphatic leukaemia; embryomas are more acute diseases than adult

cancers; and myeloid embryomas disappear from the scene earlier than lymphatic embryomas.

Myeloid embryomas are more often mistaken for other causes of death than lymphatic embryomas because the short incubation periods make them potent causes of stillbirths and infant deaths. They are also more often missed than other embryomas with equally short incubation periods because postmortem diagnosis of a blood dyscrasia poses more problems than postmortem diagnosis of a localized tumour.

If there were no difficulty in recognizing leukaemias after death we might find that myeloid embryomas produced a peak of mortality at birth—due to children who experienced a difficult delivery—and a second peak three to four months later due to children who had an easy birth followed by a cot death.⁸

If it were equally easy to recognize embryomas with short and long latent periods we might find that the early peak of leukaemia mortality at three years⁹ was as much an artefact as the late peak at 65 years. Nowadays the late peak is found only in societies which are supposed to be immune to chronic lymphatic leukaemia but are clearly living in conditions which make it easy to mistake this disease for the ill-defined signs and symptoms of senility.^{3,10}

On the assumption that we are recognizing only a quarter of the leukaemia deaths before 5 years, a haematology survey might find that the missing cases were responsible for 10% of unexplained stillbirths and a much higher proportion of cot deaths.—I am, etc.,

ALICE STEWART

Department of Social Medicine,
Oxford University

¹ Kneale, G. W., *British Journal of Preventive and Social Medicine*, 1971, 25, 152.

² Stewart, A. M., and Kneale, G. W., *Nature*, 1969, 223, 741.

³ Stewart, A. M., in *Clinics of Haematology*, ed. S. Roach, No. 1, p. 3. London, Philadelphia, Saunders, 1972.

⁴ Stewart, A. M., and Kneale, G. W., *Lancet*, 1970, 2, 4.

⁵ Data from Oxford Survey. Awaiting publication.

⁶ Hamilton, W. J., and Mossman, H. W., *Human Embryology*, 4th edn. Cambridge, Heffer, 1972.

⁷ Stewart, A. M., *Proceedings of the Royal Society of Medicine*, 1972, 65, 245.

⁸ Camps, F. E., and Carpenter, R. G., *Sudden and Unexpected Death in Infancy (Cot Deaths)*. Bristol, John Wright, 1972.

⁹ Hewitt, D., *British Journal of Preventive and Social Medicine*, 1955, 9, 81.

¹⁰ Segi, M., Kurihara, M., and Matsuyama, T., *Cancer Mortality for Selected Sites in 24 Countries*, (Nos. 1-5). Sendai, Japan, 1960-69.

Erasures from the "Register"

SIR,—We, the undersigned medical staff of the Middlesex Hospital Medical School, feel compelled to inform the General Medical Council of our profound disquiet at recent developments over the retention fee and the threat of imminent erasure hanging over many doctors who have withheld payment of the fee on grounds of principle until a public inquiry is held into the functions, structure, and financing of the General Medical Council.

There can be no doubt that the overwhelming majority of the profession would welcome such an inquiry, and Lord Cohen himself has declared publicly that he is not opposed to it. We urge the G.M.C. and the B.M.A. to take steps to ensure that this inquiry is held as soon as possible.

We must also totally deplore the threats of