for granulomata in serial sections none were found. Further, we are not aware of sub-total villous atrophy occurring in those parts of the jejunum otherwise uninvolved in small-intestinal Crohn's disease.—We are, etc.,

P. E. JONES

Medical Unit,
University College Hospital Medical School,
London W.C.1

M. H. GLEESON

Royal Infirmary,
Manchester

¹ Goulston, K. J., Skyring, A. P., and McGovern, V. J., *Australasian Annals of Medicine*, 1965, 14, 57.

² Bayless, T. M., Kapelowitz, R. F., Shelley, W. M., Ballinger, W. F., and Hendrix, T., *New England Journal of Medicine*, 1967, 276, 996.

³ Pink, I. J., and Creamer, B., *Lancet*, 1967, 1, 300.

⁴ Wall, A. J., Douglas, A. P., Booth, C. C., and Pearse, A. G. E., *Gut*, 1970, 11, 7.

Seat Belts

SIR,—As the proposer of the successful motion at the Annual Representative Meeting at Folkestone in June on the need for improvement in the design of seat belts and the advisability of their compulsory use in cars (*Supplement*, 16 June, p. 94), I was gratified to see the letter from Mr. E. S. R. Hughes of Melbourne (29 September, p. 695) on the experience in Australia of the lowered mortality and reduced incidence of injuries resulting from their legislation.

How right Professor W. Gisane (30 June, p. 772) was in commending the action of the Australian legislature in introducing compulsory belt-wear! Let us hope that our own legislators will be as discerning and perspicacious.—I am, etc.,

R. COVE-SMITH

London W.9

Hypotension in Tetanus

SIR,—I was interested to read the article by Dr. J. L. Corbett and others (25 August, p. 423), and would like to make two suggestions:

(1) Rather than leave the patients with an alarmingly low blood pressure for 15-20 minutes, to try immediate infusion of 500-1,000 ml of plasma or other fluid. A heart pumping at a control venous pressure of only 1.5 mm Hg as described in case 1 can be expected to respond to increased filling pressure with an increase in cardiac output and blood pressure, because of Starling's law of the heart. Increasing the blood volume is more effective than head-down tilt, and is surely safer than giving alpha-, beta-, or ganglion-blocking drugs to patients with hypotension. Patients on mechanical ventilation are likely to need a greater than normal blood volume, especially if autonomic function is impaired.

(2) Why not keep the patients sedated as well as paralysed? Reflex depression with barbiturates, diazepam, droperidol, or narcotics may well prevent these episodes of hypertension, which are quite common during light general anaesthesia. Perhaps normal subjects, without tetanus, would react in a similar way to prolonged curarization and mechanical ventilation as these three patients.—I am, etc.,

JAMES JUDSON

Middlemore Hospital,
Auckland, New Zealand

Hypergastrinaemia in Rheumatoid Arthritis

SIR,—We read with interest Dr. K. J. Fraser's reply (29 September, p. 696) to our preliminary communication (30 June, p. 752) concerning elevated concentrations of plasma gastrin in rheumatoid arthritis. Dr. Fraser's initial comments on our lack of awareness of the literature on the incidence of achlorhydria in rheumatoid arthritis are perhaps unfortunate as no reference is made to the work of Armstrong,¹ who demonstrated no increase in incidence of parietal cell autoantibody in rheumatoid arthritis. He also demonstrated the organ-specificity of the antibody by means of gastric function studies. The omission of reference to this work carried out in this centre is out of step with Dr. Fraser's kindness elsewhere in referring to our work in the field of Sjögren's syndrome. We would like to point out in addition that the increased incidence of hypergastrinaemia may prove to be of much more fundamental importance if it is not due to gastric atrophy.

Our preliminary studies would suggest normal acid secretory status in rheumatoid arthritis in the presence of hypergastrinaemia, and recently have shown that the induction of adjuvant arthritis in rats is associated with a concomitant elevation of fasting plasma immunoreactive gastrin.² The significance of these observations remains unclear to date, but it seems unlikely that they merely reflect the presence of atrophic gastritis.—We are, etc.,

PATRICK J. ROONEY

WILLIAM CARSON DICK

Centre for Rheumatic Diseases,
Glasgow

¹ Armstrong, E. M., *Annals of the Rheumatic Diseases*, 1967, 26, 499.

² Rooney, P. J., Dick, W. C., Buchanan, K. D., and Imrie, R. C., *Nature New Biology*. In press.

Making Hospital Geriatrics Work

SIR,—Under this heading Drs. H. M. Hodgkinson and P. M. Jefferys (2 December, 1972, p. 536) discussed the work of a new geriatric department and advocated a policy of high turnover and no waiting list. They suggested that this policy was a matter of choice by the geriatricians.

Experience at Greenwich over recent years indicated that the work of a geriatric department is not quite such a matter of choice as Drs. Hodgkinson and Jefferys state. The figure shows the changes that have occurred over a 10-year period. From 1963 to 1968 the service at Greenwich was very similar to that at the Northwick Park unit with no waiting list, direct admission on general practitioner or Emergency Bed Service request, and high turnover (five patients per bed yearly), and the wait for transfer from other wards and hospitals was less than a month. At that time the work of the department was in equilibrium with the needs of the area served.

In 1968 22 beds were closed out of a total of 200 (11%) and the admission rate has fallen from 1,000 per year to 450—a drop of 55%. This disproportionate fall stems from a simple error. When the beds were closed it was argued that an 11% reduction in beds would act as a stimulus to the efficiency of an over-bedded department (200 beds for 18,000 over-65 population). What

was overlooked was that the department had taken on a high long-stay commitment to help out other agencies under stress. There were 150 long-stay patients, including a 33-bed psychogeriatric ward with demented patients who in many parts of the country would have been placed in a psychiatric hospital, and 26 patients had been accepted from welfare accommodation for whom Greenwich would not normally have been the responsible geriatric hospital. There were therefore only 50 admission and rehabilitation beds through which 1,000 admissions were passing each year. Within these 50 beds were about 25 very-high-turnover beds and about 25 slower-stream beds with a stay of three months or more before discharge, death, or transfer to long-stay accommodation. Clearly none of the long-stay or slower-stream beds could be closed and the department was left with virtually no high turnover beds.

My choice in this situation was to accept preferentially those patients whose stay was likely to be short and to give low priority to those referrals who were likely to need a prolonged stay. Nevertheless, the department has functioned over the last five years as a low-turnover/long-waiting-list unit. In spite of not now taking ambulant demented patients and in spite of now adopting a "legalistic" approach to the geographical catchment area the number of long-stay patients is now 141 (the total bed number having fallen this year to 171). There are signs, however, that a new equilibrium is being established since the number of patients in other hospital beds waiting for transfer to the department has fallen from a maximum of 70 in 1972 to 20 in September 1973. During the whole period the staffing (medical, nursing, and paramedical) has remained substantially the same in numbers but, as would be expected, morale has plummeted.

The suggestion by Drs. Hodgkinson and Jefferys that the work of a geriatric department is a matter of choice therefore appears to be an over-simplification. Given positive attitudes and adequate staffing, there appears to be a critical number of beds in any area with which a high-turnover/no-waiting-list service can be given. An important lesson that comes out of the Greenwich experience is that the total number of beds in a geriatric department is a misleading figure. In planning changes the number of admission and rehabilitation beds is a key