is being used, so as to ensure that the recording is in fact of the fetal heart rate alone.—We are, etc.,

P. TIVY-JONES
A. KENNEY
R. D. DE VERE

Obstetric Department,
Westminster Hospital,
London S.W.1

Confidentiality

SIR,—We read about the confidentiality of the computer, but I feel we need to look at things much more human before we start on mechanization.

I have this morning received a letter from the appointments clerk to a hospital psychiatric outpatient department in which she states that a patient has an appointment to see a psychiatrist and that a note of any treatment she was having would be beneficial. The enclosed envelope is addressed to "The Psychiatric Department." I do not know what my protection society would have to say about this.

During the past month I was asked to send to the "organizing secretary" of the local mass radiography unit details of the diagnosis and subsequent treatment of a patient who had had an abnormal miniature radiograph. In this case my protection society was very doubtful that I should disclose this information to a layman.—I am, etc.,

C. LIPP

Sheffield

Methyldopa and Depression

SIR,—I read with interest the report by Mr. C. J. Bulpitt and Professor C. T. Dollery (27 October, p. 232) on side effects of hypotensive drugs and the subsequent correspondence which it aroused on the existence of methyldopa depression as a specific entity.

For the past three years I have been measuring mood changes in a group of new hypertensive referrals to the Sheffield Hypertension Clinic, using a standardized mood-rating questionnaire¹ which has been shown to be both valid and reliable. Patients completed this questionnaire at their first clinic attendance and at six-weekly intervals for a one-year follow-up period and the scores were recorded graphically.

For purposes of statistical evaluation, the patients were divided into two groups: (1) those taking adrenergic blocking agents, and (2) those taking reserpine or methyldopa. There was no significant difference in the prevalence of depression occurring in the two groups, but there was a statistically significant difference in the pattern of depression. The depressed patients on the adrenergic-blocking drugs reported themselves as mildly chronically depressed throughout the period of study. Patients who became depressed while taking reserpine or methyldopa showed a transient depressive swing within the first few months of starting the drugs. It is postulated that the two patterns of depression have differing aetiologies. Patients taking adrenergic-blocking drugs are likely to be more severely hypertensive and suffer more medical and social complications of their illness than the less severely hypertensive. This pattern of depression can be regarded as an "illness effect"

rather than a direct effect of the drugs. On the other hand, the transient depression of the hypertensive on methyldopa or reserpine is in keeping with the known amine-depleting effect of these drugs. Other factors shown to be of significance in the aetiology of both patterns of depression were personality structure and evidence of previous psychiatric disorder.

I would like to suggest that the amine-depletion pattern of depression still exists, but to a lesser degree than a few years ago, owing to a greater awareness of the problem, but there is an increasingly large group of more severe hypertensives suffering chronic depression, often unrecognized, resulting from the effects of a chronic illness on a particularly vulnerable personality. Further research on this subject is in progress.—I am, etc.,

WENDY BANT

Whiteley Wood Clinic,
Sheffield

¹ Abraham, H. C., Kanter, V. B., Rosen, I., and Standen, J. L., *British Journal of Psychiatry*, 1963, 109, 286.

Explaining to Patients

SIR,—Cancer, writes Professor W. A. Mackey (3 November, p. 297), is a "frightening, primitive word. . . . It is a pity that we do not have readily available a less horrifying name." When discussing the effect of words upon patients¹ I suggested the term growth. It is less frightening and allows discussion of different types of growth and their prognosis, starting with benign ones.

What to tell the patient is a big problem in many branches of medicine. So many diagnoses carry a halo of fear with them. Angina is associated with the thought of sudden death, a point that will be emphasized to the victim by many kind friends. Arthritis is linked with the worry of being crippled; mental or nervous symptoms with the fear of going mad. The word thrombosis, even if referring to a thrombosed pile, may strike terror, being misinterpreted as implying the same seriousness as coronary thrombosis. Perhaps someone should compile a glossary of diagnostic labels which may help to shield the patient from unnecessary and often unfounded anxiety. My contributions would be: heart attack instead of coronary thrombosis, Parkinsonianism and not paralysis agitans, rheumatism or "fibrositis" instead of arthritis (unless the doctor has time to explain the various and often harmless implications of the word), seizure rather than stroke, and growth or tumour instead of cancer.—I am, etc.,

CLIFFORD HAWKINS

Queen Elizabeth Hospital,
Birmingham

¹ Hawkins, C. F., *Speaking and Writing in Medicine: The Art of Communication*. Illinois, Thomas, 1970.

Medical Education in Italy

SIR,—Professor Alberto Zanchetti's account of Italian undergraduate and postgraduate medical training (17 November, p. 409) may appear "excessively gloomy" but is, in the Italian context, very optimistic. From experience gained and the information available to me during three months spent in Italy as a student last winter it appears that certain

facts and the translation of some Italian terms in the account are very misleading.

The official government estimate for the number of medical students for the year 1972-3 was 120,000, a third of whom were in the first year studying "A-level" subjects, and this represents a tripling of medical student numbers in four years. I agree with Professor Zanchetti's statement that ward round teaching is impossible, but would go further to say that for the vast majority of students it is non-existent. He refers to a six-month "internship." However, a more precise translation is "apprenticeship." The "apprentice," who may be a student yet to take university finals, is unpaid and is required only to be present at, but not involved in, routine hospital procedures. Even the postgraduate training does not put the "doctor" in the position of a houseman.

Worthwhile reform is unlikely since the last governmental reforms were too late and, from a British point of view, retrogressive. Therefore in arriving at future agreements and reviewing the peculiar existing arrangement the Italian medical training system should be much more thoroughly scrutinized.—I am, etc.,

MARIO BROWN

King's College Hospital Medical School,
Denmark Hill,
London S.E.5

Medically Oriented Language Courses

SIR,—Professor J. D. E. Knox asks for suggestions for medically oriented language courses (8 September, p. 546).

It is already the policy of the Junior Members Forum that such language courses should be provided, and a motion concerning this subject was carried by the meeting held at Coventry earlier this year.

From a personal aspect, as a medical officer in the armed Forces, I would find such courses, especially in German, extremely useful. Owing to the pressure on the majority of doctors' time, these courses should be available through postal means. There should be two courses for each language—one for the beginner and one for the doctor who already has a working knowledge of the particular language. These could be available on tape cassettes with a short explanatory booklet accompanying each cassette.—I am, etc.,

M. J. G. THOMAS

Chairman,
Junior Members Forum, B.M.A.

Colchester

Diabetic Pregnancy

SIR,—Thank you for the excellent survey of diabetic pregnancy by Dr. N. L. Essex and others (13 October, p. 89). It seems to me, however, that there was one important omission—namely, the extreme danger of continuing insulin in unchanged dose during and directly after delivery. I realize that this is well known to the initiated, but I take it the article was aimed largely at the uninitiated.

For reasons not known to me, for one, two, or even three days after delivery a normally insulin-requiring woman may need no insulin at all and may easily be precipitated into hypoglycaemia. Once one has

seen a young mother decerebrated by inappropriate insulin continuation one can never forget it.

May I make a second point concerning medical management? In my part of the world we can very frequently manage overweight pregnant diabetics with on 800-kcal regimen only, with considerable success, even when initial blood glucose levels have been around 300 mg/100 ml.—I am, etc.,

W. P. U. JACKSON

Department of Medicine,
University of Cape Town

Son et Lumière?

SIR,—At a time when more and more patients attending outpatient clinics appear to have functional illnesses with no physical signs it is refreshing to meet a condition presenting in a rather dramatic manner.

The patient concerned had had a vagotomy and pyloroplasty a year previously and complained of increasing vomiting. On examination there was obvious splashing in the stomach, suggesting a diagnosis of pyloric stenosis. He was asked if he experienced foul eructations and replied that he certainly did. Then, rather diffidently, he said, "On one occasion I was lighting a cigarette when I belched and the explosion burnt my nose and hair."

This sign of pyloric stenosis after vagotomy is, I feel, on a par with the "flash in the pan" of phosphorus poisoning, but I have been unable to think of a suitable name for the condition. Perhaps your readers could help.—I am, etc.,

H. A. DANIELS

Ulverston, Lancs

Independence of Review Body

SIR,—I refer to your leading article "Phase 3 and Beyond" (17 November, p. 374) and particularly to the questions you raise.

It is quite right that the profession should question the independence of the Review Body. The Review Body is supposed to make an independent review and arrive at a fair conclusion on the profession's increase in remuneration on the merits of the case presented. It is obviously unable to fulfil this remit by virtue of the fact that it is expected by the Government to seek the approval of the Pay Board for its recommendations. This is very clearly illustrated by events of the recent past and now even more openly by Sir Keith's suggestions in his letter to the Chairman of the B.M.A. Council (*Supplement*, 17 November, p. 45). It is inconceivable how one can then continue to think of an independent Review Body.

It is indeed right that the profession's rise in remuneration should take into account the prevailing national economic problems. It is also indeed right that acrimonious situations where professions and Government take issue with one another over what is a fair increase should be avoided. However, a truly independent arbitrating body would arrive at what one would justifiably expect to be a just and equitable conclusion, and so, even though the Government may subsequently reject the award and pare it down on the grounds of

of what is a fair award acrimonious situations could be avoidable. It may be that in the national interest the profession would feel morally bound to accept an award lower, much lower, than would be recommended by an independent Review Body, however unpleasant that might be. In that case the Government would be giving public recognition and the public at large would be aware that the profession's just award was being sacrificed. It is reasonable to conclude that subsequently the profession would have a stronger case in their favour and also expect and demand an amendment when better economic times prevail.

There is therefore only one course of action in reply to the questions you raise. The profession should insist on an award uninfluenced by the Pay Code and that Lord Halsbury and his colleagues make their recommendations on the merits of the case of the profession. Opting for the alternative implies acceptance of the fact that the Review Body is just an implement in the hands of the Government to keep the profession's salary increase to a predetermined level. Calling the Review Body independent under such circumstances is just a charade.—I am, etc.,

V. MANSUETO

Gillingham, Kent

Contracts for Hospital Junior Staff

SIR,—Dr. Ian Grant's letter (24 November, p. 494) demonstrates the confused attitude which arises in those who have not read carefully the reasoning behind the new contract proposals for hospital junior staff and who have therefore missed its basic philosophy of two separable loyalties. The offer of nothing more than skill and time is to the employer, not the patient.

Continuous commitment to the patient already under his care remains the doctor's ethical responsibility; far from finding this irksome or unrewarding, we trust the idealism and the conscience of the British junior doctor sufficiently to feel that there is no need to write such moral responsibilities into his contract, thus laying him open to exploitation by employers. No suggestion has been made that we should actually work only a 40-hour week or that this would provide adequate training, but the standard working week (at present up to 80 hours) which we propose, *plus* "voluntary overtime" which we assume everyone would wish to do as clinical needs may occasionally require, is a basis on which we hope to negotiate the adequate salary which Dr. Grant says he wants for the long hours which we recognize we shall continue to work.

If we should receive many letters expressing such views as Dr. Grant's, or hear these opinions through our constitutional machinery, we should, of course, hold a referendum, and would hope that we might depend on most juniors to take the trouble to read the background to the questions asked. Fortunately, however, most of the mail reaching the Hospital Junior Staffs Group Council says in some form: "Carry on the good work, and the best of luck!"—which also answers Dr. W. P. Bradford's letter (24 November, p. 494). Hospital junior staff meetings are held and publicized (through national interest, with an *a priori* knowledge the B.M.A.) in every region. Attendances

are variable, but the Group Council can hardly be blamed if juniors will not take an interest; and we hope that if they do *not* approve of our policies the juniors in the regions will come and debate them. All have chosen to accept a share of the responsibility for the running of the profession—a responsibility which cannot be undertaken entirely by non-professionals without precisely that loss of professional freedom which Dr. Grant fears.—I am, etc.,

JEAN TURNER

Chairman,

Hospital Junior Staff Group Council, B.M.A.
B.M.A. House, London W.C.1

Buying Added Years

SIR,—I wish to express complete agreement with the points raised by Dr. A. P. Grant in his letter (10 November, p. 355). I should like to ask all senior hospital medical staff over the age of 55 who are in agreement with Dr. Grant's letter to write to the Chairman of the B.M.A.'s Superannuation Committee asking him to take up the points mentioned as a matter of urgency. Very little has been achieved in negotiating a just superannuation for the older group of senior hospital staff.

I should like to mention one additional point. In the early days of the N.H.S. (1948-51) part-time appointments in junior grades involving less than a certain number of sessions (I think less than nine) were not superannuable, though the first booklet on superannuation did not mention that. In view of the shortage of appointments available at the time some of us had to accept this type of work and as a result lost a few "years."—I am, etc.,

A. MARES

Rush Green Hospital, Romford, Essex

* * An interview with the Chairman of the B.M.A.'s Superannuation Committee appears in this week's *Supplement* at p. 57.—Ed., B.M.J.

Points from Letters

Monkeys for Research

Dr. J. D. WHITTALL (Rye, Sussex) writes: I was impressed with the intelligent and humane ideas formulated by Dr. R. Schenkel in his letter entitled "Monkeys for Research" (17 November, p. 421). He says that the kind of philosophy adopted by some scientists is only concerned with the benefit to the human community. That is true. The animal—the experimental tool—is not given any consideration. That we need "a more sensitive conscience with regard to experiments with highly organized living beings" is made very evident when we look at some of the cruel and futile . . . experiments to which many primates are subjected.

Injecting Hydroceles—An Unproved Treatment?

Mr. H. G. STURZAKER (South Croydon) writes: I have no experience of injecting hydroceles or epididymal cysts, but I am appalled that Mr. C. E. Moloney (20 October, p. 170) should still use a trocar. . . . A no. 1 needle is an extremely satisfactory way of aspirating a hydrocele, and though it may take slightly longer to draw off the fluid, the whole process can be accomplished more quickly than with a trocar and causes less trauma, both physical and psychological, to the patient. . . .