Reorganization of Health Services and the N.H.S. Number

SIR,—At a time of reorganization of the National Health Services it would seem expedient to examine the benefits of restructuring the N.H.S. number. That number provides unique and permanent identification of all individuals in the country, and increasing use is being made of it in medical record linkage.¹ However, it is exceptionally cumbersome to use because of its length and the variability of its format.² Individuals seldom know their own numbers, and the space for the N.H.S. number on many medical documents is frequently left blank.¹ For example, in cancer registration in South Wales fewer than 4% of N.H.S. numbers are entered by hospitals at time of registration, and some of the entries are of their national insurance number.

One possible number that would be easier to remember and hence easier to use is suggested. It is a nine-digit alphanumeric in which the first two digits denote the year of birth and the second two are letters indicating the (new) health district of birth. Then follows a four-digit sequence allocated in turn to live births in the district in the year, and the number is terminated by a check digit.³ For example, a child born in Cardiff in early 1973 might be allocated the number 73CF00798, the 8 being the check digit. Identification of sex could also be incorporated by allocating only even numbers to boys or alternatively by allocating the first 4,999 numbers to boys.

The use of fixed-length format would overcome the difficulties of remembering and transcribing which are experienced with the present series of numbers. Inclusion of two alphabetic characters in fixed positions may aid comprehension and memorability of the whole number provided that the letters have easily identifiable significance. The proposed health districts have populations of approximately 200,000 to 500,000⁴ and 676 two-letter combinations would allow description of the majority of districts by recognizable abbreviations (for example, CF for Cardiff). Code letters could be used also for immigrants indicating country of birth, and spare codes would be kept for "unplaced" births and for new or growing districts. The number of a person living to over 100 years would need to be flagged by the issuing authority to prevent reissue of the same number. National numbering systems similar to the one outlined have been introduced recently in Denmark, Norway, and Sweden.⁵ In the Danish 10-digit number the leading six digits record the date of birth in day, month, and year but the district of birth is not identified.

In April 1974 the health services in England and Wales are to start afresh, in a re-organized form. Is this not a good opportunity to introduce a more easily useable N.H.S. number based on a simpler, fixed-length format? The same number could be adopted for national insurance purposes, and each person could be issued with a plastic card bearing this new number. If it was possible to attach unique numbers to all persons in the country at short notice in 1939, it should also be possible to renumber those already registered.—I am, etc.,

ROBERT R. WEST

Department of Social and Occupational Medicine,
Welsh National School of Medicine,
Cardiff

- ¹ Acheson, E. D., *Medical Record Linkage*. London, Oxford University Press, 1967.
- ² Farmer, R. D. T., and Cross, K. W., *British Journal of Preventive and Social Medicine*, 1973, 27, 53.
- ³ Post Office Report No. 170, *The Function and Mathematics of Check Digits*, 1962.
- ⁴ *National Health Service Reorganization, England*, Cm. 5055, London, H.M.S.O., 1972.
- ⁵ Nielsen, H., in *Record Linkage in Medicine*, ed. E. D. Acheson. Edinburgh, Livingstone, 1968.

- ⁶ Donald, K. W., *Lancet*, 1949, 2, 1056.
- ⁷ Flenley, D. C., Cooke, N. J., King, A. J., Leitch, A. G., and Brash, H. M., *Bulletin de Physiopathologie Respiratoire*, 1973, 9, 689.
- ⁸ Campbell, E. J. M., *British Medical Journal*, 1965, 1, 1451.
- ⁹ Hume, R., *British Journal of Haematology*, 1968, 15, 131.

Oxygen Therapy in Chronic Respiratory Failure

SIR,—We were interested to read the paper by Dr. R. Hume and others (20 October, p. 154). However, we feel that their postulate concerning tolerance of the respiratory centre to hypoxia in these patients needs to be challenged.

Firstly, the idea that a change in the "central oxygen state" regulates respiration is difficult to accept, as the only described effect of hypoxia on the respiratory centre is one of depression.¹ The hypoxic drive is mediated by the peripheral chemoreceptors.² Further, there was no mention of the altered response to CO₂ in these patients, which is probably the most important factor determining "oxygen tolerance." In this connexion, we should like to draw attention to work from this department.^{3,4} This demonstrated that the mean ventilatory responses to CO₂ was greatly reduced in patients with chronic respiratory failure. Add to this that these patients have a greatly diminished ventilatory capacity⁵ and it becomes clear, as was pointed out by Donald,⁶ that these patients are dependent on an intact hypoxic drive to maintain ventilation. With regard to the individual variation in the tolerance to oxygen of these patients, this was shown to exist and was measured in terms of the hypoxic potentiation of the CO₂ response.^{3,4} However, the variation was no greater than in normals,³ a fact confirmed in a much larger series more recently.⁷ This is an interesting finding and may have practical implications, as discussed,⁷ but suggests very strongly that the individual variation in the patients with cor pulmonale can have little to do with the "level of arterial oxygen to which [the] respiratory centre has become accustomed during the quiescent phase of the disease."

Secondly, the authors seem to use Campbell's⁸ criteria to define hypercapnic chronic respiratory failure. If this is so, then it appears that 15 of the 30 patients studied failed to meet these criteria as their PaO₂, as shown in the authors' figure, was greater than 60 mm Hg. Unfortunately, the PaCO₂ values for all this group are not quoted but of the 15 which are given elsewhere⁹ only six were greater than 49 mm Hg.

What we also note is that 24% oxygen given empirically would have been safe and appropriate for all the patients reported who can be seen to have had hypercapnic respiratory failure.—We are, etc.,

L. J. CLANCY
A. G. LEITCH

Department of Medicine,
Royal Infirmary,
Edinburgh

- ¹ Kronenberg, R., et al., *Respiration Physiology*, 1972, 16, 109.
- ² Heymans, J. F., and Heymans, C., *Archives Internationales de Pharmacodynamie*, 1926, 32, 9.
- ³ Flenley, D. C., and Millar, J. S., *Clinical Science*, 1967, 33, 319.
- ⁴ Flenley, D. C., Franklin, D. H., and Millar, J. S., *Clinical Science*, 1970, 38, 503.
- ⁵ Flenley, D. C., and Millar, J. S., *Clinical Science*, 1968, 34, 385.

Cancer and the Patient

SIR,—When the diagnosis of cancer is made the problem arises whether or not the patient should be told (Mr. Charles Wright, 6 October, p. 45) and much of the recent correspondence on this subject in your columns has concerned the fear the diagnosis engenders in the patient. Two elements of this fear are the fear of death itself and the fear of a pain-racked, drug-doped terminal phase (Professor G. W. Milton, 27 October, p. 221). This latter is possibly the worse fear both from the patient's personal point of view and from the great, sometimes intolerable, strain the terminal period will place on the patient's immediate family.

In all the discussion no mention has been made of the place of induced hyperthermia in the management of the problem. The most striking effect of induced hyperthermia is the relief of the pain due to tumour invasion, which is relieved or abolished by one or two episodes of hyperthermia, and this is unrelated to the heat sensitivity of the particular malignancy (pain due to bony compression of nerve roots is unaffected). This pain relief often lasts for several weeks without further hyperthermia. A safe and reliable method of raising the body temperature is now available¹ and a number of patients have been treated. After treatment the patient is alert and has a sense of well-being unrelated to the pain relief. This is in contrast to some of the after-effects of radiotherapy and anti-mitotic agents. While a single treatment with induced hyperthermia undoubtedly takes longer and requires more skilled attention than an injection of morphine, the patient after hyperthermia is able to continue a normal life, meeting people and getting out and about. This is not only of benefit to the patient, but the family also suffers less. If this therapy was more widely known and available some of the fear of cancer might be removed.—I am, etc.,

E. L. LLOYD

Department of Anaesthetics,
Royal Infirmary,
Edinburgh

- ¹ Henderson, M. A., and Pettigrew, R. T. P., *Lancet*, 1971, 1, 275.

Dermographia and Tattooing

SIR,—We have been studying the incidence of dermatographia in allergic and non-allergic subjects using the procedure described by Grolnick.¹ In the course of this study an interesting observation was that on stroking a tattooed area of skin of a subject without dermatographia, while no wealing occurred over the non-tattooed areas of the skin the tattooed skin showed wealing. In subjects with dermatographia the wealing reaction was greater over the tattoo marks than over the non-tattooed skin. We would be interested to know if this has been observed by other workers interested in this subject.