FIG. 1.

later plane S-T segment depression was still evident (fig. 1c) despite the presence of normal myocardial lactate extraction (+1.3 mg/100 ml) at this time. It could be considered that increased coronary arterial perfusion resulted from the trinitrin therapy and produced a wash-through effect, thereby preventing proper evaluation of myocardial lactate extraction status. However, after a further five-minute period spontaneous angina recurred, with further plane S-T segment depression (fig. 1d) and zero myocardial lactate extraction. Five minutes after a second trinitrin tablet had been given symptoms were relieved completely and this was associated with normal myocardial lactate extraction (+1.1 mg/100 ml) despite the persistence of plane S-T segment depression (fig. 1e). It was only after a further 20 minutes had elapsed, during which the patient remained asymptomatic with normal myocardial lactate extraction (+1.2 mg/100 ml), that the S-T segment regained its previously accepted normal J-point position (fig. 1f).—I am, etc.,

BRIAN LIVESLEY

King's College Hospital,
London S.E.5

- Gianelly, R. E., Treister, B. L., and Harrison, D. C., *American Journal of Cardiology*, 1969, 24, 161.
- Livesley, B., et al., *British Medical Journal*, 1973, 1, 375.
- Livesley, B., and Oram, S., *Lancet*, 1973, 1, 1461.

Skin Reactions to Practolol

SIR,—As Drs. R. H. Felix and F. A. Ive point out in their letter (11 May, p. 333), the psoriasiform rash which appears to be associated with practolol is quite characteristic and easily recognizable. It may be of interest to note that the rash will sometimes be superimposed on psoriasis itself, which is such a common condition. In such cases it is easy to be tardy in diagnosing the drug rash since the picture suggests merely a worsening of the psoriasis and may easily be taken as a spontaneous occurrence.

I have recently seen two patients, both middle-aged women, who had been taking practolol in standard doses for a matter of a few weeks when the psoriasis became much worse; there were no special features in one case but in the other the worsening was acute and she developed generalized fiery erythema with annular peeling and involvement of the palms, which had not previously been affected. In both the exacerbation cleared rapidly without drastic measures within a week or two of the time when the drug was stopped. Both patients

had previously had extensive but stable psoriasis and there could be little doubt that the episodes were precipitated by the practolol.—I am, etc.,

C. M. RIDLEY

Department of Dermatology,
Elizabeth Garrett Anderson Hospital,
London N.W.1

E.C.T. and Cardiac Arrhythmia

SIR,—Dr. S. M. Cannicott (31 August, p. 579) recommends routine β -adrenergic blockade for electric convulsion therapy. "Sympathetic stimulation" is invoked as a possible cause; I submit that parasympathetic overactivity, evident during a convulsion, may be an alternative cause of arrhythmia, despite premedication with atropine.

Dr. Cannicott also finds standard anxiolytics contraindicated on account of their anticonvulsant properties. Many patients are already on tranquilizers and antidepressants, and anaesthesia is frequently induced with a barbiturate, with no obvious inhibitory effect on the induced convulsion.

Dr. Cannicott may have demonstrated reduced pulse rates in his patients but he has yet to show a reduced mortality rate; β -adrenergic blockade is not without its own dangers, and since many patients submitted for E.C.T. are already on multiple-drug regimens, there may be further hazards from interactions.

I suggest that there is no case for unnecessarily complicating this very safe treatment until such time as controlled trials clearly demonstrate an improvement in the mortality rate.—I am, etc.,

P. J. WARD

Flamstead, Herts

Malaria Transmission and Fetal Growth

SIR,—The article by Drs. J. D. Macgregor and J. G. Avery (17 August, p. 433) prompts me to make the following comments on malaria and fetal growth.

When pregnant women living in endemic malarial areas are unprotected by malarial chemotherapy maternal *Plasmodium falciparum* infection causes maternal anaemia,¹ placental parasitization,² and fetal growth retardation,³ all of which are more marked during first pregnancies. So it is hardly surprising that malarial eradication in the British Solomon Islands conferred a greater benefit on the babies of primigravidae than on those of multigravidae as judged by the changes in mean birth weights.

Turning to the issue of fetal growth retardation in malaria, one factor—namely, placental parasitization—has hitherto been put forward to explain it. But there is evidence which strongly suggests that maternal anaemia can also account for the prevalence of low-birth-weight babies in endemic malarial areas. In a study in Nigeria⁴ pregnant women who were initially anaemic from a variety of causes (haemoglobinopathies, malarial infection, folate deficiency) were treated with antimalarials, folic acid, and iron tablets. Blood transfusion was given in some cases. After more than six weeks' treatment the anaemia was successfully corrected in one group of

women, and in these cases the maternal packed cell volume (P.C.V.) was more than 30% at the end of their pregnancies. In the rest the P.C.V. was less than 30% when the babies were born. When other factors known to reduce fetal birth weight were excluded it was found that the incidence of small-for-dates babies differed significantly in the two groups. It was 35% in the non-anaemic group compared with 53% in the anaemic group, and in the latter a good correlation was found between maternal haematocrit and fetal birth weight. The relationship was such that a 2% drop in maternal haematocrit reduced fetal birth weight by about 100 g.

The point that maternal anaemia is associated with fetal growth retardation has therefore been made, but it requires confirmation. The malaria eradication programme referred to by Drs. Macgregor and Avery provides an excellent opportunity to do this. For this reason it would be of interest to know what the maternal haemoglobin (or P.C.V.) levels were at the end of pregnancy before and after malaria eradication.—I am, etc.,

K. A. HARRISON

Department of Obstetrics and Gynaecology,
Ahmadu Bello University Hospital,
Zaria,
Nigeria

- Gilles, H. M., et al., *Annals of Tropical Medicine and Parasitology*, 1969, 63, 245.
- Edington, G. M., *British Medical Journal*, 1967, 1, 715.
- Bruce-Chwatt, L. J., *Annals of Tropical Medicine and Parasitology*, 1952, 46, 173.
- Harrison, K. A., and Ibeziako, P. A., *Journal of Obstetrics and Gynaecology of British Commonwealth*, 1973, 80, 798.

Schistosomiasis and Irrigation

SIR,—One of the third world's principal health problems and one which seems to present the medical profession working in affected countries with well-nigh insoluble problems is that posed by schistosomiasis.

In Africa and the Americas and in parts of Asia the most important source of dissemination of schistosomiasis results from the increasing use of irrigation and is therefore man-made. The intermediate snail host will not, as far as I am aware, breed in pipes and I have been surprised to note that the possibilities of carrying irrigation water in this way has received very little comment from the medical profession in affected countries. Such a system uses only a fraction of the land occupied by an open ditch system and therefore more land can be cultivated; it saves water in areas where there may be a scarcity of this commodity and enables water to be carried to the exact place where it is needed in the exact amount and for the exact time required for a particular crop. Loss by evaporation or permeation is avoided, man-water contact is reduced practically to zero, and the risk of infection correspondingly reduced.

While installation costs are certainly higher than for open ditch irrigation, this is in part offset by savings on cultivatable land, water used, routine control measures, and medical services, as well as by increased individual productivity. When an irrigation system is installed it is intended to be a project lasting for some centuries, and it would be right and proper, therefore, to spend more on the capital outlay in order

to provide permanent protection for the population living in the area.—I am, etc.,

HUGH L. B. RUSSELL

University Department of Community Medicine,
Usher Institute,
Edinburgh

Alpha-fetoprotein and Rh Alloimmunization

SIR,—An increase in alpha-fetoprotein (AFP) has been reported in amniotic fluid from fetuses with fetal distress, including fetal distress of Rh alloimmune origin.^{1,2}

We have assayed by radioimmunoassay³ specimens of amniotic fluid from a number of Rh alloimmune pregnant women and report the results of the assay of: (1) 72 specimens from 46 women with 27-42-week-old normal or subnormal fetuses and (2) 27 specimens from 12 women with 27-38-week-old fetuses with severe haemolytic anaemia which, 3-14 days after amniocentesis, either died or required intrauterine blood transfusion or were born with levels of haemoglobin below 10 g/100 ml. Results of AFP determinations are shown in the figure.

Amniotic AFP determination at various gestational ages in cases of severe Rh alloimmunization (X) and normal controls and/or mild cases (●). Means represented by horizontal bars (—).

There was an overall tendency to higher levels in the groups of fetuses with severe alloimmune anaemia, at least until the 36th week of gestation before which most of our specimens were taken.

From a practical point of view, we think that because of wide individual variations and consequent overlapping with controls, AFP determinations on amniotic fluid from Rh alloimmune pregnant women provides less information than bilirubin determination.

We also think that recent observations by Whyley *et al.*⁴ of significant AFP increase in the amniotic fluid of pregnant women whose fetuses did not survive may be explained by the AFP rise which takes place shortly before fetal death.—We are, etc.,

A. JOUVENCEAUX

Blood Transfusion Centre,
Hôpital E. Herriot,

P. SIZARET

N. MARTEL

International Agency for Research on Cancer,
Lyons

- 1 Seppala, M., and Ruoslahti, E., *Obstetrics and Gynecology*, 1973, **42**, 701.
- 2 Guibaud, S., *et al.*, *Lancet*, 1973, **1**, 1261.
- 3 Sizaret, P., *Journal of Immunological Methods*. In press.
- 4 Whyley, G. A., Ward, H., and Hardy, N. R., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1974, **81**, 459.

Renal Amyloidosis in Chronic Granulomatous Disease

SIR,—The association of chronic granulomatous disease and amyloidosis has not been previously documented. We report the development of renal amyloidosis in a boy with long-standing chronic granulomatous disease.

The patient had had recurrent bacterial infections, particularly with *Staphylococcus aureus* and *Salmonella typhimurium*, from the age of 8 months. The diagnosis of chronic granulomatous disease, the subject of a previous report,¹ was made at the age of 10 years on the basis of a nitroblue-tetrazolium test and cutaneous granulomas.

He presented at the age of 16 years with a pure nephrotic syndrome (oedema, serum albumin 2 g/100 ml, serum cholesterol 260 mg/100 ml, proteinuria 2 g/24 hr) without microscopic haematuria or hypertension. In contrast to his previous hypergammaglobulinaemia, low levels of serum immunoglobulins (IgG 146 mg/100 ml) with elevated urinary levels (IgG 450 mg/l.) were observed. Long-standing *S. typhimurium* septicaemia responded to rifampicin. Oedema was controlled by diuretics and a low-sodium diet. Renal biopsy revealed renal amyloidosis, predominantly glomerular but also involving interstitial, peritubular, and vascular structures.

Recent studies indicate that amyloidosis is largely composed of fragments of kappa light chains.² In patients with chronic suppuration amyloidosis has been attributed to prolonged antigenic stimulation producing excessive immunoglobulin synthesis with deposition of immunoglobulin fragments as amyloid material.³

Continuing antigenic stimulation is the most likely mechanism of amyloid production in this patient. Immune function is normal in chronic granulomatous disease⁴ and Gram-negative endotoxin has been shown experimentally to stimulate amyloid formation.⁵ Alternatively, abnormal immunoglobulin catabolism may be involved, as in hypogammaglobulinaemia or agammaglobulinaemia,⁶ or dysfunction of the reticuloendothelial system.⁷

In vitro amyloid can be phagocytosed by leucocytes⁸ and the reticuloendothelial system⁹ and this is probably the mechanism of the regression of amyloid deposits reported in liver and kidney.¹⁰ Defective catabolism of immunoglobulins and amyloid substance might also occur in chronic granulomatous disease, in which phagocytic degradative activity is known to be defective, in both leucocytes¹¹ and the reticuloendothelial system,¹² particularly in respect of bacterial phagocytosis.—We are, etc.,

R. DE SEIGNEUX
A. KANFER
P. TERRIOUX
J. D. SRAER
J. A. WHITWORTH

Service de Nephrologie,
Hôpital Tenon,
Paris

- 1 Grunberg, J., *et al.*, *Annals de Pédiatrie*, 1970, **17**, 560.
- 2 Glenner, G. G., Ein, D., and Terry, W. D., *American Journal of Medicine*, 1972, **52**, 141.
- 3 Hardt, F., and Ranlov, P., *Israel Journal of Medical Sciences*, 1973, **9**, 857.
- 4 Bridges, R. A., Berendes, H., and Good, R. A., *American Journal of Diseases of Children*, 1959, **97**, 387.

- 5 Barth, W. F., Gordon, J. K., and Willerson, J. T., *Science*, 1968, **162**, 694.
- 6 Mawas, C., Sors, C., and Bernier, J. J., *American Journal of Medicine*, 1969, **46**, 624.
- 7 Nordstoga, K., *Acta Pathologica et Microbiologica Scandinavica*, 1972, **80**, 159.
- 8 Zucker-Franklin, D., *Journal of Ultra-structural Research*, 1970, **32**, 247.
- 9 Shirahama, T., Cohen, A. S., and Rodgers, O. G., *Experimental and Molecular Pathology*, 1971, **14**, 110.
- 10 Lowenstein, J., and Gallo, G., *New England Journal of Medicine*, 1970, **282**, 128.
- 11 Holmes, B., *et al.*, *Lancet*, 1966, **1**, 1225.
- 12 Rodey, G. E., *et al.*, *Blood*, 1969, **33**, 813.

Haemodialysis and Forced Diuresis for Tricyclic Antidepressant Poisoning

SIR,—The encouraging experience of Drs. H. W. Asbach and H. W. Schüler (18 May, p. 386) in using haemodialysis for the treatment of two young children with severe imipramine poisoning has stimulated us to report a recent dramatic clinical success we have had in an adult who took a massive overdose of tricyclic antidepressants.

A 50-year-old woman with a long history of depression ingested a large quantity (precise number of tablets not known) of imipramine and amitriptyline, possibly with a few tablets of opi-pramol, trifluoperazine, haloperidol and thiothixene. About six hours later she was admitted to hospital deeply comatose, areflexic, and hypothermic, with a sinus tachycardia and a systolic blood pressure of 80-90 mm Hg. An electrocardiograph showed an intraventricular conduction defect. The patient was intubated and given gastric lavage. She was treated with continuous gastric aspiration, body warming, intermittent positive pressure respiration, and a noradrenaline infusion. Forced diuresis was commenced through the central venous pressure line.

Nine hours after admission the patient had shown no signs of clinical improvement and it was decided to commence haemodialysis. An arterio-venous shunt was placed in her leg and dialysis started using two hollow fibre kidneys in parallel (dialysate flow rate 500 ml/min; arterial flow rate 150 ml/min) and a standard dialysate solution. Dialysis was continued for six hours. Within four hours of commencing dialysis the vital signs improved significantly and there was a definite improvement in the level of consciousness. This improvement was maintained over the next 48 hours and the patient made a full recovery.

Chemical analyses of the gastric lavage showed the presence of a large amount of imipramine and amitriptyline and a very small quantity of a phenothiazine. Plasma imipramine levels were determined by a modification of the method of Grove and Halliwell.¹ No attempt was made to distinguish between imipramine and its fluorescent metabolites and results are reported as imipramine. Samples were taken soon after admission and during the period of haemodialysis. The plasma imipramine level fell from 6.3 µmol/l. to 1.7 µmol/l. within 14 hours. These results are shown in the figure. According to Niyogi,² 1.7 µmol/l. is the upper limit of the therapeutic range for plasma imipramine.

During the period of forced diuresis the urine volume was 23,370 ml, and contained 151.5 µmol (43 mg) of imipramine. No imipramine was detected in the dialysate when it was analysed by the same method as used for plasma. A 20-fold