New Curriculum

SIR,—I was interested to read Professor D. C. Morrell's letter (28 September, p. 804) in which he quite rightly pointed out that general practitioners ought to contribute to teaching in sexual matters. My letter (31 August, p. 578) was in response to a leading article concerning the teaching of sexual matters (29 June, p. 686), but I could have expanded my comments to cover other important areas of knowledge which medical students should have when they qualify but which are often inadequately taught.

The teaching of geriatric medicine should, of course, be a team effort including not only doctors (specialists and general practitioners) but also the many other members of the therapeutic team such as nurses, occupational and physiotherapists, and social workers. The main benefit of effective teaching about the elderly would be to help students who are not intending to be specialists to offer better medical care to their patients, and this would particularly apply to intending general practitioners. Another important benefit would be that a few medical students would consider specializing in geriatric medicine. Most present consultants in geriatrics went into geriatrics because of the particular circumstances in which they found themselves after qualifying rather than because they considered it as a career when they were students.

More academic departments of geriatric medicine are needed, but a multiplicity of professorial departments might defeat the object I am advocating, as research would proliferate and staff would probably be lured away from other teaching and non-teaching centres. All bodies concerned with developing good teaching programmes for undergraduates should be actively supporting multidisciplinary teaching about medicine in old age in curricula. Geriatric physicians and general practitioners should be asked to help provide this teaching.—I am, etc.,

MICHAEL GREEN

Royal Free Hospital,
London N.W.3

Hyperglycericacidaemia with Hyperglycinaemia: A New Inborn Error of Metabolism

SIR,—In a severely motor and mentally retarded patient we have found large amounts of glyceric acid and glycine in the blood and urine. Hyperglycericacidaemia combined with hyperglycinaemia has not been reported previously and our results suggests a new inborn error in the metabolism of glycine and glyceric acid.

The patient is the second child of healthy parents of Serbian origin. Birth weight 3,350 g. Delivery was normal. From birth there was severe hypotonia with weak sucking power. At 2 months old the patient was admitted to the neuropaediatric ward because of severe neurological symptoms including myoclonic jerking, cerebellar fits, and an almost classic chorea minor pattern. He remained severely motor retarded and no eye contact could be obtained. He has failed to thrive physically and mentally and is now 2 years old with severe developmental handicap. Occasional grand mal seizures have now been added to the clinical picture. There have been no acidotic episodes, however, and no symptoms of renal disease. Intravenous pyelography was normal.

By thin-layer chromatography of the urinary amino-acids large amounts of glycine were repeatedly found. Quantitative determinations by column chromatography gave the following glycine values: in serum 0.88–1.07 mmol/l. (normal controls <0.30 mmol/l.) and in urine 47.53–61.14 mmol/l. (normal controls <3.0 mmol/l.). The daily urinary output of glycine was approximately 10 mmol/24 hr (20 times the excretion in normal children of the same age). When the urine was acidified and extracted with ethyl acetate analysis by gas chromatography of the trimethylsilyl derivatives revealed a large abnormal peak. Combined gas chromatography and mass spectrometry of this peak compound gave a mass spectrum identical with that of the trimethylsilyl derivatives of an authentic glyceric acid. Analysis of eight urine specimens collected during a six-month period showed concentrations of glyceric acid ranging from 33 to 187 mmol/l. (excretion about 1.5–2.5 g/day). The concentration of glyceric acid in the serum was about 1.2 mmol/l. This compound was detected neither in urine nor in serum from control subjects. The excretion of oxalic acid and glyoxylic acid in the urine of the patient was in the low normal range.

The patient has the typical clinical symptoms of non-ketotic hyperglycinaemia.¹ Hyperglycericacidaemia and hyperglyceric-aciduria, however, have not been described in this syndrome. Hyperglycericaciduria is found in oxalosis type II (glyceric acid dehydrogenase deficiency) together with hyperoxaluria but these patients do not have hyperglycinaemia and apparently not hyperglycericacidaemia either.² The combination of hyperglycericacidaemia and hyperglycinaemia with excretion of large amounts of glyceric acid and glycine in the urine but absence of raised amounts of oxalic acid and glyoxylic acid points to a block in a hitherto undescribed pathway in the metabolism of the compounds mentioned.

This work was supported by the Danish State Medical Research Foundation.

—We are, etc.,

N. J. BRANDT

Department of Teratology

S. BRANDT

Department of Child Neurology,
Rigshospitalet,
University of Copenhagen

K. RASMUSSEN

Department of Clinical Chemistry

F. SCHØNHEYDER

Department of Medical Biochemistry,
University of Århus,
Denmark

¹ Childs, B., et al., *Pediatrics*, 1961, 27, 522.

² Williams, H. E., and Smith, L. H., *New England Journal of Medicine*, 1968, 278, 233.

Uticillin

SIR,—I have been disturbed recently to receive from Beecham Research Laboratories information regarding Uticillin (carfecillin sodium), an oral penicillin which is hydrolysed in the body to carbenicillin. Advertisements for this drug have recently appeared in the *B.M.J.* in which the specific use for this drug is stated to be the treatment of urinary tract infection.

To advertise this drug, which is merely an oral form of carbenicillin, in the way that has been done is, to put it mildly, an error of judgement and probably courts disaster. It is generally agreed that carbenicillin should be reserved for infections with *Pseudomonas aeruginosa* and cases of septicaemia where it is used in combination with gentamicin. Its use in other, less serious situations where other antibiotics may be equally or more effective is to be deprecated, as

such use can lead to populations of *Ps. aeruginosa* which are resistant to carbenicillin, thus rendering it useless when a serious infection arises.

Its routine use in acute episodes of urinary tract infection cannot be justified as there are other equally effective antimicrobials, resistant organisms to which would not be such a disaster, though admittedly undesirable. It is probably unthinkable that Uticillin should be used prophylactically in urinary tract infection in view of the development of R-factor-mediated carbenicillin resistance. Suitable prophylactic antimicrobials in this situation are nalidixic acid and nitrofurantoin, to which so far R-factor resistance has not been found, or co-trimoxazole, to which it is very rare indeed.¹

The medical profession has a great responsibility in ensuring that yet more resistant strains of bacteria do not infest our hospitals and spread within the community. This responsibility rests not only upon hospital physicians and general practitioners who are exposed to advertising pressures, but also upon those medical journals which accept such advertising copy. It is a moral responsibility that must be borne by medical men in whatever field they work.—I am, etc.,

M. J. BENDALL

St. Mary's Hospital,
Colchester

¹ Grünberg, R. N., Leakey, A., and Bendall, M. J., in Symposium on Advances in Urinary Tract Infection, International Society of Nephrology, 23–24 September 1974. To be published.

SIR,—We have recently received literature from Beecham Research Laboratories announcing the introduction of Uticillin (carfecillin sodium), a phenolic ester of carbenicillin which can be given orally and on absorption liberates a bacteriologically inactive moiety and carbenicillin. The carbenicillin is excreted in the urine, where concentrations achieved are sufficient to treat carbenicillin-sensitive strains of *pseudomonas*. We are disturbed to note that the manufacturers suggest that sensitivity tests to carfecillin should be reported in addition to those for carbenicillin and in fact that the use of carbenicillin discs for routine testing or urine is not recommended. Since the active component of Uticillin is carbenicillin this recommendation is scientifically unjustifiable. Furthermore, it is not the purpose of a bacterial sensitivity test to report which preparation of an antibiotic is suitable for administration by the various routes. In addition we would point out that at least one other oral form of carbenicillin, indanyl carbenicillin,¹ has been developed and in vitro shows a greater degree of activity against certain micro-organisms than does carbenicillin, which is the active principle liberated after absorption.

The claim that Uticillin has a different clinical useage from carbenicillin is not justified and in view of the emergence of drug resistance it is particularly important that it should not be used for treating infections other than those due to *pseudomonas* organisms or the rare strains of bacteria which are resistant to all other antibacterial agents.

Finally, we would point out that full-page advertisements for Uticillin have appeared in the *B.M.J.* which made no reference to the fact that this antibiotic