sympathetic afferent fibres or B1 receptor sites in the lung which are known to stimulate ADH secretion.³ The involvement of the vagus nerve by a bronchogenic carcinoma and associated inappropriate ADH secretion is described by Bower *et al.*⁶ However, they were also able to substantiate the antidiuretic activity of the tumour and postulated this and the possibility of distortion of vagal afferent impulses as a cause of inappropriate ADH secretion in their patient. The stress of the acute disease may also play a part, though the ADH response to stress has not been shown to be sustained.³—I am, etc.,

J. B. MYERS

University Department of Medicine,
Royal Melbourne Hospital,
Victoria, Australia

- 1 Bryant, D. H., *Medical Journal of Australia*, 1972, 1, 1285.
- 2 Phillips, M. E., *et al.*, *British Medical Journal*, 1974, 2, 149.
- 3 Schrier, R. W., and Berl, T., *Kidney*, 1974, 7, 1.
- 4 Weiss, H., and Katz, S., *American Review of Respiratory Diseases*, 1965, 92, 609.
- 5 Utz, J. P., *et al.*, *New England Journal of Medicine*, 1969, 280, 264.
- 6 Bower, B. F., Mason, O. N., and Forsham, P. H., *New England Journal of Medicine*, 1964, 271, 934.

Hypertension and Myocardial Infarction

SIR,—I was interested to read your recent leading article (5 January, p. 1) and the views of Dr. F. T. B. Lovegrove (4 May, p. 279), Dr. I. McD. G. Stewart (27 July, p. 251), and Dr. D. M. D. Lambert (14 September, p. 685).

Ex Africa semper aliquid novi; this statement, traditionally ascribed to the Greeks by Pliny the Elder, is applicable to hypertension and myocardial infarction in the African. While hypertension in the African is common, myocardial infarction is rare. In the urban African in Johannesburg hypertension and its complications is believed to be the second most common cause of death in the adult after violence.¹ Yet myocardial infarction is rare.

In my study² of 1,000 hypertensive patients (500 Africans and 500 Indians) who were followed up over a period of 13 years the incidence of cardiac changes is shown in the table. It will be noted that while

Cardiac Change	Africans (%)	Indians (%)
E.C.G. Left ventricular hypertrophy	23	22
Clinical cardiomegaly	29	26
Ischaemia heart disease	—	12
Chest radiograph: left ventricular hypertrophy	30	35
Normal chest radiograph	70	65
Congestive heart failure	16	8

congestive cardiac failure due to hypertension occurred in 16% of the Africans, ischaemic heart disease did not occur. This is in contrast to the Indian population, in whom myocardial infarction or angina pectoris occurred in 12% of cases. Our hospital has 2,000 beds and serves a population of 2 million; there are approximately 800,000 patients attending the outpatient department and about 99,000 are admitted to the hospital per year (African 83%, Indian 17%). While myocardial infarction is common in Indians, the number of cases of myocardial infarction or angina pectoris in Africans does not exceed 10 per year. The incidence of coronary atheroma in Africans

at necropsy is far less than in the European or Indian population in Durban.³ The Framingham study⁴ found that increased serum lipids, hypertension, cigarette smoking, overweight, and diabetes mellitus were predisposing factors for myocardial infarction. The African has all the above predisposing factors with the exception of increased serum lipids. The rarity of myocardial infarction in the African could be because the mean serum lipids level are lower than in other races. Walker⁵ found that in the 30-39-year age group a serum cholesterol level above 220 mg/100 ml was present in 55% of whites, 45% of Indians, 25% of urban Bantu, and 10% of rural Bantu in South Africa; the proportion for whites was much the same as that reported in the Framingham study—namely 58%. However, hypertension is not uncommonly associated with cerebral atheroma and atheroma in the aorta. In our study of 1,000 patients² cerebrovascular episodes occurred in 5% of African and 10% of the Indian patients. We feel that a high incidence of ischaemic heart disease is related to underlying diabetes mellitus and associated hypercholesterolaemia.

Our work suggests that hypertension alone is not an important predisposing factor in the aetiology of myocardial infarction. It may be an important correlate with other factors that predispose to myocardial infarction rather than a causal factor. Similar work in Zaïre⁶ has shown that while hypertension is common and not uncommonly produces congestive cardiac failure, myocardial infarction does not occur.—I am, etc.,

Y. K. SEEDAT

Department of Medicine,
University of Natal,
Durban

- 1 Seftel, H. C., *Medicine and Society in South Africa—Some Plain Thinking*, p. 14. Johannesburg, Witwatersrand University Press, 1973.
- 2 Seedat, Y. K., and Reddy, J., *South African Medical Journal*, 1974, 48, 816.
- 3 Wainwright, J., *Lancet*, 1961, 1, 366.
- 4 Kannel, W. B., *et al.*, *Annals of Internal Medicine*, 1961, 55, 33.
- 5 Walker, A. R. P., *South African Medical Journal*, 1973, 47, 85.
- 6 Marcovayannopoulou-Fojas, H., *Abstracts of Proceedings of World Congress of Cardiology*, 1974. In press.

Variability in Response to Drugs

SIR,—I find it remarkable that in his survey of "Variability in Response to Drugs" (12 October, p. 91) Professor M. D. Rawlins contrived to avoid any clear acknowledgment that psyche influences soma. Is it not general experience that "rational drug therapy"—like many another logical, technical process—often fails for "irrational" reasons?

Just as important as the patient's acetylase status is his psychological status. No doubt they interact. So let us by all means estimate plasma concentrations and gain a better understanding of tissue sensitivity and dosage control. But let us also acknowledge that a man is more than a biochemical system. Many human "diseases," like epilepsy and depression, are relieved by treatment methods which may incorporate the use of drugs but also take account of a person's inner needs and outer relationships.—I am, etc.,

J. K. W. MORRICE

Ross Clinic,
Aberdeen

Medical Management of Heartburn

SIR,—Your leading article on this subject (12 October, p. 64) begins by stating that acid reflux causes heartburn; but alkaline reflux can cause severe heartburn also. The article goes on to mention the variable and largely unpredictable response to alkalis and other remedies and discusses a number of more scientific forms of treatment based on recent research. A very curious statement is that "attempts to improve general muscle tone should strengthen the oesophageal sphincter." There is surely no evidence for this whatever, and it is a matter of common observation that many with otherwise excellent health and good muscle tone suffer from heartburn.

My main criticism is, however, that no mention is made of the fundamental reasons for the prevalence of heartburn—an incompetent sphincter and its associated condition, sliding hiatus hernia. These conditions are a feature of western civilization with its tendency to overeat. It may be that the character of the diet, with a high proportion of refined carbohydrate, leads to disorders of gastric secretion and gastro-oesophageal motility; but a simple remedy for heartburn and reflux (in the absence of gross anatomical disorders at the oesophago-gastric junction such as hiatus hernia) is to avoid eating unless appetite is present, as has been emphasized by Cleave.¹ The eating of meals as a matter of social habit rather than necessity is the cause of much heartburn, and it is not necessary to invoke complex pharmacology for its diagnosis or treatment.—I am, etc.,

KENNETH S. MULLARD

Diss, Norfolk

- 1 Cleave, T. L., *The Saccharine Disease*, p. 176. Bristol, Wright, 1974.

Anonymous Appointments

SIR,—The expansion of hospital bureaucracy is nowhere better illustrated than in the much-read appointments pages of your journal. It is frequently necessary to read the whole of a box of small print to discover at which hospital a post will be held. Occasionally this is only hinted at by the address of the personnel officer. In no such case is the name of the group hospital management committee left in doubt. This is less important information to a doctor than the name of the place where he will live and work. It could be appended or even omitted.—I am, etc.,

M. PRENDERGAST

London S.E.1

Economies in the N.H.S.

SIR,—The joint letter from certain doctors at the Royal United Hospital, Bath (12 October, p. 109), while showing a commendable concern for economy in the National Health Service, unfortunately does not show the same concern for accuracy in terminology or convey a correct interpretation of Department of Health requirements regarding the provision of sterile water for operating theatre use.

The Department of Health has not specified that sterile topical water for use in operating theatres should be particle-free but has indicated that it should be freshly