As many dental anaesthetics are still given in the upright position, it seems important in such circumstances for the anaesthetist (1) initially, to check that the back rest of the dental chair will depress quickly and smoothly (some do jam) enabling the patient to be supinated at once; (2) to be conversant with the procedure for dropping the arm rest and moving the patient sideways out of the chair if the back rest jams; and (3) to monitor the pulse throughout the anaesthetic.—I am, etc.,

BRYAN LEWIS

Birmingham

Dexamethasone in Malignant Hyperpyrexia

SIR,—A child of 6 years weighing 22.5 kg about to undergo adenoidectomy was given 100 mg thiopentone, 30 mg suxamethonium, and 1% halothane. After approximately two minutes he was sweating and felt very hot; anaesthesia was discontinued. At this stage his temperature was 39.25°C and there were no signs of muscle rigidity. In view of the report by Dr. F. R. Ellis and others (2 November, p. 270) he was given 4 mg of dexamethasone intravenously as an initial steroid dose. His rectal temperature fell rapidly to 38°C within three minutes. No further treatment was given and cooling was not required. His postoperative serum creatine phosphokinase level was 24 U/l. Further investigations are being undertaken and will be reported; it seems probable, however, that instant steroid therapy is extremely important in any patient with suspected malignant hyperpyrexia.—We are, etc.,

D. G. RAITT
A. J. MERRIFIELDRoyal Air Force Hospital,
Ely, Cambs

Florence Nightingale's Illness

SIR,—In his review of my book *Creative Malady* (16 November, p. 414), Dr. Henry Rollin takes issue with me on the nature of Miss Nightingale's illness.

The essential facts are these. In July 1856 she returned from her arduous and successful task of rescuing the remains of the British Army in the Crimea from administrative neglect and incompetence. Against intense opposition she succeeded in getting a Royal Commission established. She did most of its work and drafted its report at great speed. In August 1857 she became ill and remained an invalid till she died in 1910. I suggested that this illness was a psychoneurosis occasioned by the conflict between her passionate desire to avenge her "murdered children" (the British soldiers who died in the Crimea) and the dissipation of her time and energy, particularly by her mother and sister. Her illness enabled her to succeed. Dr. Rollin maintains "that one outstanding feature was her obsessionism, and the vulnerability of obsessionals is well known. Not surprisingly, in 1857, when her prodigious report was finished, she cracked. All her symptoms from then on could be explained in terms of an irreversible depressive illness, aggravated later, as Sir George suggests, by the menopause and possibly an addiction to morphine." What was truly remarkable was the vast output of lucid, cogent, and well-informed reports, minutes,

warrants, and regulations and summaries for the Minister's use that flooded from the invalid's couch between 1857 and 1868. Is this characteristic of a depressive illness and did Miss Nightingale crack? She certainly changed her way of life, but she was even more formidable on the invalid's couch than she had been in society, as the War Office learned to its cost.—I am, etc.,

GEORGE PICKERING

Moulsoford-on-Thames,
Berks

Safety and Fibreoptic Bronchoscopy

SIR,—The letter from Dr. I. W. B. Grant (23 November, p. 464) stressing the inadequacies and dangers of fibreoptic bronchoscopy merits further discussion. Here we may usefully contrast his experience with that of 193 American bronchoscopists surveyed in a recent paper.¹ This survey covers both large and small centres and is based on the massive total of 24,521 fibreoptic bronchoscopies, arguably giving a more solid base for generalization than a single centre's experience.

Dr. Grant implies throughout his letter that there is a measure of danger in performing fibreoptic bronchoscopy in centres without available rigid bronchoscopy. The American survey, however, found very low rates for both minor complications (0.2%) and major complications (0.08%). Even more tellingly, examination of their data shows only three cases in the 24,000 where treatment of a complication could have been helped by the presence of rigid bronchoscopy or thoracic surgery. While caution is always advisable in assessing new techniques, this widely based survey can surely offer substantial reassurance.

The letter questions the reliability of local anaesthesia for fibreoptic bronchoscopy and even labels it a "barbarous practice." This contrasts strongly with our experience in performing over 150 fibreoptic bronchoscopies satisfactorily under local anaesthesia. This impression is confirmed by the paucity of complaints found on systematically questioning patients afterwards. Local anaesthesia was also chosen for 84% of fibreoptic bronchoscopies in the American survey. Especially in a medical system based on private practice a truly barbarous method would surely be less popular.

Problems with obscuration of bronchoscope vision by mucus and blood were also mentioned in the letter. These difficulties do not appear substantial in most published series, which confirms our own experience that these problems are not insurmountable. Another suggested disadvantage was the difficulty in obtaining adequate biopsy samples with the fibreoptic technique. Our experience, on the other hand, confirms that of several large series which show high biopsy success rates, often higher than with rigid techniques.

Restriction of fibreoptic bronchoscopy to centres already equipped with rigid bronchoscopy is advocated by Dr. Grant on grounds of safety and cost. The complication rates discussed above suggest that fibreoptic centres with normal supporting facilities pose few safety problems. The costs of running such centres can be substantially reduced by sharing an endoscopy room with gastrointestinal endoscopy, which is at

present expanding rapidly. Most of the costs of this facility can then be shared by the two specialties—trained staff, cold light source, suction apparatus, resuscitation equipment, etc.

Doubtless the strongest argument for expansion of fibreoptic centres is simply the increasing number of bronchoscopies needed each year to meet the ever-growing tide of lung cancer. Some 15,000 bronchoscopies are needed annually now.² Sooner or later existing centres with rigid bronchoscopy will be unable to cope. Time will force the establishment of more fibreoptic centres. In preparation for this development, we would be well advised to include fibreoptic bronchoscopy in the training of the next generation of chest physicians.—I am, etc.,

J. B. MACDONALD

Sully Hospital,
Cardiff

¹ Credle, W. F., Smiddy, J. F., and Elliott, R. C., *American Review of Respiratory Disease*, 1974, 107, 67.

² Benetts, F. E., *British Journal of Hospital Medicine*, 1973, 10, 71.

Sticky Eye in the Newborn

SIR,—The recent comment by your expert (26 October, p. 222) that sticky eye in the newborn caused by T.R.I.C. (trachoma-inclusion-conjunctivitis) strains of chlamydia group A is of "a similar order of rarity" to gonococcal infection ("one case a year at one large maternity unit") and "usually does not appear until the second week, perhaps when the baby has been sent home" differs from our experience in Liverpool maternity units over the past 13 months.

We have investigated 25 newborn infants with mucopurulent conjunctivitis developing between three and eight days after birth; these were unselected cases except in so far as some had failed to respond to chloramphenicol or neomycin eye drops. Eye swabs were examined by the non-irradiated McCoy cell technique.^{1,2} Chlamydia were isolated, with primary inclusion counts ranging from 1 to 9,000 per coverslip (median count 600), from eight infants (32%). Chlamydia were also isolated from cervical swabs of four mothers of these positive infants (counts 300-5,000) and from urethral swabs of one of two fathers examined; two mothers yielding negative swabs had already received antibiotics for puerperal pyrexia and one refused examination. In a further 12-day-old infant with sticky eye, already treated four days with antibiotics, chlamydial culture was negative, but a cervical swab of his mother, who had an offensive lochial discharge, was positive. In another family eye swabs of a 3-day-old baby yielded only gonococci, but gonococci and chlamydia were isolated from cervical swabs of the mother. Overall, there were five proven cases of gonococcal ophthalmia, from one of which chlamydia were also isolated.

All the chlamydia-infected eyes improved clinically within a few days of beginning chlortetracycline eye ointment, given for three to four weeks. After treatment further eye swabs were negative in six cases; in the seventh case culture was again positive two weeks after treatment and the eighth infant was found positive two weeks after defaulting after two weeks' treatment.

Chloramphenicol therapy may suppress

chlamydial infection, thus making laboratory diagnosis difficult, but will not eradicate the disease. Neomycin is not active against chlamydia, but is reasonably effective against bacterial infections and should be the initial antibiotic of choice until swabs can be taken and examined for chlamydia. The persistence of conjunctivitis after neomycin should lead the clinician to investigate the possibility of chlamydial infection in both the infant and its parents.—We are, etc.,

ELISABETH REES

Department of Venereology,
Liverpool Royal Infirmary

DEREK HOBSON

Department of Medical Microbiology,
University of Liverpool,
Liverpool

1 Gordon, F. B., and Quan, A. L., *Proceedings of the Society for Experimental Biology and Medicine*, 1965, 118, 354.

2 Hobson, D., et al., *Lancet*, 1974, 2, 555.

New Curriculum

SIR,—Dr. M. F. Green (9 November, p. 344) wishes medical students to receive instruction in geriatric medicine but fears that "a multiplicity of professional departments might defeat the object... as research would proliferate and staff would probably be lured away from other teaching and non-teaching centres." I think Dr. Green's fears can be allayed. As yet only a fraction of British medical schools have established university departments of geriatric medicine, and many medical students who are graduating today and who will still be in practice well into the 21st century have received no formal instruction in the principles and practice of geriatric medicine. The proliferation of research in geriatric medicine can hardly be a bad thing, especially if the research remains closely related to improvements in clinical practice, as has been characteristic of a great deal of the research published by geriatric physicians and indeed of the important contributions made by Dr. Green himself. It is true that university departments will tend to attract the best members of the profession, but that is surely nothing to fear. They will then be in a position to influence future graduates to enter this progressive and rewarding specialty. Dr. Green is of course correct to say that the teaching of medical students in geriatric medicine and in other branches will benefit greatly from the contribution of all practising geriatric physicians as well as from that of general practitioners and members of the health professions with whom geriatric physicians are accustomed to work so closely.—I am, etc.,

BERNARD ISAACS

Department of Geriatric Medicine,
Lightburn Hospital,
Glasgow

Tuberculosis and Renal Amyloidosis

SIR,—We concur with the observation of Professor A. C. Kennedy and others (28 September, p. 795), who described pulmonary tuberculosis as the leading cause of renal amyloidosis in the west of Scotland. Renal amyloidosis is still considered to be the second commonest cause of nephrotic syndrome in India, especially in Delhi (36.2% of cases),¹ and despite earlier detection and

treatment of tuberculosis this disease will remain the commonest aetiological association of amyloidosis for some time to come in developing countries like ours.

Out of 200 cases of renal amyloidosis confirmed by biopsy seen by us in the past 10 years tuberculosis (pulmonary 125, lymphadenitis 12, spinal 8, and intestinal 5) was the commonest cause (75%). Other causes were suppurative lung diseases (13%), rheumatoid arthritis (5%), ankylosing spondylitis (3%), and metastatic adenocarcinoma of liver and lung, multiple myeloma, Hodgkin's disease, and leprosy (1% each). Most of the patients (70%) were men and their ages ranged from 20 to 72 years (mean 38 years). Most of the cases progressed to renal failure; hypertension was observed in 6% and renal vein thrombosis in 1%.

We find renal biopsy the best method of clinching the diagnosis of renal amyloidosis and when this is impossible rectal biopsy is advisable. A liver biopsy may appear normal, despite enlargement due to fatty infiltration, and hence is not always dependable. Most of our patients died within two to three years. A few still survive after more than four years. Two patients showed a regression of amyloid material in the glomeruli on serial biopsy after adequate treatment of the underlying disease and high doses of vitamin C. In established renal amyloidosis careful search may reveal tuberculous lymphadenitis, intestinal tuberculosis, occult malignancy, or sometimes healed tuberculosis. By and large the duration of the underlying disease correlates well with the severity of the amyloidosis and the prognosis.²—We are, etc.,

S. N. A. RIZVI

P. D. GULATI

HARI VAISHNAVA

Maulana Azad Medical College,
New Delhi,
India

1 Vaishnava, H., et al., *Journal of the Association of Physicians of India*, 1969, 17, 641.

2 Gulati, P. D., et al., *Postgraduate Medical Journal*, 1970, 46, 137.

Medical Centre for Unsettled Young People

SIR,—Dr. Patrick J. Day's Personal View (9 November, p. 340) described a refreshing "professional" approach to a group whose relations with standard medical institutions are deficient, to say the least. The evidence suggests to us that unattached young people, many of whom are less settled than the community Dr. Day lived with, are increasing in number. This is certainly so in London.

A working group representative of the New Horizon Youth Centre, the Soho Project, Centrepoint, the Standing Conference on Drug Abuse, and the Student Health Service is working through the Campaign for the Homeless and Rootless with various medical personnel to produce proposals for a medical centre for young people in central London. As the proposals stand it would be an extension of the Health Service providing, at street level, a walk-in clinic whose staff and procedures would be finely tuned to the needs of the patient. Not least of its assets would be its referral function, introducing young people to appropriate sources of care and treatment. The clinical function would be relatively basic, providing first aid and medical advice and, when possible, linking up the patient with a general

practitioner, whose numbers in central London are declining. Social workers at voluntary projects working with mobile young people have more and more difficulty in finding a doctor to accept cases.

The initial responses from the area health authority and from social services departments have been very encouraging and we hope for a sympathetic hearing from the Department of Health and Social Security when we present our proposals in the new year.—I am, etc.,

ANNE DAVIES

Assistant Director,

Campaign for the Homeless and Rootless

London W2

Hypoglycaemia Associated with Intrathoracic Mesothelioma

SIR,—The association of hypoglycaemia with neoplasms other than islet cell tumours is probably underdiagnosed.¹ In Hong Kong hypoglycaemia in association with carcinoma of liver in 30% of cases has been well recognized.² When an intrathoracic mesothelioma close to the right hemidiaphragm is associated with hypoglycaemia the displaced and apparently enlarged liver may cause diagnostic difficulty.³ Prompt thoracotomy may not only settle the diagnosis but also remove the cause of hypoglycaemia,⁴ as the following case shows.

A woman aged 64 years was admitted to hospital with a short history of dizziness in the morning followed by coma. On admission her blood sugar was 16 mg/100 ml. It fluctuated between 6-30 mg/100 ml over the next two weeks. A chest radiograph showed a large opacity in the right lower zone merging into the liver shadow. As the liver was palpable a diagnosis of carcinoma of liver had to be considered, but owing to the severity of the condition a thoracotomy was decided on. This was performed with the patient in coma despite an intravenous 10% dextrose infusion. A large, well-encapsulated tumour was found in the pleural cavity attached by a few small pedicles to the mediastinal pleura and the right lower lobe (see fig.). Histology showed features of mesothelioma.

The patient made a smooth recovery. The blood sugar stayed consistently above 100 mg/100 ml. There was no recurrence up to six months.—We are, etc.,

J. S. M. LEUNG
S. T. TANG

University Thoracic Unit,
Kwong Wah Hospital,
Hong Kong

1 Unger, R. H., *American Journal of Medicine*, 1966, 40, 325.

2 McFadzean, A. J. S., and Young, T. T., *Archives of Internal Medicine*, 1956, 98, 720.

3 Maier, H. C., and Barr, D., *Journal of Thoracic and Cardiac Surgery*, 1962, 44, 321.

4 Spry, C. J. F., Williamson, D. H., and James, M. L., *Proceedings of the Royal Society of Medicine*, 1968, 61, 1105.