CORRESPONDENCE

Correspondents are asked to be brief

SI Units

C. D. Needham, F.R.C.P.; G. H. Hall, F.R.C.P.;
C. G. Eastwood, M.D. 767

Cross-immunity to Influenza B Virus Strains

L. M. de Silva, M.B., PH.D., others 767

Vaccination against Whooping Cough

J. S. Dodge, M.D. 768

Respiratory Distress Syndrome in the Newborn

V. Dubowitz, F.R.C.P.; A. R. Wilkinson,
M.R.C.P.; W. P. Bevington, M.B.; Carolyn
E. M. Jones, M.R.C.P. 768

Oral Contraception and Increased Formation of High Molecular Weight Derivatives of Fibrinogen

F. Asbeck, M.D., and others 769

Normal Short Children

J. Apley, F.R.C.P., and D. R. Davis, F.R.C.P. 769

Pseudomonas Septicaemia following Superficial Colonization

A. S. D. Spiers, F.R.A.C.P. 770

Prevention of Exercise-induced Asthma by Indoramin

F. J. Prime, F.R.C.P.ED.; G. T. Dixon, PH.D.,
and others 770

Drugs for Rheumatoid Arthritis

P. J. Rooney, M.R.C.P., and others 771

Distribution and Supervision of Oral Contraceptives

E. C. Bramwell, M.B.; N. A. Chisholm,
L.R.C.P.ED. 771

Benign Intracranial Hypertension Associated with Nitrofurantoin Therapy

D. B. Sharma, M.D., and A. James, M.R.A.C.P. 771

Consultant Contract

T. S. Wilson, M.D., and E. V. B. Morton,
F.R.C.P.ED.; J. P. Hopewell, F.R.C.S.; T. F. P.
Mackintosh, M.R.C.P.ED.; J. P. Lythgoe,
F.R.C.S. 772

Sanctions

Kathleen M. Frith, F.R.C.O.G. 772

SI Units

SIR,—Professor D. N. Baron's neat presentation (30 November, p. 509) of this newest notation system reads like one of those ever-recurring enthusiastic special pleadings heard on behalf of the latest "marvellous" new international language. It has the usual neologisms and peculiar conventions which allow the "in" people cosily to twitter among themselves with a great feeling of superiority. But the prime purpose of a system of notation is efficient communication between all concerned, and that embraces a great diversity of people, only a small minority of whom have laboratory and/or mathematical expertise as their primary interest.

The multiplicity of prefixes and their related abbreviated symbols is bound to lead to confusion, and a great deal more if one adds to the fun by putting in some negative powers. This type of precisionist symbolism is perfectly good for specially trained people but is bound to lead to dangerous confusion in the hands of many simple users. That the new international notation is highly suitable for calculation of moon missile trajectories etc. does not mean that it is optimal in practical medicine, or indeed in some other spheres—for example, what will the average garage attendant say on being asked to inflate one's tyre to so many kPa? Even Professor Baron apparently does not find it all that easy, for surely a plasma concentration of phenobarbitone of 46 mg/100 ml does not become the 0.2 mmol/l that he suggests, but 1.98 mmol/l.

The argument that the SI system in its full flowering makes for clarity and efficiency lacks conviction. One can envisage a gram, and a mole of something of which one knows the molecular make-up can be meaningful—for example, NaCl; but which of us clinicians really pretends that he knows off-hand the molecular constitution of bilirubin? And again, in pressures, I think that the height of a column of mercury as seen in Aberdeen is more meaningful than some

contorted calculated derivation based on a lump of platinum held in a laboratory in Paris.

I submit, Sir, that no proper consideration has been given to (1) the views of the real practical consumers of these data and (2) the cost of indulging this newest fad.—I am, etc.,

DOUGLAS NEEDHAM

Woodend General Hospital,
Aberdeen

SIR,—From the resigned tone of your leading article (30 November, p. 490) it might be concluded that it is useless to try to resist the introduction of SI units. This is a pity as there are many weighty reasons against their use in clinical medicine.^{1,2}

(1) They are different from those we are used to. Mistakes in interpretation are bound to be made; patients will suffer. (2) SI units are often of the wrong size. Using pascals to measure blood pressure is like measuring a cricket pitch in inches and the kilopascal is too big (7.5006 mm Hg = 1 kPa). (3) Try and imagine the cost of re-printing all the pathology forms, let alone recalibrating equipment. In the petrochemical industry, I am told, the conversion of one plant will cost £2m. (4) The requirements of medicine are not the same as those of science; in particular, the use of molar measures instead of mass under certain circumstances (for example, drug concentrations in blood) is entirely inappropriate. (5) SI units are not universally employed, may never be, and may well be changed again. Further developments can be expected, with further damage to patient care.

In essence, the imposition of SI units on the medical profession is about as justifiable and logical as it would be to make all our population speak French because our Common Market negotiators in Brussels have to. "Consultation" about this vital matter

has been a hollow sham. It is essential that careful pilot studies with consumer (that is, clinical) participation be continued.—I am, etc.,

G. H. HALL

Exeter

¹ With, T. K., *Lancet*, 1974, 2, 1315.
² Hall, G. H., *Lancet*, 1974, 1, 1006.

SIR,—Surely Professor D. N. Baron (30 November, p. 509) is in error when he says that the original metric system was based upon three independent base units. As originally enunciated in the French revolutionary period the metre was the sole base unit. The three ordinary dimensions were expressed as multiples or submultiples; volume was based upon the cubic metre and mass upon that of the cubic metre of water at its greatest density (4°C). Of course additional measurements are now needed, but it seems a pity to depart from one basic unit.

The original system used to be prettily illustrated by a small ruler provided by the Metric Association. This was a decimetre long, a centimetre wide, and a millimetre thick and its weight in grams was also its specific gravity. Hung upon a cord of the right length, it became a pendulum for the measurement of time.—I am, etc.,

CYRIL G. EASTWOOD

Cambridge

Cross-immunity to Influenza B Virus Strains

SIR,—Outbreaks of influenza B infection due to two variants of the old influenza B virus were reported throughout the United Kingdom last winter.¹ Of particular interest was the occurrence of clinical illness mainly in individuals between 1 and 24 years of age, and especially in children 5-9 years old.²

We have had the opportunity recently of studying sera from an outbreak reported in September 1973 in a north of England

boarding school due to the intermediate variant strain of influenza B virus. Paired sera from six students (aged 11-15) with infection proved by virus isolation and/or a fourfold rise in antibody level (haemagglutination inhibition or complement fixation test) were subsequently tested by the fluorescent antibody technique for class-specific fluorescent antibody to the three influenza B strains (old influenza B virus and the two recent variants, the intermediate strain and strain B/HK/5/72). Virus-infected cell preparations on glass slides and standard immunofluorescent techniques were used essentially as described previously.³ Preparations were examined under a Tiyoda R1 tr microscope, using a 2 mm OX1 exciter filter and an 805Q barrier filter with mercury vapour illumination. A significant feature of the responses detected is that the IgG titre was rather similar against all three strains; any differences in the titres were fourfold or less (table I). Preliminary work done with paired sera from an outbreak due to influenza B/HK/5/72 virus gave results of a similar nature. This points to the possibility that infection due to one strain could result in some immunity to all three.

TABLE I—IgG Fluorescent Antibody Response to Influenza B in Children Aged 11-15 Years Infected with the Intermediate Strain of Influenza B Virus

Patient	Virus Strain		
	Old Influenza B*	Intermediate Influenza B	Influenza B/HK/5/72
1	7.5/160†	5/240	<5/60
2	<5/320	<5/320	<5/80
3	7.5/30	<5/60	<5/30
4	15/240	10/960	<5/640
5	5/120	15/160	5/80
6	<5/120	<5/80	<5/40

*B/England/32/71.

†Acute/convalescent serum specimens (taken 18 days apart).

TABLE II—IgG Fluorescent Antibodies to Influenza B Virus (Old Strain) in Manchester Sera, 1971

Age (years)	Percentage with Antibody*
<6/12	57
≥6/12	21 } 43
1-4	34
5-10	70
11-20	97
21-35	100
>35	89

*Sera screened at 1/10.

Fluorescent antibody studies done in 1971 on a random collection of Manchester sera⁴ are of interest in this context. If the presence of IgG fluorescent antibody to influenza B virus is taken as an index of immunity to influenza B, the results (table II) indicate that immunity should be maximal in the adult age groups, whereas a proportion of individuals less than 20 years of age should be susceptible to influenza B infection. The known epidemiological features of the disease support this hypothesis in that influenza B infections do predominantly affect the younger age groups. The epidemiological behaviour of infections due to the two new variant strains, which last winter left older age groups relatively unaffected, could also, on the basis of these figures, point to the presence of cross-immunity between the three strains of influenza B, as postulated above. Studies are now under way to assess further the usefulness of the fluorescent antibody technique in predicting immunity or susceptibility to

the various strains of influenza A and B by determining the fluorescent antibody titres to the different influenza strains in selected sera.—We are, etc.,

L. M. DE SILVA
P. R. MORTIMER
D. P. CASEMORE

Public Health Laboratory,
General Hospital,
Middlesbrough

¹ *British Medical Journal*, 1974, 2, 511.

² *British Medical Journal*, 1974, 2, 620.

³ de Silva, L. M., et al., *Journal of Hygiene*, 1973, 71, 107.

⁴ Tobin, J. O'H., and de Silva, L. M. Unpublished observations.

Vaccination against Whooping Cough

SIR,—Your leading article (31 August, p. 539) comprehensively analysed the difficulties in assessing vaccination against whooping cough. There may perhaps be some over emphasis at the beginning on laboratory methods of assessing the effectiveness of vaccination against infectious disease. Unless the specific protective property conferred by the vaccine against the particular infectious disease can be isolated and accurately assayed laboratory estimations offer only presumptive evidence.

The only valid assessment of routine immunization is the effect it produces in routine use. This, as you rightly point out, is exceptionally difficult to determine because the epidemiology of the disease is being modified simultaneously by many factors—social, nutritional, cultural, and therapeutic, to name only some—as well as by the immunization procedure itself. One of the more striking changes in the epidemiology of pertussis is the raising of the age of attack, and this has certainly contributed considerably to the reduction in death rate. There is evidence to suggest that routine vaccination has been an important factor here.

Laboratory tests have another disadvantage in that they imply that the production of absolute protection is necessary for the vaccine to succeed in practice. When a vaccine is used widely less than 100% effectiveness may well achieve the desired result. The production of undesirable side effects is not only unfortunate for the children who exhibit them but is also of importance in reducing acceptability. It may be more important to produce a vaccine which is acceptable but less effective than a vaccine which is 100% effective but non-acceptable to some parents because of side effects. The use of a two-dose schedule of the absorbed vaccine at a reduced strength could be as effective as a more frequent administration at a higher dosage, especially if it achieved a wider use because of a higher acceptability.—I am, etc.,

J. S. DODGE

University of Otago,
Dunedin,
New Zealand

Respiratory Distress Syndrome in the Newborn

SIR,—Your leading article on the present more hopeful position for the premature infant with respiratory distress syndrome (R.D.S.) (23 November, p. 428) rightly emphasizes the results that can be achieved with modern intensive care. It also raises

wider issues concerning the care of the sick newborn.

Your suggestion that "each N.H.S. area should surely come to an agreement that women in premature labour should be transferred for delivery to a unit fully equipped to deal with R.D.S. of all degrees of severity; infants with R.D.S. do not travel well" is a sensible one, and if one could be assured that not too many of these infants would be born in the ambulance en route, possibly even a practical one. It would ensure that infants not ill enough to require intensive care could return with their mothers, whereas those requiring it would have immediate intensive care available. On the other hand, your suggestion that "what is required is a technique which can be set up quickly and easily in the special care unit of any maternity hospital" may be a hopeful ideal for the future but at present is probably no more than pie in the sky.

At our current state of knowledge, treatment of severe R.D.S. still demands, as do other neonatal illnesses, the provision of medical, nursing, and laboratory vigilance of a high order, available at all times of the day and night. In addition, trained resident staff have to be available at all times to collect infants in distress in outlying hospitals. Infants with R.D.S. may not travel well, but with expert supervision en route they do at least seem to arrive safely.

Results from our own neonatal intensive care unit, and others similar, fully justify the efforts expended. It is surely unrealistic, however, to suppose that a service of this order can exist, or indeed need exist, wherever infants are born, and centralization deserves the highest priority at this time of reorganization of the Health Service.

Recent developments, such as the introduction of continuous positive airways pressure (C.P.A.P.) and the exposure to acute neonatal paediatrics which most junior staff now receive, mean that many infants can be satisfactorily managed in local hospitals. This is to be encouraged. It reinforces the mother-baby link and will improve the overall standards of practice. However, there is no doubt that some infants are too sick to be managed anywhere other than at a regional centre, and it is in their best interests to be transferred as early as possible. The difficulty, which frequently arises, is how to decide which infant should be transferred.

Professor K. W. Cross¹ has highlighted the dilemma of oxygen therapy in the newborn and pointed out that avoidance of oxygen damage to the eyes may have resulted in many infant deaths from hypoxia. The crucial deciding factor in the management of severe R.D.S. is probably whether or not the hospital concerned can provide an immediate service for blood gas monitoring throughout the 24 hours and has the necessary trained staff to deal with life-threatening emergencies such as pneumothorax, which so often occur.

The indications for transfer of an infant to a central unit thus appear to be threefold: (a) the need to ventilate the infant because of the severity of its disease or apnoeic attacks, (b) inability to manage the infant on C.P.A.P., or (c) lack of facilities for round-the-clock monitoring of arterial oxygen in babies who need it. If such a policy were adopted on a regional basis only a small proportion of infants with respiratory distress syndrome would need to leave their mothers and the specialized units would be