venously per day. We are continuing to investigate the underlying mechanisms to develop a convenient and completely effective prophylactic regimen and to assess the role of other types of parenteral nutrition in the genesis of this complication.

C A J WARDROP
R V HEATLEY
R H P WILLIAMS

M H LEWIS
G B TENNANT
L E HUGHES

Welsh National School of Medicine,
Cardiff

¹ International Society of Haematologists' Meeting, London, August 1975.

² Wardrop, C A J, *et al*, *Lancet*, 1975, 2, 640.

³ Chanarin, I, and Davey, D A, *British Journal of Haematology*, 1964, 10, 314.

Shoulder pain from subluxation in the hemiplegic

SIR,—In a study of shoulder pain in hemiplegia¹ seven other senior physiotherapists and myself found it to be a major barrier to progress in rehabilitation. In your leading article on shoulder pain from subluxation in the hemiplegic (14 June, p 581) you stated that supporting the arm in a sling made the patient more comfortable.

We designed several slings or supports for trial in each of our major hospitals. These increased sensory deprivation in the arm, did not solve the bed problems, and, from observation, had little effect on pain once it had started. The trials did not work because none of us could get total co-operation in keeping the appliance on. Some designs were discarded because nurses could not apply them. We tried to keep records of patients who could be put into a support on admission to casualty. But this scheme also failed as no hospital would run it—when life was perhaps at stake who could be bothered? Furthermore, the damaging traction could well have already happened when the family lifted the patient from the floor or into an ambulance. Many patients are treated at home and do not get to a physiotherapy or rehabilitation department till later in their illness. We also questioned the use of slings over a period as patients seemed to become addicted to them after they became unnecessary. And, lastly, wearing a support when walking, though good for the subluxed shoulder, adversely affected the gait pattern and, once formed, it could not be changed.

The practical problems of management are many. The doctor who orders a sling may be prescribing a treatment which cannot or will not be satisfactorily carried out. However, if doctors are more aware of the problems and try to tackle them from the start better solutions may be found.

J MILLER
Physiotherapist-in-charge

The Geelong Hospital,
Geelong, Australia

¹ Spencer, K, *Australian Journal of Physiotherapy*, 1974, 20, 4.

Low-dose heparin and the prevention of venous thromboembolic disease

SIR,—I would like to congratulate you on the leading article (23 August, p 447) on low-dose heparin and the prevention of venous thromboembolic disease. This presented a balanced view of the present state of prophylaxis in this most important problem. However, the following two articles

were disappointing and indeed misleading.^{1,2}

A plea for standardisation of the radioactive fibrinogen uptake test is a worthy aim. The most important components of the test are the quality of the fibrinogen and the equipment used. There is considerable variation in different batches of radioactive fibrinogen. When partially degraded fibrinogen with a high free radioactive iodine content (more than 5%) is used, bizarre counts are obtained over the limbs during the first 72 hours. This is probably due to the fact that circulating free iodine is rapidly excreted and gives low counts on subsequent days. As regards the equipment, when dealing with low counts the ratemeter is less sensitive and errors in interpretation of results are commonly seen. If the precautions outlined by Kakkar *et al*³ are followed, then such mistakes are eliminated. It is essential that after the counts in one leg have been completed the radioactivity over the reference point (heart) is again recorded to check the baseline before the second limb is scanned. This reduces the effect of "drift" in the machine. I have never observed the pattern of leg counts shown in table II of the article under discussion.¹

In the second article² the claim is made that intermittent calf compression results in a reduction of 90% in the incidence of deep vein thrombosis. It follows from this observation, therefore, that one could hope at the very most for a 10% improvement. Under such circumstances the all-important consideration is the number of patients in the trial. To be confident of showing a 10% improvement one would require about 1000 patients in the study.⁴ With 85 patients the chance of obtaining a significant result is that of the toss of a coin. No conclusions from the study can be reached.

T P CORRIGAN

Mater Misericordiae Hospital,
Dublin

¹ Roberts, V C, *British Medical Journal*, 1975, 3, 455.

² Roberts, V C, and Cotton, L T, *British Medical Journal*, 1975, 3, 458.

³ Kakkar, V V, *et al*, *Lancet*, 1970, 1, 540.

⁴ Boag, J W, *et al*, *British Journal of Radiology*, 1971, 44, 122.

Deaths in asthma

SIR,—A 46-year-old known asthmatic but otherwise healthy woman was admitted to hospital after a severe episode of asthma. She had been at a night club and had had a few drinks when the attack started. With the help of a friend, she retired to the cloakroom and was seen to take repeated puffs from her isoprenaline inhaler. On leaving the cloakroom she collapsed to the floor and made further efforts to use the inhaler. An ambulance was called but by the time of arrival in hospital she was in asystole. Despite resuscitation the patient died without recovering consciousness. Identification of propellant gases in a blood sample taken some hours after admission indicated that a substantial amount of propellant gas had been inhaled and confirmed excessive use of the inhaler. At necropsy no other underlying condition was found.

The epidemic of asthma deaths in the 1960s coincided with the increased use of pressurised aerosols.¹ The dangers associated with their overuse have become well recognised.² Selective sympathomi-

metic drugs are now available and we wondered whether our patient would have been better off using a bronchodilator with selective B₂ adrenergic stimulant effects rather than isoprenaline.

The Committee on Safety of Drugs informed us that since 1970 it has had reports of 31 asthmatic deaths among patients using isoprenaline inhalers and of a similar number of deaths (31) in patients using salbutamol inhalers. All the deaths were due to cardio-respiratory failure with underlying ventricular arrhythmias. Although in recent years selective B₂ bronchodilators may have been used in greater quantities than isoprenaline, it is also likely that abuse of either bronchodilator can have serious consequences. Another question raised by this case is whether the cardiostimulant effects of bronchodilators are potentiated in subjects drinking alcohol.

W N DODDS

N G SOLER

H THOMPSON

General Hospital,
Birmingham

¹ Inman, H W, and Adelstein, A M, *Lancet*, 1969, 2, 279.

² Committee on Safety of Drugs, *Aerosols in Asthma*, Adverse Reactions Series, 1967, No 5.

Management of acute asthma

SIR,—Your leading article under this title (11 October, p 65) was in fact concerned with *persistent* severe asthma or status asthmaticus, and the diagnostic and therapeutic suggestions which you gave were therefore relevant only to management in hospital, to which you say all such patients should be sent. But acute asthma in the sense of a sudden worsening of chronic asthma or a severe attack in a mild asthmatic (from exposure to an allergen or bronchial irritant or coming out of the blue) must be treated *before* the patient is sent to hospital. It is when general practitioners take your advice and just send for an ambulance to take such patients to hospital that they too often arrive dead, tragically and quite unnecessarily.

No patient with acute asthma should be sent to hospital without first being given 200 mg of hydrocortisone or prednisolone intravenously or intramuscularly. You suggest that intravenous steroids take an hour to have any effect, but the study on which you base this opinion was based on changes in ventilatory tests such as peak flow, FEV₁, and PEF₁.¹ Symptomatic relief may appear much sooner than that—within as little as 15 minutes in my experience. This may be due to a reduction in total lung capacity which can relieve asthmatic distress without affecting ventilatory capacity.² I have seen considerable relief even from oral prednisolone within half an hour. Even if this relief of bronchospasm may be partly psychogenic owing to the patient's confidence in the value of steroids based on past experience, it is none the less real and may be life-saving.

C M FLETCHER

Royal Postgraduate Medical School,
Hammersmith Hospital,
London W12

¹ Ellul-Micallef, R, and Fenech, F F, *Thorax*, 1975, 30, 312.

² Woolcock, A J, and Read, J, *Australian Annals of Medicine*, 1966, 15, 196.