black market dealings, but it does not stop them. (2) Does it cure? In my humble opinion, No. And I doubt if it ever will. It does lead to trading of NHS drugs. (3) Is it a good thing for us to let them receive sick pay and supplementary benefits, and stay at home on their drugs? (4) These people are a potential hazard to others. They have an "infectious," almost incurable, and very dangerous disease which will probably cause many early deaths. (5) Anyone who spreads addiction, sorrow, early death, for personal gain is far worse than someone who kills in the heat of the moment. They merit the most severe penalty. It must be made impossible for them ever to do this again. (6) Other forms of treatment: Although "patients" they must be restricted by law and treated as compulsory inpatients until the severe withdrawal symptoms are past. They should then be sent to hostels, work camps, etc, where the regimen would have to be kind but strict, administered by a dedicated staff who could not be "bought." No contact with the outside world would be allowed, no parcels, no unopened letters, visitors very rarely, and strictly supervised, only, say, a quarter of the inmates being visited at any one time.

This re-education in the broadest sense would have to continue for quite a time, and when patients were allowed to go home a kindly but strict and frequent supervision would have to be kept over them until they were considered completely and permanently cured. I realise how difficult all this would prove, and how expensive to institute, but not much less would be effective. Dedicated people are hard to find nowadays and they would be likely to have to face vilification by folks of small understanding in the outside world and vituperation from those inside. The possibility of litigation could not be ruled out. They would need wholehearted and firm backing by the law. If real cures, permanent cures, resulted, perhaps this difficult and expensive treatment would prove worth while.

C R BARRINGTON

Bromley, Kent

Toxic megacolon in Crohn's disease

SIR,—Your leading article on the above subject (27 September, p 723) refers to a recent report from this institution.¹ Our preferred method of primary surgical treatment for toxic megacolon is that of loop ileostomy and decompression blow-hole colostomy.² In a group of 49 patients recently reported³ there was one death, a mortality of 2%. All patients in the group detoxified and decompressed following this procedure, the average hospital stay being 18 days. Two patients required early colectomy during the original hospitalisation for persistent colonic haemorrhage. Colectomy was subsequently performed on 41 of these patients at a mean interval of six months, and in eight patients an ileorectal anastomosis was able to be performed. Pathological examination of the colectomy specimens showed a 2:1 predominance of transmural colitis 66% to mucosal ulcerative colitis 29%, with 5% being in the non-specific group.

On the basis of these results we feel that less than colectomy is indicated in the surgical management of the toxic megacolon

phase of inflammatory bowel disease, and that total proctocolectomy can hardly ever be justified in this condition. It seems logical to us to perform a comparatively simple and effective operative procedure in an acutely ill patient, and in this way to prepare the patient for colectomy at a time when the acute illness has passed.

D G JAGELMAN
V W FAZIO
R B TURNBULL

Department of Colon and Rectal Surgery,
Cleveland Clinic Foundation,
Cleveland, Ohio, USA

- 1 Farmer, R G, Hawk, W A, and Turnbull, R B, *Gastroenterology*, 1975, **68**, 627.
- 2 Turnbull, R B, et al, *Surgical Clinics of North America*, 1970, **50**, 1151.
- 3 Fazio, V W, *Gastrointestinal Emergencies*. Philadelphia, W B Saunders. In press.

Serum α -fetoprotein in cystic fibrosis

SIR,—Professor R K Chandra and others (29 March, p 714) reported significantly increased levels of α -fetoprotein (AFP) in the serum of patients with cystic fibrosis (CF), in the parents of the patients, and in some of their siblings in Newfoundland, Canada. These findings were confined by Dr J A Smith (17 May, p 392) in the United States for both homozygote and heterozygote carriers of CF genes. In contrast to Professor Chandra, Dr D J H Brock and others in Scotland and Dr J C Wallwork and others in England (17 May, p 392) were unable to detect an increased serum concentration of AFP in any of their CF patients or their relatives. So far as we know, serum AFP levels have not been measured in homozygote and heterozygote carriers of CF genes in central Europe.

Over the past four months we have examined the sera of 38 patients (aged 2 to 18 years) with CF of various severity of the disease for the presence of raised serum levels of AFP. Except for one Turkish boy, all patients with CF were of German origin. The diagnosis of CF was based on both clinical symptoms of pulmonary and pancreas involvement and sweat sodium-chloride levels above 80 mmol/l on pilocarpine iontophoresis. All patients with CF except one 7-year-old boy had normal values of SGOT, SGPT, and the other liver-specific serum enzymes.

The determination of AFP in the serum was carried out by radioimmunoassay with ¹²⁵I-labelled AFP in the double antibody technique. The first antibody was a rabbit anti-human AFP-serum, the second antibody a goat-anti-rabbit- γ -globulin. The test was carried out in 0.1 M borate buffer containing 0.5% beef albumin as described by Nishi and Hirai¹ and Ishii.² Incubation periods were always 24 hours for the first antibody, the radioactive labelled AFP, and the second antibody. ¹²⁵I-labelled AFP and the antisera were obtained from Dianabot-Laboratories, Tokyo, Japan. Crystalline AFP, isolated as described by Lehmann,³ 4 was used as standard by Mancini's radial immunodiffusion technique. The concentration of crystalline AFP (local standard) was 90 μ g/ml in the International Standard Preparation 72/225 of the World Health Organisation (Lyon, France).⁷ The normal range was calculated from the values of 62 healthy adults: the mean value of AFP in the serum was 4.52 ng/ml, the standard deviation 1.92 ng/ml, the upper limit of the 2.57- σ -range, including 99% of all values, 9.45 ng/ml.

The serum AFP levels in our 38 patients with CF were not different from those in healthy adults. The average serum AFP level of all patients with CF was 3.08 ng/ml (range 1-10 ng/ml). The standard deviation was 2.27 ng/ml. In infants and children the

normal range of the serum AFP levels is even higher than in healthy adults. Therefore the serum AFP levels in our patients with CF are within the normal range for healthy children of the same age. In addition, the sera of all 38 patients with CF were examined both by electroimmuno-osmophoresis with anti-human AFP-serum (limit of sensitivity 3.6 μ g/ml) and by double diffusion (limit of sensitivity 10 μ g/ml) as described by Lehmann and Lehmann.⁵ Our findings correspond with those of Dr Wallwork and others and of Dr Brock and others. These authors also, in contrast to both Professor Chandra and others and Dr Smith, have been unable to detect an increased serum concentration of AFP in any of the CF patients.

There is no explanation for this discrepancy, as Professor Chandra and his colleagues have apparently used the same rabbit antiserum and standards as Dr Brock and others. In our opinion, genetic differences cannot explain the different results concerning the serum AFP levels in patients with CF. The majority of the ancestors of the patients with CF who have been tested in the USA and in Canada may have their origin in central Europe and the north-western part of Europe.

G KNÖFFLE
H W ROTTHAUWE

University Kinderklinik,
Bonn, West Germany

F G LEHMANN

University Medical Klinik,
Marburg, West Germany

- 1 Nishi, S, and Hirai, H, *Cancer Research*, 1973, **14**, 79.
- 2 Ishii, M, *Cancer Research*, 1973, **14**, 89.
- 3 Lehmann, F-G, Lehmann, D, and Martini, G A, *Clinica Chimica Acta*, 1971, **33**, 197.
- 4 Lehmann, F-G, and Lehmann D, *Zeitschrift für klinische Chemie und klinische Biochemie*, 1971, **9**, 309.
- 5 Mancini, G, et al, *Colloquium on the Protides of the Biological Fluids*, 1965, **11**, 370.
- 6 Lehmann, F-G, and Lehmann, D, *Zeitschrift für klinische Chemie und klinische Biochemie*, 1973 **11**, 399.
- 7 Sizaret, P, et al, *Journal of Biological Standards*, 1975, **3**, 201.

Whooping-cough vaccine

SIR,—The statement by the Joint Committee on Vaccination and Immunisation (20 September, p 687) concerning the continued use of whooping-cough vaccine is important, if only for the data it contains. These show that as in previous epidemics mortality from whooping cough in the 1973-4 epidemic occurred almost entirely in infants under 12 months. Current DHSS inoculation policy leaves this age group relatively or completely unprotected by direct vaccination, and any benefits of the inoculation programme appear to depend on herd immunity promoted among children in whom mortality from pertussis has always been much less. The success of this recently modified inoculation policy operating in the context of an increasing proportion of parents born after systematic pertussis inoculation was introduced, and who probably do not now have significant protection against pertussis,¹ remains to be seen.

It is not possible to judge from the committee's data the extent to which the promotion of community protection may be offset by the hazard from a severe encephalopathic reaction in an inoculated child past the age of maximum hazard from the natural

disease. Whatever the extent of the risk (and it is not clear what the committee considers to be a "low" risk), in my view there is a very strong case for compensating the parents of those children who are handicapped by such a reaction.

Although evidence has been presented on behalf of the Association of Parents of Vaccine Damaged Children to the Royal Commission on Civil Liability and Compensation for Personal Injury, it is unlikely that legislation will ensue for some time, and at best it is unlikely to be retroactive. The Joint Committee on Vaccination would do well to back its benign reassurance about pertussis vaccine with strong support for the speedy introduction of a State compensation scheme.

JOHN WILSON

Hospital for Sick Children,
Great Ormond Street,
London WC1

¹ Lambert, H J, *Public Health Reports*, 1965, 90, 365.

Volunteers and the aftermath of stroke

SIR,—The setting up of volunteer schemes for home visitors for dysphasic patients, as described in your leading article (13 September, p 606) by Miss Valerie Eaton Griffith (p 633), is surely to be applauded in principle, but why do Miss Griffith and your correspondents (27 September, p 763) urge that such schemes should develop outside the existing speech therapy services? We see three dangers in setting up a separate autonomous system.

(1) Despite the present declarations from Miss Griffith and the Chest and Heart Association that volunteer schemes and professional speech therapy services can co-exist in parallel, economy-minded area health authorities may be tempted to jump to the conclusion that volunteer schemes can provide a cheap substitute for professional services for dysphasic patients. The glowing account that 21 out of 31 patients improved in speech after a few months of visiting by untrained and unpaid volunteers calls into question, if read uncritically, the value of the existing speech therapy services. It is true that, on several counts, the report does not stand up to critical examination—the patients were selected, the assessment procedures were not standardised, there was no untreated control group, some of the patients were also receiving professional speech therapy, and the evaluations of progress were largely made by people who had undertaken the therapy and people who were firmly committed to the scheme's success. But such reservations tend to get lost from view when schemes seem to offer solutions to difficult problems at little cost.

(2) Dysphasia is a complex subject about which continuing research is constantly providing new insights, particularly at the present time when a new discipline of "neurolinguistics" is developing; and surely it is not unreasonable to believe that the management of language therapy for dysphasic patients requires an awareness of this major area of study—requires, in fact, the professional training of a speech pathologist and therapist. Yet Miss Griffith suggests that the secret of the success of the volunteers is their very ignorance of this study and insight. One thing of which the professional speech therapist is aware is that the type of general stimulation therapy which volunteer schemes are offering may well be quite unsuitable for people with some kinds of dysphasic problems and could even be harmful. Speech therapists know all too well that they do not possess all the answers; in the present state of knowledge about dysphasia patients should be carefully assessed by modern techniques in order that therapeutic regimens can be individually designed for each patient, and

reassessed, so that these regimens can be revised as therapy proceeds.

(3) There is a continuing need for the systematic collection and collation of data so that patterns of impairment can be compared with patterns of treatment and recovery. There is also an urgent practical need for scientifically based research into the effectiveness of all ways of coping with the rehabilitation of dysphasic patients. No one, surely, would advocate that this research should be undertaken by the untrained rather than by qualified professionals, but isn't this what has been happening?

These seem to us to be compelling reasons why volunteer schemes should be fostered within the aegis of the professional speech therapy service rather than outside. If all those good intentions and hours of devoted work from volunteers are to be well directed, and if such schemes are to realise their potential usefulness, doctors who wish to refer patients to voluntary organisations should do so through a speech therapist, who can decide on whether this approach is suitable for the individual patient, plan a therapeutic programme, and supervise its implementation. If there is any magic in being untrained and uninformed, as Miss Griffith suggests (and we suspect there isn't), it needs to be complemented by informed guidance.

These views are representative of those expressed by senior colleagues in speech therapy and provide some indication of the general concern which is felt in relation to the article.

MARGARET EDWARDS

Department of Speech,
University of Newcastle upon Tyne

Sudden fetal deaths

SIR,—With the increasing use of continuous fetal heart monitoring there have been a number of anecdotal reports of sudden fetal death without warning. It is extremely important that the histories and fetal heart traces of these cases should be studied in detail if we are to make labour safe for the fetus.

I would be most grateful if any of your readers who have experience of such a case would write to me and, if possible, send a short clinical summary and the fetal heart trace. Hopefully, if sufficient cases are collected it will prove possible to make recommendations that will help to avoid these distressing unexpected intrapartum stillbirths.

R W BEARD

Department of Obstetrics and Gynaecology,
St Mary's Hospital Medical School,
London W2 1PG

Epilepsy

SIR,—I read with interest Dr F B Gibberd's article (1 November, p 270) and would like to make the following comments.

(1) It is surprising in November 1975 to find an article on the treatment of epilepsy which does not mention sodium valproate. (2) In young children presenting with convulsions it is important to consider the possibility of non-accidental injury, especially when fundal haemorrhages are seen. (3) The minor motor epilepsies, dismissed in two lines, are more common than petit mal and their treatment is different; it may include

clonazepam, nitrazepam, sodium valproate, steroids, and, in resistant and severe cases, a ketogenic diet. (4) With the availability of sodium valproate in addition to ethosuximide there must now be little place for truxidone in the treatment of petit mal. (5) The prophylaxis of febrile convulsions is an open subject; the evidence as regards regular anticonvulsant prophylaxis is contradictory and there is little evidence that intermittent phenobarbitone is useful.

D P ADDY

Department of Paediatrics,
Dudley Road Hospital,
Birmingham

SIR,—Although not mentioned in Dr F B Gibberd's article on the treatment of epilepsy (1 November, p 270), a neonatal withdrawal syndrome has been associated with barbiturate use in pregnancy.^{1,3} The prevention of this syndrome might be another valid reason for stopping treatment before pregnancy in attack-free patients.

Because the age of onset of symptoms ranges from 30 min to 14 days¹ this condition should be borne in mind both during hospital stay and after discharge in any infant of an epileptic mother.

I BLUMENTHAL

Department of Child Health,
University of Aberdeen

- ¹ Desmond, M M, *et al*, *Journal of Pediatrics*, 1972, 80, 190.
- ² Bleyer, W A, and Marshall, R E, *Journal of the American Medical Association*, 1972, 221, 185.
- ³ Martinez, G, and Synder, R D, *Neurology*, 1973, 23, 381.

Infective agent in infantile gastroenteritis

SIR,—Dr B Rowe and Mr R J Gross (18 October, p 162) are right to take up the cudgels in defence of some strains of *Escherichia coli* as causes of infantile enteritis and to remind us of the difficulties and limitations of routine bacteriological methods. Yet much of the evidence for the enteropathogenicity of *E coli* strains for man rests on similar observations of association with disease as have been reported for some of the viruses. A combined bacteriological and virological approach is required to define and distinguish the aetiological roles of these bacteria and viruses, together or separately, with due attention to the possibility of spurious associations. Difficulties of proof, particularly in neonates, may necessitate ultimately accepting such circumstantial evidence as the basis of guilt, however.

If alteration of the intestinal contents by viral diarrhoea favours selection of certain *E coli* types rather than others these types will show a secondary, not causal, association with the diarrhoea. If some of the viruses are coliphages they likewise may show spurious association with diarrhoea, unless by analogy with diphtheria the phages confer pathogenicity on the bacteria. If either bacterium or virus causes diarrhoea in a host carrying the other type of organism the diarrhoeal condition will favour simultaneous dissemination of both agents so that both may show association with diarrhoea in a particular outbreak. Similarly both agents, or whichever of them the investigator was equipped to detect, might show association with diarrhoea in an outbreak due to neither but to a third and undetected agent.