required to be united with complement in order to be an antidote, and that the complement was "fixed" by the toxin. If so, all that was required was to supply complement. I performed three experiments:

1. I put a fly blister on some healthy persons, and injected normal human serum into yaws cases. No result followed.

2. I heated the serum from cases that were being cured after "606" injections, and the serum, *even after boiling*, was efficacious.

3. I put blisters on three cases of yaws, and used the patients' own serum for their injection. No result.

I have therefore abandoned the provisional hypothesis of fixation of the complement by the toxin.

I also performed the following control experiments: I injected sterilized water with tincture of cantharides, and satisfied myself that the presence of cantharides in the serum has no effect whatever.

I injected four cases with cacodylate of soda for comparison, and I see nothing to compare with the experiments with "606" and the serum.

A further experiment with normal human serum obtained by a blister was negative.

It is my intention to inject a goat with "606," and use the goat's milk for young children with yaws. I will also blister cases that have been cured of yaws under ordinary treatment and experiment with their serum. If it be possible to cultivate spirochaetes, I suggest that spirochaetes be injected into a horse; that this be followed by "606" injections, and that experiments be made with the serum in cases of syphilis.

When the experiments were carried on, no other treatment of any kind was used; even a bath was not permitted. No complications, local or general, occurred. I believe that the dose of "606" for yaws can be safely increased, as the reaction was very slight, and in some cases absent. Cases will require a second injection. The pain after the injection of the emulsion lasts for about six hours, but is not very great. Dr. Rost intends to publish the cases injected by him with "606." We all owe him a debt of gratitude for so willingly using his "606" tubes in yaws cases. The blisters were applied on the fourth day after injection with "606," or with the serum of a "606" case.

In the light of marked effects in the secondary serum cases we may have to change the hypothesis of the formation of an antibody by the injections. The curative substance seems to have the power of increasing itself. Is it a ferment?

A REPORT ON THREE CASES TREATED WITH EHRlich's SPECIFIC FOR SYPHILIS.

BY

E. G. FFRENCH, M.D., F.R.C.S.ED., CAPTAIN R.A.M.C.,
STATION HOSPITAL, AHMEDNAGAR, SOUTHERN INDIA.

THROUGH the kindness of Mr. J. E. R. McDonagh of the London Lock Hospital, who spared me a few tubes of "606," I have been able to carry out Ehrlich's treatment in three selected cases of syphilis. I had the good fortune to see Mr. McDonagh treat some patients in London, and I have endeavoured to treat the cases after his method. The doses were received in the form of a yellow powder hermetically sealed in glass ampullae, the dose ranging from 0.3 to 0.9 gram. One end of the glass ampulla is broken off, and the contents turned out into a glass mortar which has been sterilized. The powder is dissolved first of all in as small a quantity of ethyl alcohol as possible. To this is added about 10 c.cm. of very hot water, and the fluids stirred vigorously until every trace has been dissolved; then a small quantity of solution of sodium hydrate (40 grams to the litre) is added, well mixed, and three drops of a $\frac{1}{2}$ per cent. alcoholic solution of phenolphthalein added, and titrated with normal acetic acid until the red colour has completely disappeared. Finally, a few drops of normal sodium hydrate are added until a faint rose tint comes back to remain, so that the solution is just alkaline. The solution is now ready for injection. A fairly large syringe capable of holding 12 c.cm. and fitted with a long needle is used. It is necessary to have a needle with a large bore, as

sometimes the fluid is apt to become viscid, and would certainly block an ordinary needle.

The urine must be tested in every case before injection and daily afterwards. The strictest antiseptic precautions should be taken and the needle boiled in salad oil to prevent the blunting which occurs with water.

The scapular region is a suitable site for injection. The skin over the scapular muscles is well washed with soap and water, nothing else being used. The contents of the mortar are drawn into the syringe, and the needle plunged deeply into the muscles of the scapula on the side selected. The fluid is then slowly but firmly driven into the muscles. The needle is then slowly withdrawn, and a piece of aseptic gauze held by rubber plaster put over the puncture. The patient should be kept in bed for five or six days.

CASE I.

Colour-Sergeant G., aged 35 years, acquired syphilis eleven years ago. A very short time afterwards he had seventy-two days' treatment with hyd. c. cret. in hospital. He did not have any further treatment, and kept fit until his tonsils and palate became ulcerated six weeks ago.

On admission to hospital he was found to be suffering from a large perforation of the soft palate and extensive ulceration of the tonsils and posterior pharynx. The uvula had already ulcerated away.

Five days afterwards 0.5 gram of "606" was injected into the muscles of the right scapular region. He did not have any rise of temperature, and felt no ill effects, with the exception of some slight soreness at the sight of injection.

Two days afterwards there was a distinct improvement in the appearance of the ulcers. The sloughs had disappeared, and the perforation looked smaller. At the end of the first week the tonsils and posterior pharynx had healed completely and the perforation was about one-third the original size. He was weighed, and found to have gained 5 lb. About this time a small amount of lymphoid tissue was noticed at the site of the uvula.

A fortnight afterwards the hole in the palate had almost closed, and the patient's general condition was greatly improved.

Four weeks afterwards the perforation had closed completely, and the lymphoid tissue at the site of the uvula had grown into a fairly large process. He had regained his normal voice, and on being weighed was found to have gained 16 lb.

CASE II.

Lance-Corporal R., aged 32 years, acquired syphilis five years ago. He did not receive any mercurial treatment internally, as the sore was diagnosed to be soft. He kept fit until May, 1910, when he was admitted to hospital suffering from severe ulceration of the upper gums, tonsils, and posterior pharynx. He had also disease of the nasal bone on the left side and well-marked tibial pains at night. He was given several courses of inunction periodically, but his condition did not improve.

He was sent to me as a selected case for "606" treatment. I found that he was suffering from the above-mentioned symptoms, and that quite recently he had lost five upper teeth on the right side. A few days afterwards 0.5 gram was injected into the muscles of the right scapular region. He had no pain afterwards, but had a rise of temperature for three days, the highest being 101°.

Two days afterwards the tibial pains ceased and the condition of the gums and throat was markedly improved. At the end of the first week the throat had completely healed and the gums were almost well, the teeth being quite firm. His weight had increased 2½ lb.

A fortnight afterwards the gums and throat were quite healed; a piece of necrosed bone came away from the left nostril, and the discharge, which was profuse before, was greatly diminished.

On the eighteenth day he had gained over 1 st. in weight and looked quite fit. He is now quite free from any lesions.

CASE III.

Private L., aged 28 years, acquired syphilis in October, 1910. He was admitted to hospital at Poona soon afterwards and developed a profuse papular rash over the face, neck, trunk, and upper and lower limbs. He was given three courses of ung. hydrarg., and was transferred here for "606" treatment, as his condition was not improving. On admission, he had the rash over the face, trunk, and limbs, some ulceration of both tonsils, and well-marked enlargement of the cervical and inguinal glands.

He was injected in the right scapular muscles with 0.5 gram of "606." He suffered slight pain in the muscles for about two hours afterwards. Two days afterwards the enlarged glands had completely disappeared, and the rash was fading rapidly from the face. The condition of the throat showed marked improvement. He had a rise of temperature for three days. At the end of a week the rash had completely disappeared from the face, the throat was quite well, and the rash was fading from the trunk and limbs. He had gained 1½ lb. in weight. Eighteen days afterwards, with the exception of a few stains on the skin the result of the rash, there was nothing to show that he had ever had syphilis. He had gained 3½ lb. in weight.

The cases described are the first to be treated in India with Ehrlich's wonderful specific.