March of this year he had no fits, and was then discharged as recovered.

The following case I saw in private with Dr. McMillan of Shotts, and to him I am indebted for the history.

A. B., a female aged 21, began to have fits when she was 17, and they were becoming more severe and very frequent in spite of treatment with large doses of bromide. In December, 1920, she was put on bromide and borax, and has had no fits since. For the past three months she has taken no medicine, and has had no recurrence. This patient commenced with small doses of borax and bromide, which were gradually increased, and the combination which brought about the good result in her case was borax purificatus $7\frac{1}{2}$ grains, potassium bromide 5 grains, and Fowler's solution 2 minims, three times a day.

The treatment of the eighteen patients mentioned has been as follows. To begin with they were put on potassium bromide 10 grains, borax purificatus 5 grains, and Fowler's solution 2 minims, three times a day. The dose was gradually increased, and I found I obtained the best results with potassium bromide 15 grains, borax purificatus $7\frac{1}{2}$ grains, Fowler's solution 2 minims, three times daily, and this combination gave the results as stated in the above two groups. In addition the patients had a laxative at bedtime. At present I am trying different combinations of the two drugs from the above by way of experiment.

In conclusion, I may say that the continuous treatment has had no deleterious effect on the health of the patients, there have been no gastric or skin disturbances, and, in fact, the general physical condition has improved.

TRAUMA AND APPENDICITIS.

BY

CHARLES J. G. TAYLOR, M.A., M.D.,

SURGEON, NUNEATON AND DISTRICT GENERAL HOSPITAL.

The following two cases, which have recently been under my care, raise some interesting points in regard to the etiology of appendicitis, and on that account may be worthy of putting on record.

CASE I.

A. S., aged 18, was playing football on September 30th, 1922, when, towards the end of the second half, he was charged and received a severe blow on the right side of the abdomen from his opponent's elbow, which laid him out for the moment. He vomited, but was able to rise and walk to the dressing room. Whilst changing he vomited again. He then walked about half a mile to a bus, which conveyed him to within a short distance of his home. During the whole time pain in the abdomen was severe. When he arrived home he lay down, and as the pain got no better he, later in the evening, called in his doctor, who thought that he must be suffering from appendicitis, but in view of the history, decided to wait till the following morning. The diagnosis then seemed certain. When admitted to hospital at 2 p.m. the temperature was 100° , the pulse 88. The tongue was furred, the abdomen was definitely rigid and tender in the right iliac fossa, the point of maximum tenderness being somewhat below McBurney's point.

At 7 p.m. Dr. E. N. Nason administered an anaesthetic, and the abdomen was opened through a right rectus incision. Omentum presented on opening the peritoneal cavity. A small quantity of slightly turbid fluid was mopped out, and the appendix was then found pointing downwards over the brim of the pelvis, the vessels deeply engorged, and the organ throughout its length covered with patches of lymph. The appendix was removed in the usual manner and the abdomen closed without drainage.

Convalescence was uneventful, except that two days after operation a tense swelling the size of a plum was discovered in the tissues of the right spermatic cord immediately above the testicle. This was diagnosed as a haematoma, and at the time of discharge it had shrunk almost to vanishing point.

On cross-examination after operation this youth informed me that some two months previously he had suffered from an attack of "indigestion" (pain all over the abdomen and vomiting), which lasted for two days and then cleared up. He was positive that when he walked on to the football field he was feeling as well as he had ever done in his life.

CASE II.

H. V., aged 22, at about 1 a.m. on October 18th, 1922, was working on the night shift down the pit when he had to put forth a violent effort to lift a tub on to the rails. About three minutes later he was seized with sudden acute pain across the lower abdomen and vomited. He was brought out and sent home, when he vomited again three or four times, and the pain continued to be severe. At 9 a.m. he was seen by his doctor, who advised his immediate removal to hospital, but as "his people were not willing" he refused. However, at 6.31 p.m. on the following day he was admitted. The temperature was 99.8° and the pulse 88. He complained of severe pain in right side of the abdomen.

On examination there was tenderness all over the lower abdomen, with increased rigidity; tenderness was very marked over McBurney's point. An anaesthetic was administered by Dr. Bradbury, and at 8.45 p.m. the abdomen was opened by a right rectus incision, and a short, stumpy, acutely inflamed appendix, buried in adhesions, was found behind the caecum. About half a drachm of thin pus free in the peritoneal cavity was sponged out and the operation completed in the usual way, the abdomen being closed without drainage.

He had had a troublesome cough previous to the present illness, and this somewhat complicated his recovery, as he coughed open the skin wound. In other respects convalescence was uninterrupted.

In both of these cases the attack was definitely ushered in by violence—in the one by a direct blow on the abdomen, in the other by a severe strain. An intriguing point in both cases is their bearing on the Workmen's Compensation Act. In the first case, of course, that question did not arise; but supposing it had, I do not think that compensation could have been withheld, even allowing that the patient had had a previous attack, the onset being apparently so definitely the result of the blow. In the second case the influence of trauma is more debatable.

I find on looking through my notes that I have dealt with one other case allied to these. It occurred at Scapa Flow in 1916. A boy, aged 16, from H.M.S. *Conqueror*, was kicked on the right side of the abdomen. He was admitted to hospital ship two days later with tenderness and rigidity in the right iliac fossa, but the temperature was normal. I removed a very long and congested appendix. This case would probably have cleared up without operation.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

THE VARIATIONS OF NORMAL TEMPERATURE.

A PASSAGE in Dr. E. Clark-Jones's article (BRITISH MEDICAL JOURNAL, November 25th, 1922, p. 1026), that ignorance obtains as to what constitutes a normal temperature, leads me to record some investigations I made as a student on my own body temperatures in March and April, 1892. I found that thermometers differed by as much as a third of a degree, and selected one which recorded the mean of six as I was unable to procure one with a Kew certificate of accuracy.

I made 170 records of my mouth temperature at intervals of an hour or two during a period of twenty-one days in April, 1892, and, averaging each period, I obtained the accompanying Chart A. I was led to do this because in the previous

CHART A.—Normal day temperature curve while studying, with short periods of slight exercise.

CHART B.—Rise of temperature at five-minute intervals after severe exertion in well-trained condition.

March, during training for the Hospital Rugby Cup final, I had been trying to find out the effect of exercise on the temperature, with some astonishing results. I found that the effect of slight exercise was to raise the body temperature slightly for a short time until perspiration began, when, if the exercise was continued, there was a continuous fall, and a few days before the final cup tie after a two hours' run I found my mouth temperature was 94.6° ; Chart B shows the rise afterwards at successive five-minute intervals. At the commencement of vigorous training on March 6th that year after one hour's run I recorded 95.4° , and after half an hour's interval it had risen to 99.1° , and did not return to normal till four hours afterwards. The effect of training was to diminish that rise until it was not perceptible, although the fall was lower.

On March 8th, after one hour's run, I discovered that the urinary temperature was three degrees lower than the mouth temperature at the same time, being 93.7° while the mouth was 95.7° , and when I recorded the mouth temperature of 94.6° the urinary temperature was 99.6° .