J. N. WALTON: MUSCULAR DYSTROPHY

FIG. 1.—Focal muscle-fibre necrosis and phagocytosis in a female carrier of the Duchenne-type dystrophy. (H. and E. $\times 220$.)

FIG. 2.—An area of abortive regeneration in a female carrier; the small fibre to the left of centre is basophilic and contains several large vesicular nuclei with prominent nucleoli. (H. and E. $\times 220$.)

FIG. 3.—Electron micrograph, $\times 15,550$, transverse section of muscle obtained by biopsy from patient with muscular dystrophy of Duchenne type. There is widespread loss of myofibrils with disorganization of mitochondria and of the sarcotubular system. (From Pearce, 1964.) A=Relatively normal myofibrils. B=Area of myofibril loss. C=Mitochondria. D=Collagen fibrils outside the sarcolemma. E=Sarcolemma. F=Sarcotubule.

FIG. 4.—Electron micrograph, $\times 19,430$, of muscle section from patient with muscular dystrophy of Duchenne type, showing excessive numbers of bodies, provisionally identified as lysosomes, lying between the myofibrils. (From Pearce, 1964.) A=Myofibrils. B=Z-band. C=Sarcotubule. D=Lysosomes. E=Mitochondrion.

FIG. 5.—Normal myoblasts; adult mouse muscle; phase contrast. (From Pearce, 1964.)

FIG. 6.—A myoblast, presumed to be dystrophic; explant from a dystrophic mouse of the Bar Harbor strain; phase contrast. (From Pearce, 1964.)

the time. Twin C had a precordial systolic murmur which has not persisted, Twin D a cleft of soft and part of hard palate. At seven months Twin C had two left-sided fits associated with respiratory infections. Despite phenobarbitone, fits had recurred. Twin C, who survived separation, was estimated at 17 months to have a development level corresponding to 8 months. At a month under 3 years she weighs 28 lb. 12 oz. (13 kg.). She can sit unaided for a time, rocks back and forth, crows and gurgles, says *mum* and *dad*, can hold a toy in the right hand and takes some interest in her surroundings. She has a left spastic hemiplegia.

Pre-operative studies of behaviour showed that each twin reacted as a separate individual. One slept with the other awake or sucking a bottle or crying. Pulse, temperature, respiration, haemoglobin values differed. Electroencephalograms by Dr. G. Pampiglione revealed independence of cerebral activity, including induction of sleep and appropriate changes in the one given thiopentone by injection. That the other was unaffected suggested no marked arterial communication.

The post-operative course of both survivors was stormy and difficult. Skin coverage has been made complete, and in neither has there been any evidence of destruction of the grafted freeze-dried dura mater. A series of plastic helmets of increasing size has protected the brains from further damage.

Conclusion

The paediatrician's part in the management and treatment of craniopagus twins is described together with brief case records of two pairs of such twins. If the object of the endeavour is to separate pairs of conjoined twins, the team that has one survivor from each of two pairs has scored a real success. But if the object is regarded as restoring people who have met a strange fate to a normal happy family life, these cases are disappointing.

It is a pleasure to record the untiring work during the long months of Dr. Seymour Mason, Dr. Kenneth Hugh-Jones, and the house-physicians to the children's department at St. Bartholomew's Hospital, as well as the devotion of Sister Joan Clark, Miss B. Franklin, and the nursing staff.

REFERENCES

- Franklin, A. W. (1960). *Lancet*, **1**, 1065.
 — Tomkinson, J. S., and Williams, E. R. (1958). *Ibid.*, **2**, 683.
 Kenny, A. (1961). *Daily Express*, October 20.

Muscular Dystrophy: Some Recent Advances in Knowledge*

JOHN N. WALTON,† M.D., F.R.C.P.

[WITH SPECIAL PLATE]

Brit. med. J., 1964, **1**, 1344-1348

Genetic Research

Duchenne-type Muscular Dystrophy

Evidence in favour of a sex-linked or X-linked recessive inheritance of this variety of muscular dystrophy in the majority of families comes, first, from a study of typical pedigrees (Stevenson, 1953; Walton, 1955, 1956c); secondly, from the fact that affected males have been born of the same female but of different fathers (Walton, 1956c); thirdly, from the finding of a family in which there was a suggestion of crossing-over between this variety of dystrophy and red-green colour-blindness (Philip, Walton, and Smith, 1956); and, fourthly, from the fact that the disease has been described in typical form in a "female" suffering from Turner's syndrome (ovarian agenesis) who was presumed to have an XO chromosome constitution (Walton, 1956c). Although, as Boyer and Fainer (1963) point out, none of these findings is absolutely conclusive, nevertheless the evidence in support of sex-linked recessive inheritance of these cases would seem to be convincing even though no linkage has yet been demonstrated between this form of dystrophy and haemophilia or the newly described Xg blood group. With the collaboration of Dr. Race we have examined a number of families and have found no such linkage. An analysis of pedigrees has, however, amply confirmed that in a small proportion of families there are affected females as well as affected males; this and other evidence clearly indicates that the disease may sometimes be produced by an autosomal recessive gene (Lamy and de

Grouchy, 1954; Kloepper and Talley, 1958; Dubowitz, 1960; Jackson and Carey, 1961).

Chung and Morton (1959) give reasons for suggesting that, whereas a considerable number of cases of muscular dystrophy of the Duchenne type arise as a result of a gene mutation, there are others in which the disease may be truly sporadic—that is, it may be due to manifestation in the heterozygote. Hence there is evidence to suggest that there can be fundamental genetic heterogeneity among cases of clinically similar muscular dystrophy. As Nirenberg (1963) has pointed out, intracellular transmission of abnormally coded D.N.A. information occurs within the cell when R.N.A. carries the code into metabolic steps which terminate in a quantitative or qualitative alteration of a particular enzyme. Metabolic pathways associated with the expression of various specific genes often overlap and in turn may cause an overlap in clinical stigmata. We shall not be able to understand similarities and differences between neuromuscular entities which depend upon different genes until we understand the details of such overlap. Chung and Morton (1959) and Morton, Chung, and Peters (1963) have also given cogent reasons for suggesting that although many cases of limb-girdle muscular dystrophy are clearly the results of an autosomal recessive gene and the phenotype appears in the homozygote, there are others which are truly sporadic and are presumed to be due to expression in the heterozygote.

Of more direct practical importance are recent studies carried out on female carriers of the sex-linked recessive gene responsible for most cases of the Duchenne-type dystrophy. Aebi, Richterich, Colombo, and Rossi (1962), Hughes (1963), and Pearce, Pennington, and Walton (1964c) have found that over 70% of such female carriers can be identified by means of serum creatine kinase estimation. Dreyfus, Schapira,

* Conclusion of the Goulstonian Lecture delivered to the Royal College of Physicians of London on 20 January 1964. The first part was published last week.

† Neurologist, Regional Neurological Centre, Newcastle General Hospital; Physician in Neurology, Royal Victoria Infirmary, Newcastle upon Tyne.

Schapira, and Demos (1961) detected 23 out of 26 carriers using estimations of serum creatine kinase and measurements of circulation time. More recently still, van den Bosch (1963) and Barwick (1963) have indicated that minor electromyographic abnormalities suggestive of a myopathy may also be detected in carriers, and Emery (1963) and Pearson, Fowler, and Wright (1963) have demonstrated that there may be minor clinical abnormalities such as minimal muscular wasting and pseudohypertrophy of the calves in a number of carrier females who are apparently unaffected clinically (Fig. A).

FIG. A.—Pseudohypertrophy of the calves in the mother of two sons who were suffering from muscular dystrophy of the Duchenne type. Her serum creatine kinase was 54 units, amply confirming that she was a carrier.

Dubowitz (1963) and Pearce, Pearce, and Walton (1964) have found definite histological abnormalities in muscle biopsies obtained from a series of carrier females (Special Plate, Figs. 1 and 2). These observations all give considerable support to the Lyon hypothesis (Lyon, 1961, 1962) that in a proportion of cells of the female one X-chromosome may be inactivated. The evidence strongly suggests that in carrier females a proportion of the muscle cells are in fact dystrophic but that usually this proportion is not large enough to give rise to clinically obvious weakness. Nevertheless, the methods outlined are of considerable value in the detection of carriers, and hence are useful for the purpose of genetic counselling. It is likely that within the next year or two methods of carrier detection will become even more reliable.

Fundamental Research and the Pathogenesis of the Disease

It is my firm belief that an effective treatment for progressive muscular dystrophy will be discovered only when the fundamental biochemical defect or defects responsible for the muscular wasting which occurs as a result of the disease process are also discovered. Indeed, it seems that now we are able to detect preclinical cases of dystrophy, particularly of the Duchenne type, the correct time to attack the disease process would be at this stage before clinical weakness supervenes. I have not the time to describe fully modern methods

of research, but would like simply to mention briefly some of the techniques that are being utilized in the hope of determining the nature of the fundamental pathological process. The principal methods of attack are by means of the biochemical and histochemical study of muscle-biopsy samples; by investigation of the morphology of the disease process with the aid of the electron microscope and by means of tissue culture and interference microscopy; and, finally, by electrophysiological studies of muscle-fibre and membrane potentials. A considerable impetus to research has followed the discovery of forms of hereditary muscular dystrophy occurring in mice of the Bar Harbor strain (Michelson, Russell, and Harman, 1955) and in chickens (Julian and Asmundson, 1960), but although the disease processes in these animals are clearly dystrophic it is impossible as yet to say whether results obtained with such experimental material can necessarily be relied upon in studying the human disease.

Biochemistry

Since the early observation that patients with muscular dystrophy excreted excessive quantities of creatine in the urine (Levene and Kriszteller, 1909) a vast amount of biochemical work has been carried out and the concentration of many organic and inorganic compounds within dystrophic muscle has been compared with that in normal muscle. In the table I have summarized from Dreyfus and Schapira

Variations in Constituents of Muscle Tissue in Muscular Dystrophy*

Constituent	Change	Constituent	Change
Glycogen	Decrease	Respiratory enzymes :	
Lactic acid	"	Succinic dehydrogenase	Decrease
		" oxidase	"
Glycogenolytic enzymes :		Cytochrome "	"
Phosphorylase	"	Fumarase	"
Aldolase	"	Aconitase	"
Phosphoglucumutase	"	Cytochrome C	"
Hexokinase	"		
Lactic dehydrogenase	Increase	Protein compounds :	
Total glycogenolysis	Decrease	Collagen	Increase
Phosphorus compounds and		Myosin	Decrease
other organics :		Myoglobin	"
ATP	"	Transaminases	"
Creatine phosphokinase	"	Proteolytic enzymes	Increase
Hexose phosphate	Increase		
Phosphocreatine	Decrease	Inorganic compounds :	
Creatine	"	Potassium	Decrease
Carnosine	"	Sodium	Increase
		Chloride	"
		PO ₄	"
		Iron	Decrease

* From Pearson (1963).

(1962) and from Pearson ((1963) the biochemical abnormalities observed in this disease. Considerable work has also been carried out upon the nature of the myoglobin which is present in dystrophic muscle. Perkoff, Hill, and Tyler (1962) have found an abnormal pattern of myoglobin in childhood muscular dystrophy, as have Whorton, Hudgins, and Connors (1961), but the significance of these observations in relation to the nature of the dystrophic process remains obscure and other protein studies directed particularly towards estimations of actin and myosin concentrations in dystrophic muscle have been disappointing. My colleague Dr. Pennington has demonstrated an increased concentration of cathepsins, of acid phosphatase, and of 5-nucleotidase in dystrophic muscle, but the increased concentration of these enzymes may well be simply the result of tissue breakdown. Whereas it has been recognized for some considerable time that such enzymes as aldolase and creatine kinase leak from the diseased muscle cell into the serum, and that this leakage is presumed to be due to a defect in the muscle-fibre membrane, the concentration of these and other enzymes which behave similarly in dystrophic muscle is reduced to only about 80% of the normal. Pennington (1962), however, found

that adenylic deaminase was reduced to about a third of the normal value in both dystrophic mice and in human patients with muscular dystrophy of the Duchenne type. This is the most striking reduction in enzyme content of dystrophic muscle yet discovered, but its significance is obscure, as the function of this enzyme in muscle is not yet understood. McArdle (1963) has suggested that it is of importance to study phospholipids within muscle in view of the importance of these compounds in the formation and integrity of the muscle-fibre membrane; recently my colleague Dr. Pennington has begun certain studies of phospholipid synthesis in muscle using thin-layer chromatography and a technique whereby ^{14}C -labelled phosphorylcholine is incorporated into lipid in mouse skeletal muscle, but this work is still in its preliminary stages.

Also of considerable interest is a recent finding that in various mammalian species, including man, lactic dehydrogenase exists in five electrophoretically distinct molecular varieties in nearly all tissues, and that, in contrast to normal adult muscle which contains a preponderance of LD_4 and LD_5 isoenzymes, some patients with dystrophy lack LD_5 completely and have lesser quantities of LD_4 (Wieme and Herpol, 1962; Dreyfus, Demos, Schapira, and Schapira, 1962; Pearson, 1963). Further studies of this type will be awaited with considerable interest, and it seems plain that the increasing volume of biochemical research now being carried out is likely to be fruitful.

Histochemistry

Considerable interest was aroused by the suggestion of Bourne and Golarz (1959) that dystrophy might be due to a primary proliferation of connective tissue in dystrophic muscle; they demonstrated excessive activity of 5-nucleotidase in muscle biopsy samples. Subsequently, however (Golarz and Bourne, 1963), they have somewhat modified these earlier views, and others (Dubowitz and Pearce, 1961; Engel, 1962) have failed to confirm their findings. Whereas histochemistry has revealed that there are at least two types of muscle fibre in normal human muscle, one being rich in phosphorylase and the other rich in oxidative enzymes (Dubowitz and Pearce, 1961), there is no evidence that either of these fibre types is specifically involved by the dystrophic process. Recently, too, A. G. E. Pearce (1963) has made the interesting suggestion that monoamine oxidase, which is an important component of the sarcotubular system concerned with the conduction of the electrical impulse into the interior of the muscle cell, may be reduced in amount in cases of muscular dystrophy, but this is a preliminary observation which as yet requires very much further study.

Electron Microscopy

As yet ultrastructural studies of diseased muscle are in their infancy. Following upon the important work of Huxley (1957) and others, Price (1963) has confirmed that the views expressed about the ultrastructure of the muscle fibre which were based upon the examination of rabbit muscle can also be applied to man. At last the significance of the pattern of organization in normal skeletal muscle which the electron microscope has revealed is being clarified. To date studies of dystrophic muscle with this technique have not revealed any specific changes. Van Breemen (1960), Pearce (1963, 1964), and Mair (1963) have confirmed that disintegration and fragmentation of myofibrils and myofilaments may occur (Special Plate, Fig. 3), but neither the actin nor the myosin appears to be specifically involved and there is apparently no selective damage to any single part of the sarcomere. The mitochondria and sarcotubular elements are also damaged, but the changes are not very different from those observed

in patients with other forms of myopathy such as polymyositis and even steroid myopathy. On the other hand, Pearce (1963, 1964) has drawn particular attention to the excessive concentration of vesicular bodies, provisionally identified as lysosomes (Special Plate, Fig. 4), in dystrophic muscle. In collaboration with Pennington (unpublished observations) he has found that these bodies contain proteolytic enzymes such as cathepsins and acid phosphatase; they may simply be present in a scavenging role. It is to be hoped (Pearson, 1963) that electron microscopical studies of muscle obtained from preclinical cases of muscular dystrophy may help to identify the earliest structural changes which occur in this disease.

Tissue Culture

Geiger and Garvin (1957) first suggested that the growth pattern *in vitro* of muscle obtained from muscular-dystrophy patients was different from that of normal adult human muscle, but Herrmann (1961) was unable to confirm these findings and pointed to the difficulty of culturing adult human muscle. Pearce (1963, 1964) has described a technique of tissue culture applicable to adult normal mouse muscle and to dystrophic mouse muscle. He has found that on making small crush lesions in the muscle of such mice a typical histological picture of regeneration was seen after five days. If such an area of muscle was excised and explanted there was a profuse production of myoblasts in explants obtained both from normal mice and from dystrophic mice. He compared the explants obtained from six dystrophic and six normal adult mice and found certain distinctive differences. The nuclei of myoblasts from dystrophic cultures often contained only one or two nucleoli whereas the nuclei of normal myoblasts generally possessed three (Special Plate, Figs. 5 and 6) or more. Multinucleate myoblasts were found to be less common in dystrophic cultures than in the normal, and non-striated muscle straps containing many nuclei were observed in cultures from normal muscle but never in those from dystrophic muscle. There was also an irregularity of the margins of those myoblasts which were presumed to be dystrophic. These findings strongly suggested that the muscle cells in dystrophic animals were abnormal at an early stage of development.

Ross (1964) has begun a study of the nuclei of normal and dystrophic myoblasts using the interference microscope. He has shown that at the point when myoblasts fuse there is a considerable reduction in the volume of individual nucleoli within the nuclei of the cells engaged in this activity. He suggests that this process of fusion involves considerable metabolic activity which may temporarily lead to a depletion of the R.N.A. reserves in the nucleus. His preliminary observations, which will require confirmation, upon dystrophic myoblasts, suggest that these contain significantly smaller amounts of nucleolar material than do normal myoblasts and that they fail to fuse. Clearly much further work on tissue culture requires to be done before the significance of these findings can be elucidated, but despite the serious technical difficulties it seems likely that this method will be useful in the investigation of the dystrophic process.

Electrophysiological Studies

It is now well established that by means of the insertion of micro-electrodes into individual muscle cells *in vitro* and *in vivo* it is possible to make accurate recordings of membrane potentials both in animals and in man (Johns, 1958; Elmquist, Johns, and Thesleff, 1960; Creutzfeldt, Abbott, Fowler, and Pearson, 1963). Johns (1963) has summarized the differences between end-plate potentials recorded by this method and the muscle-action potential itself.

Important observations upon the phenomenon of myotonia have already been made. Creutzfeldt *et al.* (1963) have demonstrated a consistent reduction in membrane potential in patients with hyperkalaemic periodic paralysis, many of whom also show myotonia, and, as Johns has indicated, there is a possibility that the irritability of myotonic muscle may be due to a partial state of depolarization of the fibre membrane. On the other hand, in goats suffering from myotonia, Bryant (1962) has demonstrated that the resting membrane potential is normal but that small induced depolarizations may initiate trains of repetitive action potentials.

Lenman (1963) has found that the resting membrane potential recorded from the muscle of dystrophic mice of the Bar Harbor strain is consistently about 10–20% lower than in normal mice. These results have been confirmed by McComas and Mossaw (1963) in my department; they recorded from 255 fibres in normal mice (mean 83.0 mV) and from 225 in dystrophic mice (mean 76.0 mV) and there was much greater variation between individual fibres in the dystrophic animals (Fig. B). The further application of this technique to the study of muscular dystrophy and of myotonia in both man and animals will be likely to yield information of much value. In particular, it will be important to study the propagation of the electrical impulse along the muscle-fibre membrane which is probably damaged by the dystrophic process. It should also be possible to study the inward spread of excitation from the membrane to the myofibril, a process in which the sarcotubular system is undoubtedly of great importance. These studies are as yet in their infancy, but are likely to contribute a great deal to our understanding of the behaviour of the muscle cell in health and disease.

FIG. B.—Comparison of the membrane potentials recorded with intracellular micro-electrodes in normal adult mice (above) and adult dystrophic mice (below) of the Bar Harbor strain (McComas and Mossaw, 1963).

Conclusions

I have been able to do little more than to sketch in outline some of the clinical features of the principal varieties of human muscular dystrophy and to mention only a few of the recent advances in knowledge. Unquestionably some of these have been of great practical importance. Of particular value is the introduction of the serum creatine kinase as a routine diagnostic test for clinical and preclinical muscular dystrophy and in the identification of female carriers of the gene responsible for the most severe form of dystrophy—namely, the Duchenne variety which affects young boys. Of even greater

importance, however, are the fundamental research techniques now being utilized in the hope of elucidating the nature of the dystrophic process. Many theories have been adduced to explain why the muscles waste, and there are still a great many unanswered questions. One of these, which may be fundamental, is that we have as yet no explanation why the disease process is so curiously selective, particularly in its early stages, affecting some muscles and sparing others. The suggestion has been made that those most liable to be affected are the muscles first developed in embryonic life, but this suggestion has not been substantiated.

The recent attempt by Das Gupta (1963) to see whether phasic and tonic motor units are present in man and whether one of these is selectively involved in the dystrophic process has been unsuccessful, and as yet little information is available on how individual muscles vary in enzyme constitution. It may be that there is some specific biochemical reason, depending perhaps upon differential concentrations of enzymes, which explains the selective nature of the dystrophic process. Clearly, therefore, in attempting to increase our understanding of the disease we must compare not only the different forms of muscular dystrophy but also different muscles, both in normal individuals and in sufferers. Denny-Brown's (1951) suggestion that dystrophy might be due to an inability of the muscle cell to regenerate cannot be the whole story, since Walton and Adams (1956) and others have shown that regenerative phenomena are commonly observed in dystrophic muscle, even though such regeneration appears to be singularly ineffective, a finding which is confirmed to some extent by the results to date of tissue culture. It seems probable that the dystrophic process will eventually prove to be due to the absence of one or more enzymes within the muscle cell, but despite the clinical, pathological, and biochemical similarities between the various forms of muscular dystrophy it is possible that the biochemical processes concerned in each of these may be different, and it is even possible that each of the clinical subtypes may not always be determined by the same gene at the same locus. Clearly, therefore, a multitude of problems have yet to be surmounted, but one can, I think, conclude that the climate of research which has developed within the last few years has modified, at least to some extent, the attitude of despair previously so prevalent. The increasing volume of investigative work now being carried out in many parts of the world must surely give hope that before many years have passed the cause or causes of this tragic disease will be identified, and that when that time comes there may be some prospect of developing an effective treatment.

My colleagues and I are grateful to the Muscular Dystrophy Association of America, Inc., the Muscular Dystrophy Association of Canada, and to the Muscular Dystrophy Group of Great Britain for financial support, and I am indebted to Miss R. Allan for secretarial assistance.

REFERENCES

- Adams, R. D., Denny-Brown, D., and Pearson, C. (1953). *Diseases of Muscle*. Hoeber, New York.
- Aebi, U., Richterich, R., Colombo, J. P., and Rossi, E. (1962). *Enzymol. biol. clin. (Basel)*, **1**, 61.
- Banker, B. Q., Victor, M., and Adams, R. D. (1957). *Brain*, **80**, 319.
- Barwick, D. D. (1963). In *Research in Muscular Dystrophy*, p. 10. Pitman, London.
- , Newell, D. J., and Walton, J. N. (1963). *Neurology (Minneapolis)*, **13**, 12.
- , Osselson, J. W., and Walton, J. N. (1962). Presented to the E.E.G. Society, November 1962.
- , and Walton, J. N. (1963). *Amer. J. Med.*, **35**, 646.
- Batten, F. E. (1909). *Quart. J. Med.*, **3**, 313.
- , and Gibb, H. P. (1909). *Brain*, **32**, 187.
- Becker, P. E. (1953). *Dystrophia Musculorum Progressiva*. Thieme, Stuttgart.
- (1957). *Acta genet. (Basel)*, **7**, 303.
- , and Kiener, F. (1955). *Z. Neurol.*, **193**, 427.

- Bourne, G. H., and Golarz, M. N. (1959). *Nature (Lond.)*, **183**, 1741.
- Boyer, S. H., and Fainer, D. C. (1963). *Amer. J. Med.*, **35**, 622.
- Bryant, S. H. (1962). *Fed. Proc.*, **21**, 312.
- Buchthal, F. (1962). *Wld Neurol.*, **3**, 16.
- Chung, C. S., and Morton, N. E. (1959). *Amer. J. hum. Genet.*, **11**, 339.
- Creutzfeldt, O. D., Abbott, B. C., Fowler, W. M., and Pearson, C. M. (1963). *Electroenceph. clin. Neurophysiol.*, **15**, 508.
- Das Gupta, A. (1963). In *Research in Muscular Dystrophy*, p. 256. Pitman, London.
- Denny-Brown, D. (1951). *J. Neuropath. exp. Neurol.*, **10**, 94.
- Dowben, R. M. (1963). *New Engl. J. Med.*, **268**, 912.
- Dreyfus, J. C., Demos, J., Schapira, F., and Schapira, G. (1962). *C.R. Acad. Sci. (Paris)*, **254**, 4384.
- and Schapira, F. (1958). *Ann. N.Y. Acad. Sci.*, **75**, 235.
- and Demos, J. (1961). *2nd Int. Congr. Hum. Genet., Rome*, p. 80.
- Dubowitz, V. (1960). *Brain*, **83**, 432.
- (1963). *J. Neurol. Neurosurg. Psychiat.*, **26**, 322.
- and Pearce, A. G. E. (1961). *J. Path. Bact.*, **81**, 365.
- Duchenne, G. B. (1868). *Arch. gén. Méd.*, **11**, 5, 179, 305, 421, 552.
- Ebashi, S., Toyokura, Y., Momoi, H., and Sugita, H. (1959). *J. Biochem. (Tokyo)*, **46**, 103.
- Elmqvist, D., Johns, T. R., and Thesleff, S. (1960). *J. Physiol. (Lond.)*, **154**, 602.
- Emery, A. E. H. (1963). *Lancet*, **1**, 1126.
- Engel, W. K. (1962). *Neurology (Minneapolis)*, **12**, 778.
- Foster, J. B., Hughes, B. P., Huxley, H. E., and Mahler, R. (1961). *Brain*, **84**, 167.
- Erb, W. (1884). *Disch. Arch. klin. Med.*, **34**, 467.
- Evans, J. H., and Baker, R. W. R. (1957). *Brain*, **80**, 557.
- Fuchs, E. (1890). *Arch. Ophthalm.*, **36**, 234.
- Geiger, R. S., and Garvin, J. S. (1957). *J. Neuropath. exp. Neurol.*, **16**, 532.
- Gilliatt, R. W. (1962). *Brit. med. J.*, **2**, 1073.
- Golarz, M. N., and Bourne, G. H. (1963). *Muscular Dystrophy in Man and Animals*, Chap. 3. Karger, Basel.
- Gowers, W. R. (1902). *Brit. med. J.*, **2**, 89.
- Gresham, G. A., and Cruickshank, J. G. (1960). *J. Neuropath. exp. Neurol.*, **19**, 342.
- Herrmann, H. (1961). *Res. Publ. Ass. nerv. ment. Dis.*, **38**, 697.
- Hughes, B. P. (1963). In *Research in Muscular Dystrophy*, p. 167. Pitman, London.
- Hutchinson, J. (1879). *Med.-Chir. Trans.*, **62**, 307.
- Huxley, H. E. (1957). *J. biophys. biochem. Cytol.*, **3**, 631.
- Jackson, C. E., and Carey, J. H. (1961). *Pediatrics*, **28**, 77.
- Johns, R. J. (1958). *Res. Publ. Ass. nerv. ment. Dis.*, **38**, 704.
- (1963). *Amer. J. Med.*, **35**, 611.
- Julian, L. M., and Asmundson, V. S. (1960). *Anat. Rec.*, **136**, 218.
- Keevil, T. (1953). *Bull. Hist. Med.*, **27**, 201.
- Kiloh, L. G., and Nevin, S. (1951). *Brain*, **74**, 115.
- Kloepfer, H. W., and Talley, C. (1958). *Ann. hum. Genet.*, **22**, 138.
- Kugelberg, E., and Petersén, I. (1949). *J. Neurol. Neurosurg. Psychiat.*, **12**, 268.
- Lamy, M., and de Grouchy, J. (1954). *J. Génét. hum.*, **3**, 219.
- Landouzy, L., and Dejerine, J. (1884). *C.R. Acad. Sci. (Paris)*, **98**, 53.
- Lenman, J. A. (1963). In *Research in Muscular Dystrophy*, p. 230. Pitman, London.
- Levene, P. A., and Kriszteller, L. (1909). *Amer. J. Physiol.*, **24**, 45.
- Leyden, E. (1876). *Klinik der Rückenmarks-Krankheiten*, vol. 2, p. 531. Hirschwald, Berlin.
- Lyon, M. F. (1961). *Nature (Lond.)*, **190**, 372.
- (1962). *Amer. J. hum. Genet.*, **14**, 135.
- McArdle, B. (1963). In *Research in Muscular Dystrophy*, p. 145. Pitman, London.
- McComas, A. J., and Mossaw, S. J. (1963). Unpublished observations.
- Mair, W. G. P. (1963). In *Research in Muscular Dystrophy*, p. 112. Pitman, London.
- Meryon, E. (1852). *Med.-chir. Trans.*, **35**, 72.
- Michelson, A. M., Russell, E. S., and Harman, P. J. (1955). *Proc. Nat. Acad. Sci. (Wash.)*, **41**, 1079.
- Möbius, P. J. (1879). *Samml. klin. Vortr.*, **171**, 1505.
- Morton, N. E., and Chung, C. S. (1959). *Amer. J. hum. Genet.*, **11**, 360.
- and Peters, H. A. (1963). *Muscular Dystrophy in Man and Animals*, Chap. 8, edited by G. H. Bourne and M. N. Golarz. Karger, Basel.
- Nirenberg, M. W. (1963). *Sci. Amer.*, **208**, 80.
- Oppenheim, H. (1900). *Mtschr. Psychiat. Neurol.*, **8**, 232.
- Pearce, G. W. (1963). In *Research in Muscular Dystrophy*, p. 75. Pitman, London.
- (1964). In *Disorders of Voluntary Muscle*, Chap. 9, edited by J. N. Walton. Churchill, London.
- Pearce, J. M. S., and Walton, J. N. (1964). In preparation.
- and Walton, J. N. (1962). *J. Path. Bact.*, **83**, 535.
- Pearce, J. M. S., Pennington, R. J., and Walton, J. N. (1964a). *J. Neurol. Neurosurg. Psychiat.*, **27**, 1.
- (1964b). *Ibid.*, **27**, 96.
- (1964c). *Ibid.* In press.
- Pearse, A. G. E. (1963). In *Research in Muscular Dystrophy*, p. 189. Pitman, London.
- Pearson, C. M. (1957). *New Engl. J. Med.*, **256**, 1069.
- (1962). *Brain*, **85**, 109.
- (1963). *Amer. J. Med.*, **35**, 632.
- Fowler, W. M., and Wright, S. W. (1963). *Proc. Nat. Acad. Sci. (Wash.)*, **50**, 24.
- Pennington, R. J. (1962). *Proc. Ass. clin. Biochem.*, **2**, 17.
- Perkoff, G. T., Hill, R. L., and Tyler, F. H. (1962). *J. clin. Invest.*, **41**, 1391.
- Perlstein, M. A., Gibbs, F. A., Gibbs, E. L., and Stein, M. D. (1960). *J. Amer. med. Ass.*, **173**, 1329.
- Philip, U., Walton, J. N., and Smith, C. A. B. (1956). *Ann. hum. Genet.*, **21**, 155.
- Price, H. M. (1963). *Amer. J. Med.*, **33**, 589.
- Ross, K. F. A. (1964). *Quart. J. micr. Sci.* In press.
- and Schapira, G. (1962). *Biochemistry of Hereditary Myopathies*, p. 146. Thomas, Springfield, Illinois.
- Shy, G. M., Engel, W. K., Somers, J. E., and Wanks, T. (1963). *Brain*, **86**, 793.
- and Magee, K. R. (1956). *Brain*, **79**, 610.
- Sibley, J. A., and Lehninger, A. L. (1949). *J. biol. Chem.*, **117**, 859.
- Steinert, H. (1909). *Disch. Z. Nervenheilk.*, **37**, 58.
- Stevenson, A. C. (1953). *Ann. Eugen. (Lond.)*, **18**, 50.
- Thomson, W. H. S., and Guest, K. E. (1963). *J. Neurol. Neurosurg. Psychiat.*, **26**, 111.
- Leyburn, P., and Walton, J. N. (1960). *Brit. med. J.*, **2**, 1276.
- Turner, J. W. A. (1940). *Brain*, **63**, 163.
- (1949). *Ibid.*, **72**, 25.
- and Lees, F. (1962). *Ibid.*, **85**, 733.
- Tyler, F. H., and Wintrobe, M. M. (1950). *Ann. intern. Med.*, **32**, 72.
- Van Breemen, V. L. (1960). *Amer. J. Path.*, **37**, 333.
- van den Bosch, J. (1963). In *Research in Muscular Dystrophy*, p. 23. Pitman, London.
- Walton, J. N. (1952). *J. Neurol. Neurosurg. Psychiat.*, **15**, 219.
- (1955). *Ann. hum. Genet.*, **20**, 1.
- (1956a). *J. Neurol. Neurosurg. Psychiat.*, **19**, 106.
- (1956b). *Lancet*, **1**, 1023.
- (1956c). *Ann. hum. Genet.*, **21**, 40.
- (1957). *J. Neurol. Neurosurg. Psychiat.*, **20**, 144.
- (1961). *Res. Publ. Ass. nerv. ment. Dis.*, **38**, 378.
- (1963). In *Muscular Dystrophy in Man and Animals*, Chap. 7, edited by G. H. Bourne and M. N. Golarz. Karger, Basel.
- (1964). In *Disorders of Voluntary Muscle*, Chap. 11, edited by J. N. Walton. Churchill, London.
- and Adams, R. D. (1956). *J. Path. Bact.*, **72**, 273.
- (1958). *Polymyositis*. Livingstone, Edinburgh.
- and Nattrass, F. J. (1954). *Brain*, **77**, 169.
- Wayne, H. L., and Browne-Mayers, A. N. (1959). *Dis. nerv. Syst.*, **20**, 288.
- Welander, L. (1951). *Acta med. scand.*, Suppl. No. 265, p. 1.
- (1957). *Acta Genet. (Basel)*, **7**, 321.
- Whorton, C. M., Hudgins, P. C., and Connors, J. J. (1961). *New Engl. J. Med.*, **265**, 1242.
- Wieme, R. J., and Herpol, J. E. (1962). *Nature (Lond.)*, **194**, 287.