Kelvin, and Sir George Stokes failed to ascertain, or even recognize when they had been pointed out, the Royal Society should have made some acknowledgement of his work, especially as it utterly discredited him and ruined his career. As I have shown, many have heard of his discredit, but are unaware of the complete change of opinion. I have written this article to show these people exactly how matters stand.

COLECTOMY FOR INTESTINAL STASIS.

BY

A. WILLIAM SHEEN, M.S., F.R.C.S.,
SURGEON, KING EDWARD VII'S HOSPITAL, CARDIFF.

I HAVE done four cases of colectomy for intestinal stasis—one in a man and three in women. Three cases recovered and one died.

History, Symptoms, and Progress after Operation.

CASE I.—W. C., a mason, who had not worked for three months and had done only light and intermittent work for eight years, complained of abdominal pain one hour after meals. He was very constipated, had often gone a week without bowels acting, and was constantly taking castor oil or salts. He had had sickness after nearly every meal, but not lately; he sometimes made himself sick; he had vomited blood (a teacupful) on six or seven occasions. He suffered from distension, flatulence, had got thinner, and was very miserable, said life was not worth living, and felt inclined to finish himself off. He had taken much medicine, and was treated medically in hospital four years ago without relief. He was a thin man, with sallow complexion and dark rings round his eyes; the tongue was coated, the right kidney was movable. There was pain, tenderness, and gurgling over the caecum, tenderness beneath the upper part of the right rectus, and pain in the epigastrium. The urine contained creatin and creatinin in large quantities. A test meal gave total acidity 0.1606 per cent. X-ray photographs of bismuth meals showed great delay, and dropping of the transverse colon so that its centre reached the bottom of the pelvis. After operation, on November 6th, the parietal wound suppurated slightly. The patient got up on November 25th, and on December 11th, five weeks after operation, went home.

CASE II.—A. L., housewife, had been "ailing" many years; suffered from indigestion, and was very constipated. Eight years ago she was fitted with a supporting ring for "womb trouble." The symptoms had been worse for the last four months, and for three months she had passed blood frequently from the bowel, and had grown much thinner. She was a thin, anaemic woman, with lemon-yellow complexion, dark rings beneath the eyes, cold extremities, and a miserable appearance. The lower abdomen was distended, and there was much gurgling of gas; tenderness was most marked in the left iliac region, but was present also over the caecum and pylorus. X-ray photographs of bismuth meals showed, after half an hour, the stomach dropped, so that the greater curvature reached the pelvis, and, after twenty-four hours, large masses in the caecum, with distension of colon and sigmoid. Examination with the sigmoidoscope revealed nothing abnormal. After operation (January 22nd) she at first went on well, but later developed progressive asthenia and cardiac weakness without active symptoms; there was no parietal infection. She died on January 31st, nine days after operation. An examination *post mortem* through the operation wound only showed thin pus in the pelvis and lower abdomen; the bowel wounds were intact.

CASE III.—A. M. S., a married woman, complained that for four or five years she had had pain in the left side of the abdomen, particularly in the lower part. It was worse directly after food; she sometimes vomited, and was only able to take "baby's food." She was very constipated, and passed a quantity of whitish material and "strings" with the motions and at other times. She had had an enema once daily for two years. She was very thin, with a yellow-brown complexion and darkish red areas on the cheeks. There was much tenderness all over the course of the colon, and mucus and shreds in the motions. The sigmoidoscope showed the bowel to be very red. An x-ray photograph of a bismuth meal showed that the transverse colon reached to the pubes. Operation was postponed because after a preparatory enema the patient became dizzy, lost consciousness for two hours, and had great cardiac weakness. It was eventually performed on March 2nd. She did well; no parietal infection occurred, and she left the hospital one month after operation.

CASE IV.—C. A. had been operated on three years earlier for "ulcerated stomach," and had been better until five months ago, when vomiting and pain commenced after food. She had been very constipated for years, and had had three slight attacks of haematemesis. She was a thin woman with a coated tongue and earthy complexion. She vomited frequently after admission, sometimes a little blood. There was tenderness over the caecum and duodenum, a vertical median scar in the epigastrium, and slight oedema of legs. Test meal: Total acidity 0.1752 per cent.; urine, trace of creatin. Mobile right kidney. The case was transferred from a medical ward after prolonged

treatment, including morphine frequently. After operation (April 30th) she did well except for slight parietal infection and obstinate cystitis. She left hospital seven weeks after operation.

Details of Operation.

The conditions found in the abdomen and the operative details were substantially the same in all four cases. The preparation, which commenced three days before operation, consisted of enemata, castor oil, clear soup, albumin water, tea, brandy. Case 3 had kerol gr. v thrice daily for a fortnight before operation. Anaesthesia was by scopolamine and open ether. A continuous submammary saline injection was given throughout the operation. The patient was placed in a partial Trendelenburg position, with the legs over the end of the table. A long vertical incision was made through the inner part of the left rectus. In each case external adhesions were found between colon and parietes and kinking at hepatic and splenic flexures, particularly the hepatic, where the transverse and ascending colon were bound together, ileal kink, dilated duodenum, prolapsed stomach, and transverse colon. In Case 4 (previously operated on) the stomach was adherent to the anterior parietes and the duodenum was most markedly dilated. There were various omental, gall bladder, duodenal, and other adhesions and varying degrees of membrane overspreading the caecum and ascending colon. After separating external and other adhesions, so as to render the colon freely mobile and permit of its being lifted out of the abdomen, it was removed from 8 in. to 9 in. proximal to the ileo-caecal valve to the middle of the pelvic colon; the distal end of the colon was closed and the ileum joined to the colon by an end-to-side anastomosis, taking care that the ileum was not twisted. The hole in the mesentery was carefully closed and an oesophageal tube passed up the rectum and through and about a foot beyond the anastomosis, the operator manipulating the tube and an assistant pumping paraffin in to help its progress. This tube was tied to the anal margin. The parietal wound was closed in layers without drainage, except in Case 3, in which a tube was kept in the abdomen for forty-eight hours.

The dressing throughout was sterilized, frequently changed, hot boracic fomentations. The patient was placed in a marked Fowler's position. The rectal tube drained well in all cases, a few ounces of paraffin being pumped up it every twenty-four hours. It was taken out in seven or eight days. All cases showed some shock after the operation, but it was only marked in Case 2, where the pulse was very weak and rapid from the first.

End Results.

The three cases which recovered were all seen and examined by me on October 5th, 1914, when their condition was as follows:

CASE I (eleven months after operation).—Says he is much better than before the operation, that he can eat practically anything, and that he never vomits. He still feels a weakness in his abdomen and has loose motions, the bowels acting two or four times a day. He has tried two or three times to do his ordinary work, but cannot manage it. He has not got fatter or thinner. On examination he is pale, his eyes are clear, with no dark rings round them. He wears a belt; the recti are slightly separated. There is epigastric tenderness.

CASE III (eight and a half months after operation).—Says she is much better and much fatter; can eat anything now. All her people say how very different she looks. She sometimes has a pain in the back and in the lower abdomen when she walks, and she suffers from palpitation. She takes a tablespoonful of paraffin every morning, and this keeps the bowels regular. She is never sick. On examination she is a fat, healthy looking woman with a good complexion and clear, bright eyes. There is a sound abdominal scar; some tenderness over sigmoid region. She is emphatic in saying that she is better than she has been for many years.

CASE IV (five months after operation).—Much better. Put on a stone in weight after she went home. Her friends say she looks quite different. She is sick at her periods, which are profuse. She takes syrup of figs occasionally for the bowels. She is a healthy looking, well-nourished woman, with bright eyes and clear complexion. There is a strong abdominal scar, with no tenderness.

The operation is a serious one. It requires careful preparation of the patient and careful technique. It should be reserved for cases with marked signs and symptoms of stasis. It contrasts, in my experience, with many cases in which various lesser measures have brought about cure. In all these abdominal cases—which can be grouped under

no short title (unless "stasis" be used), but which present symptoms suggesting trouble now in one region, now in another—it should not be the practice to approach and expose simply the part where the trouble is thought most likely to be, but by a wide incision to investigate the whole of the abdominal contents and to recognize and, as far as possible, remedy the various abnormalities.

The end results, as far as can be judged from these few cases, show that, while great improvement may be expected, a perfect condition of health is not necessarily attained.

AN OPERATION FOR MOVABLE KIDNEY.

By G. S. THOMPSON, F.R.C.S. Eng.,
KIMBERLEY.

In this operation the essential desiderata are (1) a guarantee against recurrence of the undue mobility, and (2) an assurance that the fixation shall be satisfactory, by restoring and maintaining the normal position of the organ.

None of the ordinary operations—extracapsular, capsular, suture and tendon-fixation, etc.—can be said to fulfil these essentials. All the usual methods are faulty owing to their uncertainty, whilst their very number suggests that no one is adequate and sure, which is, indeed, true. Hence we may say that so far no good operation has been generally practised for nephroptosis.

The procedure to be described gives a virtually normal fixation of the organ, and if properly carried out can be relied on almost certainly to ensure a satisfactory and permanent cure. By this method the kidney retains with certainty the position in which it is fixed, and it cannot become displaced again; and, owing to this very fact, care must be taken to see that the correct or proximate position is obtained before anchoring in place. Moreover, although the pathological mobility is corrected by this operation, the normal slight respiratory excursions of the organ are not hindered. One advantage more is that the patient can be got up very much earlier than in the ordinary way—in fact, soon after the wound is healed—namely, by the third week, in contrast with the inevitably prolonged, but not always successful, detention in bed with the other methods.

The principle of this operation is the insertion of the kidney into, and its suspension by, a sling, and the fixation of the latter to the posterior abdominal wall.

The sling consists of a net made from chromicized catgut, or preferably floss silk, the material at present varying according to the predilection of the surgeon, as time alone can show which is to be preferred. These slings are made in various sizes, so that there shall be no difficulty in adapting one to the requirements of the particular case. A gap is left at the hilum 2 in. long by 1 in. broad, so that the border of the net here falls well short of the vessels and duct, and thus no pressure can be exercised on these important structures. The convex border is left open to facilitate the insertion of the kidney into the sling, and at the ends of this opening two free ligatures are left open for the purpose of lacing up the gap, and then using to sling the net to the abdominal wall. The complete sling is thus reniform with a permanent gap at the hilum and a temporary slit along the outer border, the size of the mesh being about $\frac{1}{2}$ cm.

The kidney, having been exposed in the usual way, is inserted through the gap and by the help of the slit into the sling; it is then laced in, but loosely, and when this is completed the ligatures are knotted and thus fixed, the lower ligature ending above opposite the costal groove on the kidney. The lacing should not be tight so that the sling fits the kidney somewhat loosely, in order to allow of the expansion of the organ which is known to occur. It must not be loose enough, however, to permit the net to rotate and impinge by its free inner border on the structures of the hilum, thereby compressing them; and in order to obviate this it is as well to insert one or two fixation sutures so as to include the net and kidney and maintain the proper relative positions of both. The kidney having been invested, the next step is to fix it in position by the free ligatures along the outer border. One is passed external to the normal position of the kidney over the twelfth, or if necessary the eleventh rib,

the other through the muscles near the lower end of the viscus, and the two ends tied loosely under the skin; loosely, again, in order that the kidney may be free to move with respiration and adapt itself spontaneously to the exact position, after the patient has been placed in bed with the foot blocked up, should such not have been quite correctly obtained at the time of operation—not always an easy matter to be certain of with the patient in the lateral position. The remaining steps of the operation are as usual.

The netting by its presence will provoke a celluloplastic exudation which will surround and infiltrate the meshwork, and in time form by organization a firm adventitious tissue capsule, in addition to the extraneous capsule already present. Probably the former would be sufficient to permanently fix the organ if a slowly absorbable material, as chromic catgut, be used. This seems very likely, but to obviate any possible doubt on this point perhaps it would be better, as suggested, to use floss silk. I say floss silk, and not the ordinary twisted surgical silk, as I think the former smooth material less liable to determine haematogenous bacterial deposit than the rough irritating surface of the latter. In both cases a fibrous tissue capsule will form; with catgut this alone will be permanent, but with silk both will remain. I do not think it likely that floss silk, by its permanent presence, would provoke an excessive formation of fibrous tissue, and thus produce an unyielding covering not altogether desirable in an expansile organ like the kidney; but this might occur with the roughened variety. The only disadvantage of silk in contrast to catgut is the slight tendency of the latter to become the seat of remote haematogenous infection, and give rise to abscess, assuming, of course, the wound heals, as it should, by primary union, and there is no immediate soiling from ungloved hand contamination, as is the rule when such occurs, and not through any imaginary difficulty in sterilizing silk. But the use of floss silk will also tend to obviate this contingency.

When I first developed the above method, in 1909, I used for the posterior base of the sling a reniform plate of thin sheet celluloid, riddled with punch-holes, the remainder of the sling being as above and the netting attached to the border of the plate. When placed in the body this plate soon becomes fixed to the neighbouring parts by infiltration. But I think the elimination of the plate an improvement in technique, and have not mentioned it nor used it latterly.

The first patient operated on, in 1909, is still quite well and free from all her previous troublesome symptoms, whilst the organ still remains quite fixed. The wound healed by primary union, and the patient left the hospital at the end of three weeks.

I think it can be claimed that this operation is an advance on any other, and is sure, quick, rational, simple, as well as satisfactory from the patient's point of view. I consider its merits should cause it to supplant the other faulty and bad procedures which are in vogue at present. A good operation for nephroptosis has long been required, and in the belief that this one meets the case and fulfils all necessary requirements it is submitted with confidence.

L.S.L., Lacing and suspending ligatures; Sl., kidney sling containing kidney (dotted); O.S., outer slit; H.G., hilum gap; H.S., structures of hilum—vessels and duct; K., kidney fixed over twelfth or eleventh rib, triple anchorage by ligatures and pedicle.