milk to ordinary diet, and entirely recovered. But Mr. Woodhouse quite tired himself out by the care and vigour he had to exert to get this plan carried out. Several times we were almost reduced to despair of success. You must prepare for a course of decisive and sometimes unrequited labour in managing these cases.

The conclusions to which I have endeavoured to lead you by aid of three patients under your eyes are—

1. Hysteria is a disease (a word which I always use as synonymous with deficiency of life) as much of the mind as of the body.

2. In some cases the mental, in others the bodily, phenomena predominate.

3. The predominance of one or other must be our guide whether moral or physical agents are most required in the treatment.

4. The aim of the moral treatment must be the teaching the patient to exert the will; and that is best done by a change of habits and scene.

5. The organ which aids us most in our physical treatment is the stomach.

Original Communications.

THE TREATMENT OF DIPHTHERIA.

By JOHN WEST WALKER, M.B.Lond, etc., Spilsby.

NOTWITHSTANDING the great accumulation of knowledge bearing on the pathology of diphtheria, I believe there is a general feeling of doubt and dissatisfaction throughout the profession as to the most efficacious mode of treatment. I, therefore, feel justified in calling attention to a plan of treatment the decidedly beneficial effects of which I have had ample opportunity of testing.

Avoiding all discussion as to the nature of diphtheria, all will agree with me in noticing the three following salient characteristics of the disorder, each of which possesses a decided therapeutic indication, namely: 1. A more or less acute unhealthy form of inflammation affecting the faucal region, and which, if considered singly, would lead us to associate the complaint with common cynanche tonsillaris. 2. The presence of the peculiar leathery exudation which again associates the disorder with canker, stomatitis, and other allied affections. 3. The great, often suddenly induced, debility and tendency to death by exhaustion, indicate the typhoid character of the complaint.

For each of these several characteristics, our materia medica possesses a remedy of almost specific virtue.

I have long looked upon guaiacum as a specific in common sore throat; for, no matter in what place, what the age, sex, or condition of the patient, or what the stage of the complaint, I have always administered it, and had reason to be satisfied with the result; indeed, I consider the effect of this remedy in such cases to furnish an example of cure as distinguished from treatment. (*Vide* Dr. Latham's Lectures.) Frequently have I seen cases of cynanch and diphtheria existing at the same time and in the same family; and I was first induced to give guaiacum in the more severe affection by observing its good effect in cases which, at the onset, were supposed to be examples of common sore throat. I now unhesitatingly consider guaiacum as the remedy *par excellence* for diphtheria. The presence of the peculiar exudation almost instinctively suggests the employment of the chlorate of potash. For the debility, etc., our sheet-anchors are ammonia and bark.

My prescription for all cases of diphtheria is the following:—

R Potassæ chloratis ℥iv; tincturæ cinchonæ comp. ℥ss; tincturæ guaiaci comp. ℥ss ad 3vj; mellis q. s.; aquæ ad 3vij. M.

Of this mixture from one tea- to two tablespoonfuls may be given, according to age; but the interval between each dose, from one to four hours or three times a day, must be regulated according to the severity of the case.

These, then, I consider the therapeutic essentials in the treatment of diphtheria; but, in a complaint so terrible, it behoves us to resort to every possible adjuvant, however slightly it may contribute to the general good; and under this heading I am disposed to place all topical applications. My plan is to apply (by no means frequently) a concentrated solution of nitrate of silver, believing it to hasten the removal of the exudation, and to induce a more healthy action over the denuded surface. Of gargles, I consider weak vinegar and water, or what is more elegant, dilute acetic acid with infusion of roses, to be as good as any, producing slightly stimulant and astringent effects on the parts, and washing away offensive detached particles. Externally, warmth applied after the manner of poultices, of which a bag of chamomile-flowers is as convenient as any, together with mild terebinthinate embrocations applied two or three times a day, are to be recommended. Like most of my fellow-practitioners, I avoid mercurials; and, should the bowels require moving, castor-oil or the milder laxatives suffice.

The diet should be nutritious, with an increasing allowance of wine from the first. Lastly, we should ever be on the alert to remove as far as practicable the primary cause of the complaint by attention to every sanitary particular.

In advocating the guaiacum treatment of diphtheria, I must disclaim all credit for originality. Others have recommended it to the notice of the profession; and more, I doubt not, are daily in the habit of carrying it out in practice. What I desire is, that many of my fellow-practitioners may be induced to try a remedy which, from what I have lately read on the subject, is not so extensively employed as it deserves to be.

EXCERPTS FROM DAILY PRACTICE.

By T. HERBERT BARKER, M.D., F.R.S.Edin., Bedford.

IV.—*Case of Poisoning by "Almond Flavour." Treatment. Emetics. Cold Affusion. Stimulants. Recovery. Remarks. Composition of "Almond Flavour." Analogous Cases.*

IN the evening of the 6th of December last, I received an urgent message to see a young lad residing near to my house. The messenger said he had been poisoned by drinking "almond flavour." I took at once with me a bottle containing ipecacuanha powder. The little patient, a youth of six years of age, was much prostrated, very pale; the eyes bright, prominent, and glassy, and the pupils dilated. The pulse was small and quick. He appeared to have considerable uneasiness in the throat; but fortunately the power of swallowing had not been lost. A brimming teaspoonful of the powdered ipecacuanha was immediately mixed with warm water and given to him, and the cold affusion had recourse to. Within three minutes from the first dose, as vomiting had not commenced, I mixed another teaspoonful of powdered ipecacuanha with warm water, adding to it a small quantity of mustard and common table salt. I then gave him brandy and water at frequent intervals. Within two minutes after the second dose, free vomiting commenced. The ejected matters smelled strongly of bitter almonds. The cold affusion was persevered with, and the frequent administration of warm brandy and water,