

## Asthma induced by epoxy resin systems

We are all environmentalists nowadays, but an interest in the air we breathe has long been of concern to chest physicians, particularly in the management of asthmatic patients. In addition to the common atmospheric and domestic allergens, patients may be exposed in their work to chemicals such as the isocyanates, piperazine, and formalin, which may provoke asthmatic attacks. More recently, interest has been focused on the epoxy resin systems.<sup>1 2</sup>

Epoxy resin systems are used in industry as adhesives, reinforced plastics, moulding resins, and surface coatings. The resins are converted to the finished product by adding curing agents or hardeners. When curing agents such as the acid anhydrides are used the process requires additional heat, while when cold curing agents such as the amine compounds are used the process generates its own heat. In either case, fumes of curing agent may be given off and may provoke asthma in a susceptible person.

This asthmatic response appears to be a true hypersensitivity,<sup>1 3</sup> for there is no reaction on first exposure, but after a latent period asthma develops on exposure to low concentrations, though only in a minority of the exposed population. The bronchial reaction may be immediate, non-immediate, or dual, and may be elicited by a few breaths of the sensitising fumes. We still do not know the precise immunological mechanisms, but Maccia *et al*<sup>4</sup> reported an instance of phthalic acid anhydride sensitivity in which the patient had a raised titre of specific IgE. Another patient did not have a raised titre,<sup>2</sup> but the delayed part of the dual broncho-constrictor response was inhibited by pretreatment with sodium cromoglycate.

A careful history, with particular attention to chemical processes at work, will usually alert the physician to the possibility that asthma is occupational. The typical pattern is of wheeze occurring either immediately or several hours after exposure with partial or complete recovery at weekends and during holidays. The latent period between first exposure and asthma may vary from a few days to months or even years. A known exposure to a known causative agent makes the diagnosis more likely, while objective confirmation of an occupational cause may sometimes be obtained by asking the patient to keep a work diary and record his peak expiratory flow three or four times a day. Absolute identification of the precise cause requires referral for careful bronchial provocation testing,<sup>5</sup> when a fall in peak flow or airway conductance after inhalation of controlled amounts of the suspected agent confirms the diagnosis. The procedure is potentially hazardous—a severe attack of asthma may ensue—and should be undertaken only in hospital, with observation overnight in case a delayed reaction occurs.

Provocation testing may help in individual management by confirming a suspected cause and may occasionally show that the putative cause is innocent. Further testing may incriminate some unsuspected agent. Provocation tests are also important in identifying new causes of occupational asthma and may prove of value in clarifying the mechanisms responsible.

Acute attacks of asthma provoked by epoxy resin systems respond to the usual treatment with bronchodilators. Systemic steroids should be added if the attack is severe or protracted. Sodium cromoglycate and inhaled steroids may help to control recurrent or continuous asthma. Clearly, however, the best

treatment is complete avoidance of exposure, which may require a change of occupation. Such advice should not be given lightly, and if possible it should be backed by objective evidence of an occupational factor identified by formal provocation testing. With complete exclusion of the precipitating cause the asthma may resolve completely, and the outlook is then excellent. In contrast, continuing exposure may cause increasing sensitisation with more severe and chronic asthma requiring continuous suppressive treatment, leading eventually to permanent deterioration of lung function.

<sup>1</sup> Fawcett, I W, Newman Taylor, A J, and Pepys, J, *Clinical Allergy*, 1977, **7**, 1.

<sup>2</sup> Chester, E H, *et al*, *Clinical Allergy*, 1977, **7**, 15.

<sup>3</sup> Dernehl, C U, *Journal of Occupational Medicine*, 1963, **5**, 17.

<sup>4</sup> Maccia, A C, *et al*, *American Review of Respiratory Diseases*, 1976, **113**, 701.

<sup>5</sup> Pepys, J, and Hutchcroft, B J, *American Review of Respiratory Diseases*, 1975, **112**, 829.

## Soft contact lenses

Recently there has been a sharp swing towards the use of soft, as opposed to hard, contact lenses. This has come about because these can be worn for longer periods, and soft lenses are now widely used in three different sorts of conditions: for low refractive error in normal eyes, in aphakia, and as a protective membrane in many corneal disorders. After eliminating any discussion of the third category, we may ask how far the results of wearing soft contact lenses for the first two compare with those of other established methods of optical correction.

Low refractive error in normal eyes may, of course, be corrected by spectacles, but many patients prefer contact lenses because of their convenience, efficiency, and cosmetic appearance. In this group soft contact lenses therefore have to be compared with hard contact lenses. To take the disadvantages: firstly, most eyes cannot tolerate them for extended periods even with the high water content Sauflon 70 and Sauflon 85 lenses. There is a limit to the degree of hypoxia that the endothelium of the cornea will tolerate. Secondly, even if a particular eye can tolerate extended wear, however, there is still a danger of bacterial or fungal keratitis. The latter may be disastrous and may be masked by the lens while it remains in wear. Thirdly, complicated routines for cleaning the lenses are necessary. Fourthly, the lenses are fragile and they tend to cost more than hard lenses. Fifthly, when a lost or damaged soft lens is replaced by another apparently identical lens both the comfort and vision may be poorer than with the first lens. Soft lenses lack the reproducibility of hard lenses. On the other hand, soft lenses do have certain definite advantages for low refractive errors in normal eyes: they are easier to get used to than hard lenses, and they are more easily managed by the intermittent wearer.

In aphakia optical correction by spectacle lenses admittedly has certain disadvantages. Nevertheless, it is wrong to say that patients are seldom happy with their post-cataract glasses, since after an initial period of learning the great majority manage very well indeed. This fact is forgotten, however, by those who advocate optical correction by intraocular acrylic lenses or extended-wear soft contact lenses. Both methods have inherent dangers, which are avoided with conventional spectacles. Intraocular acrylic lenses and soft contact lenses should therefore be compared in the knowledge that both introduce avoidable dangers. Even when introduced skilfully,

intraocular acrylic lenses increase the death rate of endothelial cells (as does any intraocular procedure) to an extent which may irreversibly cloud the cornea. Cystoid oedema of the macula is commoner when such lenses are used than in the simple cataract extraction. Many operative and postoperative mishaps seem to be associated with the use of intraocular lenses, and the number of modified varieties testifies that something is still amiss with the whole principle.

Extended-wear soft contact lenses, on the other hand, are probably much less prone to give rise to disastrous complications, provided they are well-fitted and are of the high water content type. Even so, a patient with soft lenses should have frequent and prolonged follow-up in the clinic, and should not manipulate the lenses himself. He should be told that any discomfort or change in vision needs urgent attention.

In summary, therefore, hard lenses remain the contact lenses of choice for the ordinary phakic wearer with no disease. Soft lenses should be reserved for those who cannot tolerate hard lenses and for those who want to wear lenses intermittently. Extended-wear, sadly the most attractive feature of soft lens philosophy, is wise only if the patient follows instructions closely and has immediate access to skilled advice when something seems to be wrong. For aphakic patients spectacles are much safer. Nevertheless, the use of soft lenses should be explored further, if only to emphasise that we need to evaluate intraocular acrylic implants much more critically.

## Foreign bodies in the rectum

The eight external orifices of the human body seem to attract strange foreign bodies just as honey does bees. For size and variety of objects, pride of place goes to the rectum; but the circumstances responsible are varied. Firstly, thermometers, enema tips, and catheters may disappear within the rectum, and inspissated masses of barium may be left behind after radiological examination. Next is the therapeutic group: to relieve pruritus and prolapsing piles patients may use all sorts of blunt objects up to lamp bulbs, bottles, and broom-stick handles, and indeed several old-fashioned proprietary "treatments" for haemorrhoids consisted of obturators for insertion into the anal canal. Criminal assault is an occasional cause, and another source of trouble is swallowing sharp foreign bodies which then impact in the lower rectum to present as either an abscess or a fistula. Finally—and today it seems most common—there is the introduction of a wide variety of objects into the rectum for sexual gratification. Haft and colleagues<sup>1</sup> have recently reported two examples of women who presented with battery-driven vaginal vibrators lodged in the rectum during intercourse. One was removed per anum but the other required a laparotomy to dislodge the vibrator from the sigmoid colon into the rectal ampulla; in this instance the patient reported that the motor had continued to operate for five hours after insertion, surely a tribute to modern electric batteries.

Most reports of rectal foreign bodies are anecdotal and many more are simply recounted at medical mess dinners. A useful service has therefore been performed by Eftaiha and his colleagues<sup>2</sup> from the section of colon and rectal surgery at Cook County Hospital, Chicago (one of the largest hospitals in the world), who recently reviewed a five-year experience of the removal of 31 colorectal foreign bodies in 30 patients, all men. They suggest a classification according to physical properties (sharp objects, large round foreign bodies, and glass) and location (whether easily palpable in the rectal ampulla or out of reach in or proximal to the rectosigmoid junction). After clinical examination, radiographs were taken of the abdomen and pelvis in two planes in order to establish the type, number, and location of the objects. Low-lying foreign bodies were removed transanally under spinal or local infiltration anaesthesia, which allowed complete relaxation of the anal sphincter. Sharp objects (bone spicules, toothpicks, glass fragments, etc) were delivered through a proctoscope with every care to prevent further mucosal laceration. Large rounded objects were easily removed with the help of forceps when there was no risk of breaking the foreign body, but glass containers required special care to avoid fracture. The suction effect created by the upward direction of the mouth of the container was ingeniously released by the use of a couple of Foley catheters passed around the container and extended into the lumen of the bowel above the foreign object. After inflating the balloons of the catheters, air was injected. Applying traction to the catheters will also help in removal of the object. Breaking the suction effect of the glass container in this manner will usually be successful. Others have also described the use of obstetric forceps to remove bottles in the rectum.<sup>3 4</sup>

For high-lying foreign bodies the Chicago group advocate spinal anaesthesia. The patient is placed in the lithotomy position, the object is located through the sigmoidoscope and manipulated, if possible, by abdominal palpation into the rectal ampulla. In three patients, however, laparotomy proved necessary. In one the object could then be manipulated into the rectum and removed, but in two others colotomy was needed, one to remove a ballpoint pen and the other to evacuate a large triangular-shaped glass bottle. The introduction of more and more ingenious attachments to the flexible fibre-optic colonoscope will no doubt reduce still further the need to resort to laparotomy for removal of highly situated objects.

After removal of a foreign body sigmoidoscopy should be carried out to exclude mucosal lacerations, perforation, or a missed second foreign body, and the patient should remain in hospital for a day or two postoperatively to be observed for delayed symptoms and signs of perforation or of perirectal suppuration.

<sup>1</sup> Haft, J S, Benjamin, H B, and Wagner, M, *British Medical Journal*, 1976, **1**, 626.

<sup>2</sup> Eftaiha, M, Hambrick, E, and Abcarian, H, *Archives of Surgery*, 1977, **112**, 691.

<sup>3</sup> Moynihan, N H, and Thomas, S H, *St Thomas's Hospital Gazette*, 1951, **49**, 152.

<sup>4</sup> Peet, T N D, *British Medical Journal*, 1976, **1**, 500.