Emergencies in the Home

How I would treat hypothermia and exposure

JOHN C FRANKLAND

When asked how I would treat hypothermia and exposure my first thought was—with respect, as old adversaries who can be rather challenging. My second thought was that in the home I would not expect to meet this condition very frequently but, because of a special interest in the Yorkshire Dales, would see roughly 80 cases out-of-doors to one in the home.

Every social worker is now aware of hypothermia in the elderly, but it is still neglected in the training of doctors. I recently asked an audience of young general practitioner trainees who were assessing the adequacy of their emergency equipment how many carried a subnormal thermometer and found that none did. Hypothermia is defined as a core temperature below 35°C. A problem with such terminology is that it implies that everyone with a core temperature marginally above this level is free from risk. In the intensive care unit they may well be, but in a threatening and cold environment, where deterioration may be rapid, this is far from the case. Exposure lacks a precise definition and perhaps the "exposure-exhaustion syndrome" is more apt. It certainly is a likely.

Any general practitioner near to mountains, beaches, caves, or open water may be called to treat victims of hypothermia. Death may be rapid—for example, within one hour in caves—and there may be several victims, which is unlikely at home. On one day we have had nine exposed fell runners and on another up to 10 exposed caves. In the home the victims are infants, drunk or suicidally-minded adults, or more commonly the elderly, many of whom will be gravely ill from other causes.

Diagnosing hypothermia

First a degree of suspicion is essential. If your own hand is warm then laying it on the patient's forehead will give the first clue. If it feels cold to the touch then place the hand on an area of the trunk that is normally covered. This also feels cold then you must assume that hypothermia is present and proceed otherwise by means of a subnormal thermometer inserted into the rectum and read after three minutes. Any chemist can supply a subnormal thermometer. Buy at least two, because you may need them to check your own hands are cold and you may break it through fumbling.

Managing patients with hypothermia

The doctor must insulate the patient, warm the room, and then transfer the patient to hospital. As a first wrapping a "space blanket" which may be bought for under £1 in most climbing or outdoor pursuit shops, will minimise radiant heat loss in patients at any age. If you do not have one then minimum cooking foil or a space blanket is a good substitute. Then wrap the patient with blankets or whatever dry fabric is to hand. Surprisingly, many forget to dry the hands and feet. Outdoors with wind blowing a heavy duty polyethylene bag will give much more protection than a space blanket.

Lancaster LA1 3AQ

JOHN C FRANKLAND, MB, FRCP, general practitioner

Preventing well-meaning neighbours from applying hot water bottles indiscriminately may be difficult and they should be discouraged from doing this if one has to leave the patient to telephone. In hypothermia a large pool of cooled blood in the body shell is isolated from the circulation owing to peripheral circulatory status. Applying heat indiscriminately to the skin may just open up this reservoir and the return of this cooler blood to the core can be disastrous.

The patient should be handled as little as possible, and ambulance colleagues should be requested to keep the patient horizontal if this is feasible. The ambulance heater may have been set just for the comfort of the crew and it should be moved to maximum output and the doors kept closed until the patient is at the ambulance.

Perhaps all infant hypothermia is a due to negligence and most likely if the parents are educationally subnormal or mentally distressed, so you should look for other features of child abuse and consider calling in social workers. Ambulance services will supply a warmed cot for transport.

It is difficult to justify home treatment for elderly patients when studies done in hospital show death rates of 50%. Obtaining a hospital bed for elderly patients is always a problem for some doctors. Geriatric units may keep one special bed for hypothermic patients to that admission is facilitated. If such patients are occasionally found to have "warmed up in the ambulance" and are not too profoundly chilled on arrival then maybe the fault is with the system and not with the sentiment of the referring doctor.

DEATHS

The place of early drug treatment for cold patients is controversial and is well reviewed by Maclean and Emble-Smith.¹ Many elderly patients have life-threatening drug toxicity which must be treated initially, bearing in mind that drug actions may be unpredictable at low temperatures, that oral medication and even parenteral medication injected into cold muscle may be minimally absorbed, and that drugs are best given intravenously in titrated small doses. This advice must be qualified by the fact that finding veins in hypothermic patients may be extremely difficult and it is not always clear whether the most promising sites for later intravenous infusions in hospital.

The once routine use of parenteral hydrocortisone for elderly hypothermic patients is now advised against, perhaps as a last resort, but because many subsequently develop respiratory infection, it is worth while giving parenteral penicillin. Intravenous glucose has been advocated,² but the brutal Dachau experiments showed that hyperglycaemia develops in direct proportion to the degree of hypothermia so that primary care need not seem to demand intravenous glucose. My last hypothermic patient at the home, however, was also a diabetic, and only with intravenous dextrose did the regaining consciousness, start to shiver, and ceased to miss his meal the previous evening.

Treating exposure

For victims of hypothermia in remote spots the basis of treatment is still insulation and evacuation. With a core temperature below 35°C that is, hypothermia as defined—forcing them to exert themselves will cause a net heat loss by opening up the peripheral circulation and thus accelerate deterioration.³ Patients must be carried. Traditionally removing wet clothing and replacing it with dry insulating clothing

is taught,⁴ but in practice most rescue teams will minimise further chilling by just removing outer wet clothing and putting their dry insulation round the more wet clothing, which will be at skin temperature. The type of fabric used, but Keighley⁵ has shown that a synthetic fibre called "Ektelon" sleeping bag is the most efficient, which as a bonus retains high insulating qualities when wet. In cave rescue thick Neoprene exposure bag is almost routinely used because further immersion during rescue is likely. The "wind chill factor" can be more confusing than "wind speed" as it is a measure of what fabric is used must have a totally windproof outer layer.

In English mountains many rescue teams now consider delaying evacuation to allow a patient to establish a "space blanket" as a survival bag (Karrimor)⁶ round both the victim and rescue team, wind chill is prevented, the air temperature is raised, injuries can be tended to, and adequate insulation is provided. Evacuation by helicopter is justified if weather conditions permit.

In 1972 Lloyd⁷ described airway rewarming as the first on-site method of treating hypothermia. In 1974 a report of an expedition of its successful use in the field.⁸ Since then the method has aroused controversy. In 1975 a portable device (the Reviva)⁹ was marketed and recommended as the method of choice for on-site treatment of acute onset hypothermia.¹⁰ Since then Audi¹¹ has studied the technique in dogs, claiming negligible benefit, and Collis¹² also reported a significant benefit in hypothermic patients. It is mandatory in man, making it the preferred first-aid technique. In the world of mountain and cave rescue many anecdotal stories attribute survival to this technique, and American rescue teams are rapidly acquiring such equipment. The heat supplied is only of the order of 50 kJ/m² an hour but it can also prevent the normal respiratory heat loss of around 35 kJ/m² an hour,¹³ which is probably the largest source of further heat loss to those with peripheral circulatory stasis and profound skin cooling. Preventing this loss and giving some added heat as a bonus is all that can be done for the victim on site apart from insulation, to which they are adjuncts and not substitutes. Perhaps "airway insulation" should be used instead of airway rewarming.

Other available warming techniques are chemical heat packs, used by the US military and European rescue teams, but these are not supported with precise details of effectiveness, and the Thermal Sargol,¹⁴ whereby heated ethylene glycol circulates through tubing in a sleeping bag. Both of these carry the hazard of dropping core temperature by opening up the flow of cold, pooled blood in the periphery if used indiscriminately.

I have just been given but not yet used a simple piece of equipment for airway insulation during evacuation. A light-weight face mask has a 6 in length of anaesthetic tubing with a cardboard air filter attached. The tubing is placed inside the victim's anorak or sleeping bag. He then breathes the air slightly warmed by his own body and with high humidity retained by the filter (ELI Lloyd, personal communication). Reducing respiratory heat loss is the sole aim.

A long rescue lasting several hours through hostile conditions may be inevitable for victims of mountain or down cave rescue where the victim cannot expect official hospital care. Most will have detectable ketones on the breath. We feed and offer warm drinks to the victims but the only acceptable source of energy is the victim's own fat stores. Such a measurement should not be done within half an hour of giving a warm drink.

I have not personally seen a victim in the state described as "warm euphoria" where cerebellar and judgment are impaired by cold so that clothing is discarded. Often those collapsed have been carrying

but have not donned insulating clothing or have ignored a nearby source of shelter. Scottish teams have described following a trail of cast-off clothing to find victims.¹⁵ A difficult decision with hypothermic victims is in determining whether life still exists. Nazi and Lewis¹⁶ described recovery after one hour of cardiac arrest. Golden¹⁷ has pointed out that many found apparently drowned wearing lifejackets are hypothermic and savable, and children submerged in cold water for up to 45 minutes have recovered without cerebral impairment.¹⁸ Within the constraints of common sense the diagnosis of death from hypothermia can only be that of failure to resuscitate.

In severe exposure hypothermia of young and fit people who were rapidly chilled and particularly who are deteriorating clinically, active rewarming as soon as is practical is desirable and lifesaving.¹⁹ If helicopter evacuation is not feasible then immersion in a warm bath at the nearest house may be necessary and facilities for this should be organised during, not after, rescue. Place the trunk but not the arms and legs in a bath of water at 37°C. If no bath is available and if no thermometer is available, a cold victim will rapidly cool the surrounding water which must be constantly stirred and its temperature measured by "topping up" so that around 40 gallons of water are necessary. Many victims show aggressive, uncontrollable movements after immersion so that the head must be kept as low as is safe and be firmly supported. Medical supervision is desirable during this treatment as hypotension, cardiac arrhythmias or arrest, and hypoglycaemia are theoretically possible.²⁰ The shortening of the "after drop" in core temperature is the immediate aim. General practitioners in mountainous and marine areas should be aware of the benefits of this technique.

Fortunately many of our victims out of doors are young and resilient, with a better prognosis than the cold elderly.

I thank Dr Evan L. Lloyd, South Lothian Health District, Edinburgh, for his helpful comments on this article.

References

- Lloyd DC. Cavers dying of cold. *British Med J* 1964;79:1-5.
- Maclean D, Emble-Smith D. *Accidental hypothermia*. Blackwell Scientific Publications, 1977.
- Day AE, Morgan EB. Accidental hypothermia—report of a case following alcohol and barbiturate overdose. *Ann Intern Med* 1974;73:72-7.
- Fuller RH, Gunning AJ, Bellamy KD, Trigg DR. Severe hypothermia as a result of barbiturate overdose complicated by cardiac arrest. *Lancet* 1969;ii:907-9.
- Alexander L. The treatment of shock from prolonged exposure to cold especially in water. *Report No 250*. Washington, DC: Office of the Assistant Surgeon General, Department of Commerce, 1945.
- Freeman J, Pugh LGCE. Hypothermia in mountain accidents. *Int J Aviat Space Med* 1967;38:253-6.
- Handbook of mountain and cave rescue 1978-80. In: Mountain Area, Temple Meads, Buxton, Derbyshire SK17 9BE: Mountain Rescue Committee, 1979.
- Keighley JH. A comparison of mountain rescue bags. *Gloving Research J* 1980;8:40-56.
- Frankland JC. Caving: problems and pleasures. *Practitioner* 1972;210:211-8.
- Combs GL, Johnson RE, Peoria JL. *Physiological measurements of metabolic functions in man*. New York: McGraw-Hill, 1963.
- Lloyd EL, et al. A comparison of the effectiveness of central rewarming as a first aid measure. *Scott Med J* 1972;17:83-91.
- Lloyd EL. *Handbook of mountain and cave rescue*. In: Mountain Area, Temple Meads, Buxton, Derbyshire SK17 9BE: Mountain Rescue Committee, 1979.
- Anonymous. Immersion and drowning in children. *Br Med J* 1977;iii:77.
- Auld CD, Light LM, Norman JN. Accidental hypothermia and rewarming in dogs. *Am J Vet Res* 1979;40:601-8.
- Collis ML, Stronman AM, Chaney RD. Accidental hypothermia: an experimental study of practical rewarming methods. *Aviation Space Environ Med* 1977;48:825-32.
- Lloyd EL. Accidental hypothermia treated by central rewarming through the armpits. *Br J Anaesth* 1977;48:41-5.
- Stewart T. Clinical signs of exposure to cold. *General Practitioner* 1977;23:23.
- Nazi SA, Lewis JF. Profound hypothermia in man—report of a case. *Am Surg* 1958;14:264-6.
- Golden FSC. Resuscitation and treatment of immersion hypothermia. *Proc R Soc Med* 1968;61:1058-61.
- Derbyshire DK, Clark RG. Cerebral recovery after prolonged global brain ischaemia. *Lancet* 1980;ii:831.
- Anonymous. Severe accidental hypothermia. *Lancet* 1972;ii:337.

*Mountain Instructors Survival Bag from Karrimor Products, Avenue Parade, Lancaster, Lancs.

†Made by Peter Bell Engineering, The Slack, Ambleside, Cumbria. Similar equipment is marketed under the name "Thermoguard" made by Electro Med & V Tech Service, 932 Equatorial Road, Victoria BC, Canada and as "The Little Dragon" made by M P Michels, Kendal, Cumbria.

‡Manufactured as "Warm-Rite" by Energy Systems, c/o Pine Street, Nashua, New Hampshire 03060 USA at about \$1800.

Pitfalls in Practice

Finding a practice

II: The interview

JOHN OLDROYD

This article is based on an oral presentation made for vocational trainees in general practice by the MSD Foundation.

Last week Dr Ian Keen, young and newly married, searched the advertisements in the *BMJ* for a place as a family practitioner on a committee list. He decided to apply for this one:

THIRD PARTNER to join two established principals in Elysium on Thames. The practice is conducted from owned premises in this market town which has excellent ranges of housing and good schools. Ancillary staff employed, attached health visitor and district nurse. Salary £8000 at start with review. Curriculum vitae please to Dr Sharp & Lide, Regency House, The Market Cross, Elysium on Thames.

Ian's curriculum vitae was one of many to arrive at Regency House, and Dr Jack Sharpe and Dr Alfred Lide considered that on paper he showed promise. The fact that Jack Sharpe, like Ian, was also an old boy of Workhouse and St Enoch's Hospital and thought Keen would be the right type for Elysium was, of course, an undisturbing factor. And so far as Lide was concerned, any doctor with a young and newly acquired wife was well worth meeting.

Dr and Mrs Keen were given a date and time to visit the practice. The interview progressed well, though Lide—illustrious Grammar and Redbrick—was decidedly cynical about the frequent references to the Master's house and the hospital facilities. Nevertheless Dr Lide assessed Keen as full of energy and enthusiasm for work and gave Sharpe the nod to drop the interview which gave him the go-ahead to offer to Keen the post of junior partner.

SHARPE Well, look here, Ian—I'm sure you don't mind me calling you Ian—you're just the chap we are looking for. I think both Alfred and I agree we'd like to invite you to join us as a partner.

KEEN Thank you very much, Dr Sharpe. I am sure it's what I am looking for. You have been very clear with your description of the practice.

LIDE Well, you've certainly asked a lot of searching questions, and I am pleased you like the way we have it organised. One thing we have to talk about is money. Jack and I have built up a practice we are proud of and as senior partners we are glad to have an enthusiastic like you joining us. But, of course, with your abilities you will quickly be in a good financial position.

SHARPE Yes, of course, we think our offer is fair. We want you to be a salaried partner, though I believe the fashionable term is "a partner receiving a guaranteed salary." There you'll know just what we mean. I wish I had known when I was setting up—so you won't have to

Secretariat for London Local Medical Committees, Tavistock House North, London WC1H 8ET

JOHN OLDROYD, MB, FRCP, secretary

worry about your income being dependent on profit or losses. We'll offer you £8000 a year for starters and, if all goes well, increase this by £500 a year to £9500. In addition to this we'll go further and promise that in any year that your salary doesn't come up to a third of Alfred's and my share we'll bring it up to that point.

KEEN That sounds about what is on offer in the *BMJ*, but some of the ads carried higher rates.

LIDE Ah, but did they say whether that included the expenses? We shall be paying all the practice expenses and, anyway, I bet some of them were in pretty grotty places.

KEEN Yes, that's true, but is there to be any increase to parity subsequently?

SHARPE Ian, none of us knows what things will be like in practice in three years. We'll all have to get together and think about that again. We shall give you three weeks' holiday a year and, if you want it, some study leave as well. You'll get a half-day a week, and there'll be some weekends off as well. We'll work out a rota, anyway. Of course, we don't want to see these deputising services—our patients deserve better than that.

KEEN Well, that sounds reasonable.

SHARPE Good. Then this is splendid. We can shake hands and you're starting on the first of the month after next, then? I must say the amiable way we've settled this is a good omen for the future. The word of an old Workhouse man is good enough for me and even Alfred won't object even though...

During this "all good fellows together" discussion, Ian Keen's mind, which doesn't often turn to legal matters, remembers he has received a written golden rule on his vocational day-release course about partnership agreements. One pops into his head now:

All potential future partners are not necessarily gentlemen and honourable

While discussions are going on all appear to be so, but it may take some months for their true natures to emerge. Acronym creep into many partnerships: if the partnership is merely one of will it offers no security to anyone. Where the conditions of the agreement are written down in an agreement, then, when disputes occur, at least there is a definite basis on which to resolve them. Dr Keen should therefore insist on the partnership being based on a drawn-up partnership agreement so that he can study the details.

KEEN Yes, but I'm sure you must feel as I do that it's wise for us to have a partnership agreement in writing so we can get down on paper details about my income and my duties in the practice and something spelling out all our responsibilities.

While saying this in a somewhat embarrassed fashion—after all Sharpe and Lide, who seem decent fellows, are looking rather pained—Lide also recalls:

A medical partnership is not only an association between professional colleagues

... it is also a business arrangement. What's more, whether or not there is a written agreement, if the conduct of the parties makes it

appear that they are acting in partnership in law states that they are partners to the outside world. Partnership is a legal subject that much of our clothing is made of. The fact that Jack Sharpe, like Ian, was also an old boy of Workhouse and St Enoch's Hospital and thought Keen would be the right type for Elysium was, of course, an undisturbing factor. And so far as Lide was concerned, any doctor with a young and newly acquired wife was well worth meeting.

Prompted by Keen's wish to get things down in writing, Lide gives Sharpe a quarter of an hour to get the agreement in writing. Sharpe, like Jack Sharpe, learns as chairman of all those committees, doesn't desert him at this stage.

SHARPE Quite right, Ian. If you feel it necessary, you say so. Alfred and I can, I am sure, rely on you. If you feel you happier we can go along with you. Let's write it out here and now. You can spare another 20 minutes, I suppose—the train to catch when you have your car outside. I suppose you'll have to take to your rounds in, eh?

Sharpe glances out of the window at Ian's beloved, but now aging, Capri. An amalgam of his 21st birthday presents and life savings. However, Ian has also heard that:

Doctors are rotten businessmen and abysmally ignorant of partnership law

... consequently it's worth paying a small fee to draw up an agreement now rather than pay the "princes" of the legal profession enormous fees when disputes occur later.

KEEN Don't you think it would be better if perhaps we had a solicitor draw it up? I am a bit of a novice when it comes to legal matters.

Clinical Curios

We have used ultrasound to treat nine patients for pain in the early and late stages of herpes zoster and have achieved sufficiently dramatic results to report on this. We know of others who have had success with this technique, but know of no published report.

We apply ultrasound over the painful area and not necessarily over the nerve root in a dose of two minutes of 1 MHz of 0.25 W/cm² for each 12 cm² (2 square inches) using liquid paraffin as a couplant. For a large area we have used the same dose per area. Treatment has been given daily except at weekends. There has been only one side effect when a second treatment was given only 17 hours after the previous treatment—"a drawing feeling," which disappeared within a day. Tolerance to treatment of the most we have used.

A 58-year-old woman with herpes zoster of the right infraorbital nerve needed analgesics day and night. After the first treatment with ultrasound the pain was free of pain for two months. He held his first night after nine treatments he was free of pain. He reports that particularly on cold days he has discomfort and needs some analgesics.

A 50-year-old woman became depressed and unable to face doing housework four weeks after pain from herpes zoster had started despite the pain being dulled by analgesics. After the first ultrasound treatment she was free of pain for the first time. Soon she was doing housework, and after 12 treatments she was completely free of pain.

An 87-year-old woman had had pain for two years after an infection with herpes zoster. She had not had one night's rest without drugs. After the first treatment with ultrasound the pain was free of pain after six treatments she was so relieved that she now needs only one-quarter of the amount of analgesics to relieve the residual discomfort.

A 90-year-old woman with a large area of buttock and groin affected by herpes zoster started to develop pain three days after onset of the infection. Ultrasound treatment was given at the active blistering stage and the pain stopped. After three treatments she said she did not need any more and does not need analgesics.

A 45-year-old male had a small area of herpes zoster growing near his eye. When the pain started he was given analgesics, but after the first and only treatment with ultrasound they were not needed.

Ide is looking singularly uncomfortable and is clearly beginning to regret his previous compliance, but Jack is on familiar ground.

SHARPE We seem to have got ourselves a very young man as well as a good doctor, Alfred. Look Ian, I think you're right. I'll be seeing Mr Bonamie tomorrow. He's our local, have what local "regal" and I'll get him to draft it out and send it in the post. That should suit you, eh? I am sure there won't be any problems. As soon as it is all signed, sealed, and delivered...

There are smiles all round. Sharpe and Lide can rely on Bonamie, their long-standing friend and associate, to produce what they want. Ian is happy. He hasn't ruined his chances by being punctilious, but he has forgotten another golden rule:

The lawyer drawing up an agreement will not necessarily look after the interests of all parties

He will do his best, but the basis of the agreement is what the parties instruct him to put in under the various headings he knows are necessary. Also, because few lawyers understand the way in which GPs are remunerated, it's as well to ask around for firms who have experience in this field. But each of the partners should ask his own and separate adviser to scan this legally. Ian should obtain an independent opinion on the legalities and also ask a colleague with experience, perhaps the secretary of the local medical committee, to advise on the contract for him personally and an accountant to advise on the financial and tax implications.

They part amicably with handshakes and congratulations all round. This is the second of four articles on finding a practice.

The vesicles continued to increase as expected, but the pain has been minimal.

In every case there was immediate relief after the first treatment. Some pain usually occurred after a few hours but progressively lessened with further treatment. Idoxuridine 5% (Herpid) had been used in all cases before the pain started but had not prevented pain. We think the ultrasound should be further studied for treating patients with herpes zoster, and we have the impression after treating four early cases, two of which are included here, that it is even more effective for the early cases than the late cases—ARTHUR GABBERT, general practitioner, MARY GABBERT, physiotherapist, Reapham, Norfolk.

A man of 65, who seldom complained of anything, came to my surgery one summer saying that he had a boil. It was much bigger than any boil he had before and was giving him a lot of pain and disturbing his sleep. He did, indeed, have what looked like a large boil on the right lower part of his anterior abdominal wall. A course of antibiotics and kaoline poultices seemed to produce some localisation and the discharge of some pus, but the lesion persisted. As there didn't appear to be a very active tissue reaction round the "boil," only the local treatment was maintained, and the lesion discharged pus intermittently but did not resolve.

This lack of progress led me to reconsider the diagnosis, but there was no history of any injury, either internal or external, and all his other bodily systems appeared to be behaving normally. I therefore referred him to a friendly surgeon, who found in addition to a large quantity of pus a fishbone some 4 cm in length. The abscess then healed promptly without any trace of a fistula from the bowel developing and the patient swore that he hardly ever ate fish—MIKE COURTNEY, general practitioner, Battersea.

We will be pleased to consider for publication other interesting clinical observations made in general practice.—ED, BMJ.

Correction

Beyond the Surgery: General practitioner as coroner

In this article by Sir Montague B Levine (17 January, p 166) the final sentence in the third paragraph should have read: "It is an excellent course and Britain is unique in providing this special qualification."