CORRESPONDENCE

Colposcopy and cytology Jocelyn Imrie, MRCPATH; B A Evans, FRCP ...	1547	Clinical Algorithms: Infertility M Elstein, FRCOG, and others	1550	Subcutaneous calcium heparin versus intravenous sodium heparin in treatment of established acute deep vein thrombosis of the legs N Scolding, MRCP; M G Walker, FRCS, and others.....	1551
Early onset pre-eclampsia: recognition of underlying renal disease M D Lindheimer, FACP, and L C Chesley, MD	1547	Irritable bowel syndrome as a cause of chronic pain in women attending a gynaecological clinic P Hogston, FRCS	1550	Medical staffing and training in the West Midlands region J Parkhouse, FFARCS, and others.....	1552
Hospices for children? D Stevens, MRCP, and Marie Owen, DCH; Valerie Broadbent, MRCS	1548	Unemployment and mortality G Costa, MD, and N Segnan, MD.....	1550	Points Explanation and management of neurological disability (N Gordon); Torsion of the testis: why is the prognosis so poor? (E M Kiely); Osteoporosis: cause and management (B McConkey); Inappropriate use of confidence intervals (P MacFarlane)....	1552
Atlantoaxial instability in Down's syndrome Y T Kontinen, MD, and S S Santavirta, MD; B Jagiivan, MB, and others; I Calder, FFARCS	1549	What are health authorities doing about the health problems caused by unemployment? J Middleton, MB, and others.....	1551		
Asthma and bronchial reactivity J B L Howell, FRCP, and W E Waters, FFCM.....	1549	Promoting prevention in primary care Ann-Louise Kinmonth, MD, and others	1551		

- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.
- Because we receive many more letters than we can publish we may shorten those we do print, particularly when we receive several on the same subject.

Colposcopy and cytology

SIR,—The edition of 23 May highlighted the current dilemma in colposcopy and cytology with four articles approaching the subject from different angles: Professor H Fox's leading article (p 1307), the article by Dr H C Kitchener and others (p 1313), Dr Jane Chomet's Practice Research article (p 1326), and the Lesson of the Week by Dr M J Campion and colleagues (p 1337). As a cytologist, I would endorse Professor Fox's opinion that standardisation of terminology in cytology is important, but this would do little to alter the basic problem, which is the borderline ("atypical") smear in patients with preneoplastic lesions (cervical intraepithelial neoplasia grade 2 or 3).

Few would disagree that colposcopy should be more widely available, but surely no one can envisage its use as a screening test. The provision of training for new colposcopists, whether in general practice, family planning clinics, or genitourinary medicine clinics, will, at least initially, increase demands on colposcopy centres. In addition, the spin off effects of expansion will increase the workload of cytology and histopathology laboratories as well as that of primary care doctors responsible for follow up.

A cytological report recommends a course of action in local conditions. If the colposcopy clinic has a four month waiting list (which may be undesirable but is not unknown) or there is no local colposcopy service a repeat smear may be a more reasonable course of action than referral in the first instance for borderline changes.

The present state of disarray can be resolved only by closer links between colposcopy and cytology, with the colposcopist indicating to the cytologist what advice he considers to be most appropriate (partly depending on what numbers the colposcopy service can cope with). Various

other factors, including correlation of cytological and histological findings, are necessary to determine the optimum advice so that patients may not develop invasion lesions. In view of the apparent trend towards "false negative" cytology — that is, borderline changes — in cervical intraepithelial neoplasia grade 2 or 3 such advice cannot be a static decision but must constantly be reviewed.

JOCELYN IMRIE

Monklands District General Hospital,
Airdrie ML6 0JS

SIR,—Dr Jane Chomet and Dr H C Kitchener and colleagues each report a year's experience of colposcopy in general practice and family planning clinic settings, respectively. In the same issue Dr M J Campion and colleagues emphasise the increased risk of invasive cervical cancer associated with human papillomavirus infection and suggest that such disease may be the earliest stage of cervical intraepithelial neoplasia.

Genital wart virus infection is the third commonest sexually transmitted disease in the United Kingdom after non-specific genital infection and candidal infection, yet in neither of these colposcopic studies did the authors assess the history or prevalence of wart virus infection. If this aspect of cytological changes in the cervix is not appreciated and appropriate action taken the burgeoning threat of cervical cancer will continue to grow.

The increasing prevalence of precancer in young women is faced principally by departments of genitourinary medicine, where training in colposcopy has been an integral part of higher specialist training for the past two years, and clinics have been operating in several centres for much longer. Our waiting list for colposcopy at the West London

Hospital, where we run three clinics a week, is now six months. More resources are needed urgently.

B A EVANS

West London Hospital,
London W6 7DQ

Early onset pre-eclampsia: recognition of underlying renal disease

SIR,—Dr B U Ihle and coworkers (10 January, p 79), described 84 women with pre-eclampsia before 37 weeks' gestation, noting a high degree of recurrent hypertension in subsequent pregnancies and, surprisingly, the presence of chronic renal disease or essential hypertension in 90% of the subjects. In two thirds of these patients the lesion was glomerulonephritis, usually of the IgA variety. We wonder whether these data represent a selected population, perhaps relating to the referral pattern of the University of Melbourne renal department.

One of us (LCC) followed the clinical course of 270 women surviving eclampsia (the convulsive phase of the disorder) between 1931 and 1951, and most of these women were re-examined periodically for almost 40 years.¹² Unfortunately, the prenatal records did not always reflect when the disease started, but we ascertained that it was before 36 weeks' gestation in at least 42% of the women. (As eclampsia represents a severe stage of the disorder it is reasonable to assume that many more had had hypertension for some weeks before the putative convulsion.)

When urine samples from 174 of these women, still alive 23-43 years after the eclamptic gestation, were examined for protein only four yielded positive results. Of the 96 who died, 17 had a

history of proteinuria. These data confirmed an earlier follow up study published in 1941, in which measurement of specific gravity, proteinuria, and urea clearances was used to survey 165 women who had had eclampsia: 94% had normal renal function.³ Assuming that only a third of all the above cases (a probable underestimation) had early pre-eclampsia, we would have expected more evidence of compromised renal function if the incidence of underlying parenchymal kidney lesions were as high as suggested by Dr Ihle and colleagues.

FIG 1—Distribution of birth weight of 157 infants in relation to gestational age. Different diseases identified by renal biopsy examination are illustrated. The 10th, 50th, and 90th centile growth curve lines were adopted from Lin C-C *et al.*,⁵ *Am J Obstet Gynecol*, by permission.

FIG 2—Birth weight in relation to gestational age for all live infants. Lines represent mean weight (1 and 2 SD) of 25 608 live births at university hospitals of Cleveland between 1958 and 1968. From Katz AI *et al.*,⁶ *Kidney Int*, by permission.

The second of us (MDL) reported on the gestational course and remote prognosis of over 150 women in whom the aetiology of their hypertensive complications of pregnancy was determined by postpartum renal biopsies.^{4,5} Fig 1 shows fetal weight and outcome in relation to the biopsy diagnoses. Most of the women who delivered before 37 weeks' gestation had the renal lesion of pre-eclampsia alone. When women with known renal diseases were studied (fig 2) most delivered after the 37th week. Choice of who should undergo biopsy, even in centres with liberal criteria, is usually quite selective, often based on an unusual facet of the case; thus the degree of underlying renal disease in our patients with pre-eclampsia is probably overestimated.

Finally, Dr Ihle and coworkers are to be congratulated as they have elegantly confirmed the dictum that women with early and severe pre-eclampsia represent those most likely to harbour other diseases (usually essential hypertension later in life.) Furthermore, this is the group most likely to have recurrent hypertension in subsequent gestations (the severity of which is not predictable). We disagree, however, with the contention that 90% will have another disorder, believing that in fact most will still have pure pre-eclampsia and that most of the subsequent pregnancies will be quite successful.

MARSHALL D LINDHEIMER

University of Chicago

LEON C CHESLEY

Downstate Medical Center,
University of the State of New York

- Chesley LC, Anntito JE, Cosgrove RA. The remote prognosis of eclamptic women: sixth periodic report. *Am J Obstet Gynecol* 1976;124:446-59.
- Chesley LC. *Hypertensive disorders in pregnancy*. New York: Appleton-Century-Crofts, 1978.
- Chesley LC, Somers WH. Eclampsia and post eclamptic hypertension: a follow-up study with an analysis of factors affecting the remote prognosis. *Surg Gynecol Obstet* 1941;72:872-88.
- Fisher KA, Luger A, Spargo BH, Lindheimer MD. Hypertension in pregnancy: clinical pathological correlations and remote prognosis. *Medicine* 1981;60:267-76.
- Lin C-C, Lindheimer MD, River P, Moawad AH. Fetal outcome in hypertensive disorders of pregnancy. *Am J Obstet Gynecol* 1982;142:255-60.
- Katz AI, Davison JM, Hayslett JP, Singson E, Lindheimer MD. Pregnancy in women with kidney disease. *Kidney Int* 1980;18:192-206.

Hospices for children?

SIR,—The excellent leading article by Dr T L Chambers (23 May, p 1309) emphasised the desirability of caring for terminally ill children at home. In Gloucestershire over the past 10 years nearly all children dying of cancer and leukaemia have been nursed at home by their parents supported by the paediatric team who were responsible for the child's treatment. The paediatric team of consultant, clinical assistant, day care nurse, and, more recently, domiciliary paediatric oncology nurse work with the primary care team to provide a 24 hour service for advice and domiciliary care. The parents always need a great deal of support and may need a member of the team to be present at the moment of death. For the small group of parents who wish the child to be nursed in the hospital we have recently furnished a family flat near to the children's wards, where they can look after their child with the close support of the hospital staff.

Increasing numbers of children with life threatening illnesses are receiving all their medical care in regional centres, which may be many miles from home. Treatment often lasts for several years, and the parents and children become completely dependent on the regional centre for medical care and emotional support, during which time the

district paediatrician and family doctor become strangers to the child, the family, and the illness. When the child becomes terminally ill it is very difficult for the regional centre to provide the necessary support for parents to look after their dying child many miles away, yet it is only with such support that many parents will feel able to look after their child at home.

In Gloucestershire we have worked with the regional paediatric oncology centre at Bristol to provide a system of shared care that seems to have overcome these problems while maintaining the undoubted benefits of centralised care. All children with cancer and leukaemia are under the shared care of a regional paediatric oncologist and the district paediatric oncology team led by one consultant and clinical assistant. Much of the treatment is given locally, yet the regional specialist sees the children in monthly clinics, makes all strategic decisions, and is always available for counselling and advice. The district team is experienced in delivering treatment locally with minimal disruption to family life and schooling and in the event of terminal illness gives the domiciliary care and support to enable the child to be nursed at home.

DAVID STEVENS
MARIE OWEN

Gloucestershire Royal Hospital,
Gloucester GL1 3NN

SIR,—I wholeheartedly agree with Dr T L Chambers's view that institutional arrangements for the care of dying children are second best to home care (23 May, p 1309).

Dying children differ from dying adults in that they are more portable and thus more easily nursed, have (usually) two able-bodied parents to care for them, and have fairly brief periods of incapacitating terminal illness, often attending school up to a few days before death. On these grounds alone the necessity for institutional care is less than that in adults.

Most families are frightened and feel unable to cope when faced with the prospect of their child dying at home. Time for honest discussion of fears and to dispel myths, however, invariably results in a very satisfying and even joyful experience. This can be achieved only by sound teamwork and liaison among all health care workers.

Many primary care teams have little experience of caring for a dying child as death in childhood is fortunately rare. Hospital based community nurses are invaluable in providing continuity of care between hospital and home and updating the home care team in the use of modern technology (syringe drivers and central venous lines, for instance) as well as in pain control. Many nurses provide this and other services, such as babysitting and night care, on a purely voluntary and unofficial basis. Such visionary and ideal practices are often hampered, primarily by lack of funds but also by bureaucratic rigidity.

If institutional care is necessary most children (and families) would choose a familiar hospital ward. In this case even a busy acute ward can be made to feel as much like home as possible with thought and imagination. This sometimes requires the devious practice of ensuring that the nursing hierarchy are absent when a beloved bouncy golden labrador pet is brought in or enabling secret cooking sessions in the ward microwave oven for a midnight feast.

Mother Frances has said that hospice care is a philosophy, not an institution. It is a pity that the general public prefers to make its emotional response to dying children by putting money into bricks and mortar rather than supporting the